University of Reading

On thinning ice: effects of atmospheric warming, changes in wind speed and rainfall on ice conditions in temperate lakes (Northern Poland)

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Bartosiewicz, M. ORCID: https://orcid.org/0000-0001-5197-1630, Ptak, M., Woolway, R. I. ORCID: https://orcid.org/0000-0003-0498-7968 and Sojka, M. (2021) On thinning ice: effects of atmospheric warming, changes in wind speed and rainfall on ice conditions in temperate lakes (Northern Poland). Journal of Hydrology, 597. 125724. ISSN 0022-1694 doi: 10.1016/j.jhydrol.2020.125724 Available at https://centaur.reading.ac.uk/100092/

It is advisable to refer to the publisher's version if you intend to cite from the work. See Guidance on citing.

To link to this article DOI: http://dx.doi.org/10.1016/j.jhydrol.2020.125724

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

On thinning ice: Effects of atmospheric warming, changes in wind speed and rainfall on ice conditions in temperate lakes (Northern Poland)

Maciej Bartosiewicz[1*], Mariusz Ptak[2], R. Iestyn Woolway[3] and Mariusz Sojka[4]

[1-] Department of Environmental Sciences, University of Basel, Basel, Switzerland

[2-] Department of Hydrology and Water Management, Adam Mickiewicz University, Krygowskiego 10, 61-680 Poznań, Poland

[3-] European Space Agency Climate Office, ECSAT, Harwell Campus, Didcot, Oxfordshire, UK

[4-] Institute of Land Improvement, Environmental Development and Geodesy, Poznań Life Science University, Piątkowska 94, Poznań, Poland

*Corresponding Author: maciej.bartosiewicz@unibas.ch
Department of Environmental Sciences,
University of Basel, Basel, Switzerland

Key words ice thickness, ice duration, winter limnology, seasonal effects, climate change

Abstract

Northern Hemisphere lakes are losing their ice cover due to climate change. Here we explored six decades of observational data (1961-2017) showing trends in air temperature, wind speed and precipitation over northern Poland, as well as changes in the ice conditions for five lakes with different morphometry. We evaluated whether and to what extent climatic effects, including atmospheric warming, changing wind speed and precipitation during fall and winter, influence ice conditions in morphometrically different lakes in Northern Poland. Our analysis demonstrated that ice cover duration and thickness in decreased at rates of 5.4 days decade^{-1} and 2.5 cm decade^{-1}, respectively. Ice conditions were influenced (65-75%) by the direct effects of air temperature change and to some extent by an interaction of warming with wind speeds and rainfall (5-10%). While stronger autumnal winds result in longer ice cover duration, the effect of precipitation is bimodal with either an enhancement of ice formation by autumnal rain or accelerated ice loss during spring. To project future changes in ice conditions, we used a 1D hydrodynamic lake model forced with four climate model projections under low, medium and high Representative Concentration Pathway (RCP) scenarios. Our simulations demonstrate that current ice conditions will stabilize under the low emission scenario (RCP 2.6) but decrease under both the medium and high emission scenarios (RCP 6.0 and 8.5). During the 21st century, the studied lakes are projected to lose their ice at a rate between 4.5 and 10 days decade^{-1} and ice thickness will decrease by between 3.0 and 5.0 cm decade^{-1}. The rate of change will be more rapid in smaller rather than larger lakes and more so for those situated further inland. The probability of ice-free winters will increase for all lakes and among all future scenarios by between 4 and 69% with the highest potential frequency of ice-free winters in smaller and deeper but relatively wind-exposed lakes.

1. Introduction

Under the effects of global climate change, surface waters are currently warming at an unprecedented rate (Czernecki and Ptak, 2018; Woolway et al., 2019), and consequently lakes are losing their ice cover (Kainz et al., 2017; Ptak et al., 2018). Frozen lakes function differently than those without ice as turbulent mixing and light-dependent processes under the ice are greatly reduced (Ptak et al., 2019a). Ice and snow also reduce the interactions between the lake and the atmosphere; thus, hydrodynamic conditions are controlled by the solar radiation penetrating through the ice (Mironov et al., 2002), heat flows from sediments (Terzhevik et al., 2009) and lateral inflows (Bengtsson 1986; 1996). As the under-ice temperature at the water-ice interface is fixed at the freezing point the stratification and mixing patterns are controlled by convective motions and conductivity gradients (Kirillin et al., 2012; Bouffard et al., 2019). Thickness and duration of the ice cover have thus consequences not only for the hydrodynamics and biogeochemistry of lakes (Williams et al., 2004) but also for their oxygenation and plankton communities (Adrian et al., 1999; Bartosiewicz et al., 2019a). These effects, most pronounced in winter, have important implications for the functioning of lakes in the warm productive season (Hampton et al., 2017).

Ice conditions can influence the phenology and strength of thermal stratification (Preston et al., 2016), water retention times and associated changes in the water chemistry, the extent and composition of phytoplankton blooms (Adrian et al., 1999) and related shifts in the abundance of zooplankton and fishes (Helland et al., 2011). The cascading effects of changes in the ice cover duration on biological productivity are also reflected in its relevance for the economic (Prowse and Brown 2010) and socioeconomic importance of lakes (Orru et al., 2014). In this context it is important to underline that while most lakes in the northern hemisphere are gradually losing their ice, many that were regularly frozen in the past already remain now completely ice free during

exceptionally warm winters (Sharma et al., 2019). The process and consequences of decreasing ice cover in lakes have been reported previously (Magnuson et al., 2000; Knoll et al., 2019). However, the processes responsible for the increased probability of complete ice loss in the near future (Benson et al., 2012; Sharma et al., 2016; Wu et al., 2018), which may have irreversible effect on, among other things, planktonic food webs as well as organic matter processing in lakes, are not well documented.

Changes in lake ice phenology are known to be influenced by air temperatures and ambient wind speeds (Arp et al., 2013), but also depend on lake size and shape in relation to prevalent wind directions (i.e., effective fetch, Magee and Wu, 2017). On the one hand, while changes in zonal winds related to large-scale climate oscillations drive the interannual variability in ice-off dates (Schmidt et al., 2019), changes in wind speed above individual lakes, as well as associated patterns of mixing, may drive regional differences in timing of lake ice formation. In addition, while rainfall in spring may accelerate ice cover thinning until breakup, rainfall in autumn also influence ice conditions (Leppäranta, 2010) by altering mixing intensity and temperatures in the water column (Rooney et al., 2018). Despite the potentially important interactions between warming temperatures and other components of recent climate change, such as atmospheric stilling (Woolway et al., 2019) and changes in precipitation (Caine, 2002), the mechanisms behind the interactive control of hydroclimate on lake ice remains largely unexplored. Localized effects such as distance from the coastline or regional weather patterns may control the responsiveness of the ice cover to interannual changes in atmospheric temperature and circulation (Weyhenmeyer et al., 2004). Notwithstanding these indirect effects, predictions show that temperate lakes will have less and thinner ice in the near future and that the rate of this decline may depend on the interaction between limnological characteristics and warming effects (Yao et al., 2014; Tan et al., 2018).

In this study we explored a rich meteorological dataset, including records of air temperature, wind speed and precipitation, collected between 1961 and 2017 in the vicinity of five temperate lakes in Northern Poland, where associated changes in ice cover phenology and thickness were available. We investigated the interaction between the effects of air temperature, rainfall and wind speed on ice phenology and thickness, as well as evaluated how the interactions between changes in these atmospheric drivers influence ice cover in the morphometrically different lakes. We also used a 1D hydrodynamic model forced by an ensemble of four climate model projections under low, medium and high Representative Concentration Pathway (RCP) scenarios to simulate changes in the ice conditions for the studied until the end of the 21st century.

2. Methods

2.1. Lakes

We studied five temperate lakes located in northern Poland (Fig. 2). Study lakes represent a gradient of depth, mixing regimes (Table 1), water transparency and productivity (as Secchi depth between 0.7 and 2.7m). Lakes also differ in morphometry from symmetric round to complex shapes with multiple sub-basins and these differences are reflected by wind exposition with fetch ranging from 3 to 13km.

2.2. Ice conditions and meteorology

The paper is based on data collected within a monitoring program at the Institute of Meteorology and Water Management – National Research Institute (IMiGW-PIB, Poland). Within this long-term monitoring program, the observations of ice conditions are conducted daily from the moment of ice formation to its disappearance (in the case of the analyzed lakes from November

131 to April). Monitored parameters cover, among others, the term of formation of the ice cover, term
132 of its breakup, and measurements of its thickness performed every fifth day at one sampling point
133 located near the water gauge. Meteorological data were within the same long-term monitoring
134 collected by the IMiGW-PIB (Poland) in the scope of standard monitoring for a network of
135 stations throughout the country. Briefly, each IMGW meteorological station is located on a flat
136 surface at least 100 m away from any group of buildings or tree stands and at least 100m from
137 any open water surfaces (i.e., lakes, rivers, reservoirs). Each meteorological station is equipped
138 with semi-automated (1965-1996) or fully automated (1996-2015) temperature logger positioned
139 2m above the ground and sheltered from the direct sunlight. Surface air temperatures in the
140 historical period (prior 1990) were recorded with traditional manual methods (at least every 8h).
141 More recently stations were equipped with automated temperature sensors and suitable data
142 loggers (i.e., LB-710R thermo-hygrometer with LB-480 data logging module). Each
143 meteorological station is equipped with Hellmann Rain Gage that allows to monitor daily
144 rainfalls and with anemorumbometer (i.e., M63 M1) allowing to continuously monitor speed and
145 direction of winds. Continuous, long term meteorological records were not always available from
146 the stations located in the immediate proximity of the studied lakes; hence for some lakes we
147 analyzed data from stations further afield (up to 20km for Lake C). However, the monthly
148 average observations used for the studied lakes, all of which are located in the lowland area of
149 Northern Poland, are representative of long term climatic effects in the region. Trends and effects
150 between meteorology and ice conditions for the studied lakes were analyzed by correlation
151 analysis (Pearson's) and multiple regression models (XLStat 2019). To remove any potential
152 autocorrelation between the predictor variables within the regression models (i.e., unconclusive
153 Durbin Watson test - DW), we used a Cochrane-Orcutt procedure ($1.95 > DW < 2.1$). Following
154 this procedure, all residuals from the regression models were normally distributed; thus all of the
155 model assumptions were fulfilled.

156

157 *2.3. Simulating future ice conditions*

158 To simulate historic and future (2020 to 2099) ice conditions (i.e., duration and thickness) in the
159 studied lakes, we used the 1D hydrodynamic Freshwater Lake model, FLake (Mironov, 2008;
160 Mironov et al., 2010), which has been tested extensively in past studies (Woolway and Merchant,
161 2019). In brief, FLake is process-based model which solves the heat budget of lakes at a daily
162 resolution. The integrated approach implemented in FLake allows a realistic representation of
163 the major physics behind turbulent and diffusive heat exchange in lakes; it includes an ice
164 module, and a module to describe the vertical temperature structure of the thermally active layer
165 of bottom sediments, as well as its interaction with the water column above. Noticeably, Flake is
166 is one of the most commonly used models in lake studies and is also used as a module in
167 numerical weather prediction (i.e., Rooney and Jones, 2010). The meteorological variables
168 required to drive FLake are air temperature at 2 m, wind speed at 10 m, surface solar and thermal
169 radiation, atmospheric pressure, and specific humidity. These atmospheric drivers were extracted
170 for this study from four bias-corrected (to the EWEMBI reference dataset; Frieler et al., 2017;
171 Lange, 2019) climate model projections from the Inter-Sectoral Impact Model Inter-comparison
172 Project phase 2b (ISIMIP2b), HadGEM2-ES, GFDL-ESM2M, IPSL-CM5A-LR, and MIROC5
173 (see Supplementary Information S1). Future projections, which represent the evolution of the
174 climate system subject to three different anthropogenic greenhouse gas emission scenarios
175 covering the period 2020 to 2099, RCP 2.6 (low-emission scenario), 6.0 (medium-emission), and
176 8.5 (high-emission), are also investigated. These data were available at a daily time step and at a
177 grid resolution of 0.5°. Time series data were extracted for the grid point situated closest to the
178 center of each studied lake. Aside from meteorological data (see Supplementary Information
179 File), FLake requires estimates of water transparency, lake depth and fetch. Given that Flake
180 equations account for the influence of length and depth of the lake basin on temperature and ice

conditions we assume that simulations, to some extent, account for the effects associated to differences in the size and shape of simulated lake basins.

Historical meteorological data (1965-2005) generated by the four climate models described above showed comparable rate of warming as observed over northern Poland during this study (Fig. S1). Future simulations showed either stabilization of air temperatures (RCP 2.6) or warming throughout the 21st century (RCP 6.0 and 8.5, Meinshausen et al., 2011; respectively). Comparison between Flake-simulated ice conditions (1965 to 2005) to future (2020-2099) predictions allowed us to also to estimate the relative increase in the probability of the ice-free winters for each of the monitored lake. This was done by assessing the frequency (in %) of events when the potential changes in the ice over duration or thickness may reach zero calculated as:

$$F_{\text{ice thick}} = \left\{ \frac{n|\overline{Ice} - SD(\overline{Ice}) \leq 0|}{N} \right\} \times 100$$

where $\overline{Ice}$ is the expected (predicted average) ice thickness, SD is the standard deviation from the four climate models and under one of the three greenhouse gas emission scenarios, n gives the number of observation when annual ice thickness was reaching zero within the range of one standard deviation and N is the total number of observations.

3. Results

3.1. Changes in regional and local meteorological conditions (1961-2017)

Observed climate change in northern Poland (Kolendowicz et al., 2019; Tomczyk and Szyga-Pluta, 2019) is consistent with trends observed elsewhere, including positive trends in air temperature, a variable course of rainfall and a negative trend in wind speeds between fall and winter (i.e., the period between ice formation and ice break-up, Fig. 3F). However, on a local

scale the magnitude (and to some extent even the direction) of these changes differ. For example, while warming over the last sixty years was significant for all weather stations, its magnitude ranged between 0.03 and 0.04°C y^{-1} (Fig. 3A and B, respectively). Similar variability was observed for trends in wind speed during fall and winter, which ranged between an increase of 0.009 ms^{-1} y^{-1} (Fig. 3A) and stilling between -0.001 and -0.02 ms^{-1} y^{-1} (Fig. 3B and E, respectively). Fall and wintertime rainfall increased throughout the region at rates between 0.07- and 0.2-mm y^{-1} (Fig. 3 E and B) and only decreased locally closer to the coast (Fig. 3A).

3.2. Changes in lake ice conditions and relationship to climate

All lakes considered in this study are losing their ice cover (Fig. 4A to F). Over the last sixty years, ice cover has formed later (on average by 1.4-day $decade^{-1}$) and disappeared faster (by 4-day $decade^{-1}$), which results in a total decrease of the duration of ice cover by 5.4 days $decade^{-1}$. On a more resolved spatial scale this decrease ranged between 3.9- and 7.8-day $decade^{-1}$ (Fig. 4. B and E). The ice cover for all lakes is also becoming thinner by 2.5 cm $decade^{-1}$ with rates of thinning ranging between 1.6- and 3.3-cm $decade^{-1}$ (Fig. 4B and D).

The multiple regression analysis revealed that ice conditions can be well simulated (predicted) for differently shaped temperate lakes (Fig. 5A to E) using a combination widely available meteorological data (temperature, wind, rainfall). The decrease in ice cover duration for all lakes is mostly influenced, and can be well predicted, by the effects of atmospheric warming (Fig. 5F and Table 2) but also by the effects of wind (September) and rainfall in autumn (November) and early spring (March). For all lakes, stronger winds in early autumn (i.e., October) had a negative influence on ice duration and thickness. However, stronger winds later in the season (i.e., November) resulted in the formation of thicker ice cover. Throughout the entire dataset the effect of rainfall was bimodal, with November rainfall resulting in an earlier

formation of ice cover and rainfall in March stimulating ice loss. Generally, between individual lakes the thickness of ice cover for temperate lakes was influenced by the influence of changing temperature ($R^2 > 0.65$) with wind speed and rainfall contributing an additional 5-10% to variability explained by the model (Fig. 5; Table 2). On a seasonal timescale, and for individual lakes, the influence of different components of climate change varied. For example, while warming had a persistent negative effect on both ice cover duration and thickness, the effects of wind and rainfall were bimodal, having either a positive (autumn) or a negative (autumn and spring) influence on the ice cover duration and thickness (Table 2). The goodness of fit for the regression model also varied between individual lakes (Table 3). For example, while all the F values were significant, they ranged from 7.6 to 17.8 for regressions of ice cover thickness and from 6.7 to 15.2 for these on ice cover duration (Lake C & A, respectively). Similarly, the square root of the variance of residuals (RMSE) varied between 4.8 and 7.7 for thickness and between 13.4 and 19.3 for duration. The mean absolute percentage error (MAPE) varied from 13 to 39.5 (Lake E & B) for ice thickness and from 14.1 to 30.8 for ice cover duration (Lake E & D, respectively).

Future climate and ice conditions in morphometrically different lakes (2020-2100)

The performance of the FLake model, which was validated using historical ice cover observations between 1961 and 2005, was found to be moderately good (Ice Thickness $0.12<R^2<0.24$; $p<0.001$, Ice duration $0.14<R^2<0.18$, $p<0.001$, Fig. S2), with the range and rate of change in ice conditions over the observational record both well reflected by the simulations (Fig. S2). The FLake future simulations demonstrated that ice cover is likely to remain relatively stable under RCP 2.6, with potentially marginal increase in the average thickness (Fig. 6) and duration (Fig. 7) of ice cover by 0.3 to 1-cm decade^{-1} and 0.6 to 1.7-day decade^{-1}, respectively. By contrast, under RCP 6.0 and 8.5, ice cover thickness and duration are projected to decrease in all studied lakes. The rate of change in ice conditions vary between individual lakes as well as

the RCP scenarios. That is, for the simulated future changes in ice cover thickness, the most pronounced decrease of 5 cm decade^{-1} was observed in Lake E under RCP 8.5. On the other hand, ice thickness in Lake A is suggested by the model to only decrease by 2.8 cm decade^{-1}. Overall, lake ice thickness in Northern Poland will likely decrease by between 3.7 and 3.9 cm decade^{-1} (RCP 6.0 and 8.5, respectively) within the current century. Decrease in ice cover duration will range between 4.5- and 10-day decade^{-1} for Lake A and E (RCP 6.0 and 8.5) with a mean ice cover duration loss for an average lake in this region between 7.1- and 9.5-day decade^{-1}.

For the modelled historical dataset (1965-2005) none of the monitored lakes demonstrated stable ice-free conditions. Therefore, the probability of ice-free winters increased for all the lakes and in all future climate scenarios. For RCP 2.6 the increase ranged from 4 to 46% in Lake A and B, respectively. For RCP 6.0 the probability of ice-free winter increased by between 8 to 58% and under RCP 8.5 the probability for these lakes to remain ice free increased by between 18 and 69%.

4. Discussion

The observed decrease in lake ice cover duration and thickness in Northern Poland is similar to those observed in lakes around the globe (Magee et al., 2016; Lopez et al., 2019; Sharma et al., 2019). The impact of recent climate change allowed us to explain a large fraction of this negative trend over the last sixty years. However, while most of the variability in ice conditions can be attributed to the effects of warming, seasonal changes in wind speed as well as rainfall also accounted for part of the trend. Interestingly, while lower winds during the freezing period stimulated ice formation, the relationship was reversed in fall when, in some of the studied lakes, stronger winds accelerated freezing likely due to enhanced evaporation and cooling. This effect illustrates the seasonality of meteorological influence on lake ice phenology when mixing and

283 heat exchange (i.e., thermal homogenization) is controlled by shear forcing. For instance, calm
284 conditions will trigger ice formation in lakes under freezing temperatures but only when surface
285 waters previously cooled down more than deeper ones through wind-enhanced heat loss.
286
287 Conversely, effects of increasing precipitation were evident mostly during the break-up
288 of ice cover. Precipitation can stimulate the thawing of lake ice as much or more than the effect
289 of warming. We also provide some indications that the location and shape of the basin in relation
290 to prevalent wind direction may be an important factor to consider for better predictions of future
291 lake ice phenology. Notwithstanding the dominant effect of warming, associated changes (i.e.,
292 atmospheric stilling, higher winter and springtime rainfall) need to be accounted for on a possibly
293 more temporally resolved (i.e., monthly) scale as similar trends may have contrasting effects on
294 ice formation depending on the season and stratification stage (Table 2). These seasonal effects
295 should be more closely considered to better understand and predict the rate of lake ice loss in the
296 future.
297
298 The ubiquitous observational evidence of decreasing ice conditions in temperate and
299 boreal lakes over the last century implies that ice cover is responding rapidly to the effects of
300 global climate change. The rate at which ice cover is decreasing in lakes is globally variable
301 (Sharma et al. 2019) and depends on the strength of climatic effects on the ice formation and
302 break-up times as well as on the size and shape of each individual lake. Previous studies have
303 reported a decline in the ice cover duration for lakes in Poland by between 1.0-day decade^{-1} (Ptak
304 et al., 2017) for the deep alpine lake Morskie Oko and 8.2-day decade^{-1} for Lake Ełckie (Choiński
305 et al., 2015). For the temperate lowland lakes considered here, the calculated trend varied
306 between 3.9- and 7.8-day decade^{-1} (for lake B – Charzykowskie and E – Studzieniczne). These
307 rates are comparable to those reported for other lakes in temperate latitudes (i.e., Bernhardt et al.,

2012; Apsite et al., 2014; Soja et al., 2014) but a rather large range of changes is apparent despite comparable effects in average air temperature. For instance, in the relatively shallow Lake A (1.3 m deep), where the circulation is influenced by the oceanic climate and the water column mixes multiple times between summer and fall (polymixis; Ptak et al., 2019b), the ice cover formation is usually delayed until January. This lake already remains ice free during exceptionally warm winters (e.g., 2007/2008). Bottom waters of Lake B during the stratification period remain warm (>8ºC, Garbacz et al., 2008), and the hypolimnetic volume is moved upwards by sinking surface waters in fall. This buoyancy flux is the main cause of the mixing and likely one of the most important reasons for observed delay in the ice cover formation. By contrast, Lake C freezes relatively fast (on average frozen by mid-December). This lake, also polymictic, is at times exposed to strong winds along rather than across the lake and, thus, the wind-induced mixing is efficient (Woolway and Simpson, 2017). The shape and complexity of the lake basin in relation to mixing efficiency apparently also influence the responsiveness of the ice cover to changes in climate (Magee and Wu, 2017).

In our regression analyses we used the Cochrane-Orcutt method to eliminate autocorrelation. The model estimates including effects of temperature, wind and rainfall best explained the observations when compared to any reduced model configuration (i.e., relatively high F and low p values; Table 3). The analysis of errors (i.e., MSE, RMSE) generally resulted in higher values for the estimates of ice cover duration as compared to the estimated ice thickness. This potentially indicates that there are some other factors or interactions that were not considered in this study but could have improved the goodness of fit for the predicted changes in ice cover duration. Noticeably, greatest errors were estimated for analyses in Lake C (Table 3) which is, in fact, the studied lake with the most complex morphometry (Fig 2). This may potentially indicate that the effect of morphometry in lakes with complex shape should be considered more

333 directly to improve our predicted ice phenology. Improved process understanding and future
334 predictions will also require accounting for these effects and potential biases.

335 All study lakes will be losing their ice cover during the 21st century according to current
336 climate projections (GHG, Figs. 6 & 7). In fact, among the lakes in Northern Poland, the deep
337 and symmetrical Lake E will be losing its ice most rapidly (i.e., 4.8 cm decade^{-1} under RCP 8.5)
338 and is likely to experience ice-free winters with much higher probability than in the past (up to
339 53% increase). Shallow westerly Lake B will be losing its ice cover at an average rate of 2.7 cm
340 decade^{-1}, almost half the rate of Lake E. This inter-lake difference appears to result from the fact
341 that the ice cover in Lake E is strongly influenced by the effect of changes in air temperatures
342 (rapid warming under RCP 8.5) as compared to the relatively more important influence of
343 hydroclimate in shallow Lake B. However, this also results from the fact that ice cover formation
344 in Lake B was much delayed when compared to Lake E already by 2015 and the lake ice remained
345 respectively thinner. Notwithstanding difference in rates of ice cover decrease, the probability of
346 ice-free winters will also increase rapidly in Lake B (up to 69% under RCP 8.5). Effects of future
347 climate on ice conditions will depend on the location and shape of the lake but also on the
348 duration of recent ice cover that may influence the responsiveness of spring mixing/stratification
349 patterns to atmospheric warming.
350
351 Recent studies have suggested that ice cover in shallow lakes that mix frequently will be
352 relatively less influenced by atmospheric warming compared to lakes that are strongly stratified
353 in summer (Magee and Wu, 2017). This conclusion, in part consistent with our findings (i.e., in
354 Lake A), stems from the assumption that, as climate warms, deeper lakes will gain more heat
355 throughout the upper and mid-water column since the diurnal heat losses will be less than in
356 shallower polymictic lakes. This is correct when comparing deeper and shallower lakes under
357 similar winds (and effective fetch) inducing heat loss or thermally homogenizing the water

column. Once the lake size or fetch is considered, it may be more a function of prevalent wind direction and intensity of surface warming in the day that will control diurnal and seasonal heat exchange (Waples and Klump, 2002) and thus directly influence the exact timing of overturn and subsequent ice formation. We argue here that the stratification, mixing patterns and efficiency of heat exchange in summer and fall rather than size or depth may also have a more direct impact on future lake ice phenology.

Stratification in lakes is controlled by an interaction of heat fluxes with the wind energy inputs and inflows following heavy rainfall events (Laborde et al., 2010). Heavy rainfalls in autumn which occur simultaneously with strong winds at the lake surface, can be particularly important in triggering mixing and cooling throughout the water column (Kimura et al., 2017). In fact, large rainfall inflows (relatively to lake volume) have been suggested to control the lake-wide circulation patterns (Carmack 1979; Killworth and Carmack 1979). The changes in autumnal rainfalls may thus indirectly influence ice formation through accelerated or delayed mixing and temperature homogenization. Recent studies have also suggested that raindrops falling into water-unsaturated air will cool through evaporation thus their passage can lead to cooling of the air (Rooney et al., 2018). This mechanism may further enhance convective mixing in the surface at the end of a heavy rainfall event in autumn and allow for a more rapid ice formation. The timing of lake overturn and subsequent freezing during and after autumnal rain- and snowstorms will ultimately depend on the amount of energy required to overcome the density gradients between surface and bottom waters. Standard indices used in physical limnology (i.e., Schmidt or Wedderburn numbers; Imberger and Patterson, 1989) to describe the ratio between meteorological forcing and the gradient of pressure established by the stratification may provide an excellent proxy to investigate the effects of future stronger stratification on lake ice conditions.

Lakes are likely to stratify more strongly and for a longer period in the near future as a consequence of direct and indirect effects of climate and global environmental change (Woolway and Merchant, 2019, Bartosiewicz et al., 2019a, b). The seasonal as well as interannual changes in the strength of thermal stratification and efficiency of heat exchange should be considered for better predictions of the future lake ice phenology. For instance, while in currently polymictic lakes heat is gained and lost rapidly throughout the water column between summer and fall, these water bodies are likely to retain part of the heat for longer when the water column is most stably stratified. The efficiency of heat retention and downward transport will depend on the water column transparency (Ptak et al., 2018). In browning or greening lakes (Leech et al., 2018) the effects of atmospheric warming will accumulate rapidly in surface waters, leading to a thinner and warmer epilimnion ("thermal shielding"). Under such conditions lakes stratify early in spring and are more likely to remain stratified for longer in fall (Bartosiewicz et al., 2015). Stratification precludes thermal homogenization and can effectively result in delayed ice formation. On the other hand, thermally shielded bottom waters of less transparent lakes are likely to gain less heat during the summer and thus remain relatively cold throughout the summer (Bartosiewicz et al., 2016). Therefore, there will be less heat from deep waters to be lost during the overturn before the ice is formed (Ye et al., 2018). These contrasting effects, likely to influence lake ice phenology in the near future, need to be further explored to improve our understanding of ice processes and changes.

While changes of the stratification and heat retention in lakes will affect the timing of ice formation, the duration of the ice cover will in turn control the onset of stratification. This potential feedback effect, which to our knowledge has not been yet comprehensively studied, can result in a cascading change of the ice phenology in many shallow lakes around the globe. If a shallow lake mixes less often or even remains stably stratified during the summer in a warmer

408 climate, the outcome for the timing of ice formation will depend on the amount of heat
409 (temperature) of bottom waters. There are two possible scenarios that are worth considering.
410 First, if bottom waters gain sufficient heat in springtime as stratification develops slowly, and
411 still gain some heat over the summer (as a function of high transparency), then the ice formation
412 will be delayed upon autumnal mixing until this heat is lost to the atmosphere. Second, if bottom
413 waters do not gain much heat during the springtime when stratification develops rapidly and do
414 not warm up over the summer (as a function of low transparency), ice formation will follow the
415 autumnal overturn in short order. The overall outcome depends on whether faster and more
416 enhanced stratification in less transparent lakes will be sufficiently strong to delay mixing (and
417 following freezing) as much as upwelling of warmer bottom waters in more transparent lakes.
418
419 The enhanced warming of surface waters in less transparent lakes may interact with the
420 effect of increased rainfall in the catchment and decreasing wind speeds over the lake to result in
421 faster and stronger thermal stratification. While such a direct effect of these co-occurring
422 processes on ice conditions in temperate lakes are apparent from the current study, their indirect
423 effects through changes in water transparency and differential heating in the upper and lower
424 water column require further investigation in the future. The arising feedback effect appears
425 particularly important if we consider the major implications of the changing ice conditions for
426 the functioning of lakes. For example, the duration of the ice cover, directly related to the onset
427 and duration of thermal stratification, has been recognized as one of the major determinants of
428 the springtime and summer warming trends (O'Reilly et al., 2015). The cascading influence of
429 the ice phenology also affects the formation and magnitude of spring and summer growth of
430 phytoplankton blooms (Adrian et al., 1999, Blenckner et al., 2007), their functional diversity
431 (Özkundakci et al., 2016) and the abundance of zooplankton (Dokulil et al., 2014). Pronounced
432 and long-lasting effects of warming air temperatures on the functioning of lakes may be further

enhanced by accelerated warming of surface waters in ecosystems that are or will be losing their ice completely (Kintisch, 2015). Delayed ice formation and decreasing ice thickness and duration will all lead to less stable thermal conditions under the ice (Bruesewitz et al., 2015). These changes, in turn, may affect the primary production and oxygen dynamics in frozen lakes and are potentially related to effects throughout the aquatic food webs (Beall et al., 2016) and accelerated emission of greenhouse gases (Denfeld et al., 2016). Adding to potential effects on biology and biogeochemistry of lakes, future ice decline and predicted increased in the frequency of ice-free conditions will be likely to also affect ecosystem services and regional economy (Knoll et al., 2019)

Conclusions

This study supports the interactive effects of warming, stilling and changing precipitation patterns on the ice conditions in temperate lakes. Polish lakes are losing their ice, and this change is driven mostly by direct effects of temperature and to some extent by the effect of increasing rainfall. The effect of wind is either negative when it delays ice formation in fall or positive when it stimulates ice thickening throughout the winter. Predictive simulation, based on simple hydrodynamic model and an ensemble of four climate model projections, suggests that under continuing emissions, most lakes in northern Poland will be losing their ice cover rapidly and may become largely ice-free by the end of the century. The large spectrum of responsiveness in ice phenology and conditions to changing weather conditions most likely results from the interactive effects between surface meteorology, lake size and shape as well as the strength of thermal stratification. In this context it is important to underline that predicted changes in the mixing regime of global lakes will most likely have important consequences for ice phenology. Changes in ice phenology may also result in accelerated shifts toward different mixing of lakes. This potential effect needs to be further explored. Lake ecosystems that are prone to remain ice-

458 free in consequence of warming or increased precipitation will function differently than in the
459 past.
460

461 **Acknowledgements**

462 This research was supported by SNF Project 169552 (2016-2019) and its ongoing extension

463 (2020-2021).

464

References

Adrian, R., Walz, N., Hintze, T., Hoeg, S., Rusche, R., 1999. Effects of ice duration on plankton succession during spring in a shallow polymictic lake, Freshwater Biology, 41, 3: 621–632.

Apsite, E., Elferts, D., Zubaničs, A., Latkovska, I., 2014. Long-term changes in hydrological regime of the lakes in Latvia, Hydrology Research, 45, 3: 308–321.

Arp, C.D., Jones, B.M., Grosse, G., 2013. Recent lake ice-out phenology within and among lake districts of Alaska, U.S.A., Limnology and Oceanography, 58, 6: 2013–2028.

Bartosiewicz, M., Laurion, I., MacIntyre, S., 2015. Greenhouse gas emission and storage in a small shallow lake. Hydrobiologia, 757, 1: 101–115.

Bartosiewicz, M., Laurion, I., Clayer, F., & Maranger, R., 2016. Heat-wave effects on oxygen, nutrients, and phytoplankton can alter global warming potential of gases emitted from a small shallow lake. Environmental science & technology, 50, 12: 6267–6275.

Bartosiewicz, M., Przytulska, A., Deshpande, B. N., Antoniades, D., Cortes, A., MacIntyre, S., Laurion, I., 2019a. Effects of climate change and episodic heat events on cyanobacteria in a eutrophic polymictic lake. Science of The Total Environment, 693, 133414.

Bartosiewicz, M., Przytulska, A., Lapierre, J. F., Laurion, I., Lehmann, M. F., & Maranger, R., 2019b. Hot tops, cold bottoms: Synergistic climate warming and shielding effects increase carbon burial in lakes. Limnology and Oceanography Letters, 4, 5: 132–144.

Beall, B.F., Twiss, M.R., Smith, D.E., Oyserman, B.O., Rozmarynowycz, M.J., Binding, C.E., Bourbonniere, R.A., Bullerjahn, G.S., Palmer, M.E., Reavie, E.D., Waters, L.M., Woityra, L.W., McKay, R.M. 2016. Ice cover extent drives phytoplankton and bacterial community structure in a large north-temperate lake: implications for a warming climate, Environmental microbiology,18, 6: 1714–1719.

Bengtsson, L., 1986. Dispersion in ice-covered lakes. Hydrology Research, 17, 3: 151-170.

Bengtsson, L., 1996. Mixing in ice-covered lakes. In: The First International Lake Ladoga Symposium (pp. 91-97). Springer, Dordrecht.

Benson, B. J., Magnuson, J. J., Jensen, O. P., Card, V.M., Hodgkins, G., Korhonen, J., Livingstone D. M., Stewart K. M., Weyhenmeyer G.A., Granin N.G., 2012. Extreme events, trends, and variability in northern hemisphere lake-ice phenology (1855–2005). Climatic Change, 112, 2: 299–323.

Bernhardt, J., Engelhardt, C., Kirillin, G., Matschullat, J., 2012. Lake ice phenology in Berlin-Brandenburg from 1947–2007: Observations and model hindcasts, Climatic Change, 112, 3–4: 791–817.

Blenckner, T., Adrian, R., Livingstone, D. M., Jennings, E., Weyhenmeyer, G. A., George, D. G., Straile, D., 2007. Large-scale climatic signatures in lakes across Europe: A meta-analysis. Global Change Biology, 13, 7: 1314–1326.

514 Bruesewitz, D.A., Carey, C.C., Richardson, D.C., Weathers, K.C., 2015. Under-ice thermal
515 stratification dynamics of a large, deep lake revealed by high-frequency data, Limnology and
516 Oceanography, 60, 2: 347–359.
517
518 Bouffard, D., Zdorovennova, G., Bogdanov, S., Efremova, T., Lavanchy, S., Palshin, N.,
519 Zdorovennov, R., 2019. Under-ice convection dynamics in a boreal lake. Inland Waters, 9, 2:
520 142–161.
521
522 Caine, N., 2002. Declining ice thickness on an Alpine Lake is generated by increased winter
523 precipitation, Climatic Change, 54, 4: 463–470.
524
525 Carmack, E. C. (1979). Combined influence of inflow and lake temperatures on spring circulation
526 in a riverine lake. Journal of Physical Oceanography, 9(2), 422–434.
527
528 Choiński A., 2006. Catalogue of Polish Lakes, Wyd. Nauk. UAM, Poznań.
529
530 Choiński A., Ptak M., Skowron R., Strzelczak A., 2015. Changes in ice phenology on polish
531 lakes from 1961-2010 related to location and morphometry, Limnologica, 53: 42–49.
532
533 Czernecki, B., Ptak, M., 2018. The impact of global warming on lake surface water temperature
534 in Poland - the application of empirical-statistical downscaling, 1971–2100, Journal of
535 Limnology, 77, 2: 330–348.
536

537 Denfeld, B.A., Kortelainen, P., Rantakari, M., Sobek, S., Weyhenmeyer, G.A., 2016. Regional
538 Variability and Drivers of Below Ice CO_2 in Boreal and Subarctic Lakes, Ecosystems, 19, 3:
539 461–476.
540
541 Dokulil, M.T., Herzig, A., Somogyi, B., Vörös, L., Donabaum, K., May L. & Nõges, T. 2014.
542 Winter conditions in European shallow lakes: a comparative synopsis. Estonian Journal of
543 Ecology 63: 111–129.
544
545 Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L. & Geiger, T. 2017.
546 Assessing the impacts of 1.5 C global warming–simulation protocol of the Inter-Sectoral Impact
547 Model Intercomparison Project (ISIMIP2b). Geoscientific Model Development.
548
549 Garbacz J.K., Cieściński J., Ciechalski J., Dąbkowski R., Cichowska J., 2018. Thermal and
550 oxygen conditions of Lake Charzykowskie in the years 2014-2016, Polish Hyperbaric Research,
551 62, 1: 85–96.
552
553 Hampton, S.E., Galloway, A.W.E., Powers, S.M., Ozersky, T., Woo, K.H., Batt, R.D., Labou,
554 S.G., O'Reilly, C.M., Sharma, S., Lottig, N.R., Stanley, E.H., North, R.L., Stockwell, J.D.,
555 Adrian, R., Weyhenmeyer, G.A., Arvola, L., Baulch, H.M., Bertani, I., Bowman, L.L., J, Carey,
556 C.C., Catalan, J., Colom-Montero, W., Domine, L.M., Felip, M., Granados, I., Gries, C.,
557 Grossart, H.-P., Haberman, J., Haldna, M., Hayden, B., Higgins, S.N., Jolley, J.C., Kahilainen,
558 K.K., Kaup, E., Kehoe, M.J., MacIntyre, S., 2017. Ecology under lake ice, Ecology Letters, 20,
559 1: 98–111.
560

561 Helland, I.P., Finstad, A.G., Forseth, T., Hesthagen, T., Ugedal, O.,2011. Ice-cover effects on
562 competitive interactions between two fish species, Journal of Animal Ecology, 80, 3: 539–547.
563 Imberger, J., and Patterson, J. C. 1989. Physical limnology. In Advances in applied mechanics,
564 27: 303–475.
565 Kainz, M.J., Ptacnik, R., Rasconi, S., Hager, H.H., 2017. Irregular changes in lake surface water
566 temperature and ice cover in subalpine Lake Lunz, Austria, Inland Waters, 7, 1: 27–33.
567
568 Killworth, P. D., & Carmack, E. C. (1979). A filling-box model of river-dominated lakes.
569 Limnology and oceanography, 24(2), 201–217.
570
571 Knoll, L.B., Sharma, S., Denfeld, B. et al. 2019 Consequences of lake and river ice loss on
572 cultural ecosystem services. Limnology and Oceanography Letters 4: 119–131.
573
574 Kimura, N., Wen-Cheng, L., Jeng-Wei, T., Chih-Yu, C., Kratz, T. K., & Tai, A. (2017).
575 Contribution of extreme meteorological forcing to vertical mixing in a small, shallow subtropical
576 lake. Journal of Limnology, 76(1).
577
578 Kintisch, E. 2015. Earth's lakes are warming faster than its air: First ever global survey reveals
579 summer lake temperatures rising at an alarming rate, Science, 350, 6267: 1449.
580
581 Kirillin, G., Leppäranta, M., Terzhevik, A., Granin, N., Bernhardt, J., Engelhardt, C.,
582 Zdorovennova, G, 2012. Physics of seasonally ice-covered lakes: a review. Aquatic sciences, 74,
583 4: 659–682.
584

585 Kolendowicz, L., Czernecki, B., Półrolniczak, M., Taszarek, M., Tomczyk, A.M., Szyga-Pluta,
586 K., 2019. Homogenization of air temperature and its long-term trends in Poznań (Poland) for the
587 period 1848–2016. Theoretical and Applied Climatology 136, 3:1357–1370.
588
589 Laborde, S., Antenucci, J. P., Copetti, D., & Imberger, J. (2010). Inflow intrusions at multiple
590 scales in a large temperate lake. Limnology and Oceanography, 55, 3: 1301–1312.
591
592 Lange, S. 2019. EartH2Observe, WFDEI and ERA-Interim data Merged and Bias-corrected for
593 ISIMIP (EWEMBI). V.1.1. GFZ Data Services. http://doi.org/10.5880/pik.2019.004.
594
595 Leppäranta, M., 2010. Modelling the formation and decay of lake ice. In The impact of climate
596 change on European lakes. Springer, Dordrecht pp. 63–83.
597
598 Lopez, L.S., Hewitt, B.A., Sharma, S., 2019. Reaching a breaking point: How is climate change
599 influencing the timing of ice breakup in lakes across the northern hemisphere? Limnology and
600 Oceanography, 64, 6: 2621–2631.
601
602 Magee, M.R., Wu, C.H., Robertson, D.M., Lathrop, R.C., Hamilton, D.P., 2016. Trends and
603 abrupt changes in 104 years of ice cover and water temperature in a dimictic lake in response to
604 air temperature, wind speed, and water clarity drivers, Hydrology and Earth System Sciences,
605 20, 5: 1681–1702.
606
607 Magee, M.R., Wu, C.H., 2017. Effects of changing climate on ice cover in three
608 morphometrically different lakes, Hydrological Processes, 31, 2: 308–323.
609

Magnuson, J.J., Robertson, D.M., Benson, B.J., Wynne, R.H., Livingstone, D.M., Arai, T., Assel, R.A., Barry, R.G., Card, V., Kuusisto, E., Granin, N.G., Prowse, T.D., Stewart, K.M., Vuglinski, V.S., 2000. Historical trends in lake and river ice cover in the Northern Hemisphere, Science, 289, 5485: 1743–1746.

Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J. F., ... & Thomson, A. G. J. M. V. (2011). The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. Climatic change, 109, 2: 213–241.

Mironov, D., Terzhevik, A., Kirillin, G., Jonas, T., Malm, J., Farmer, D., 2002. Radiatively driven convection in ice-covered lakes: Observations, scaling, and a mixed layer model. Journal of Geophysical Research: Oceans, 107, 4: 7–1.

Mironov, D., Heise, E., Kourzeneva, E., Ritter, B., Schneider, N., Terzhevik, A. 2010. Implementation of the lake parameterization scheme FLake into the numerical weather prediction model COSMO. Boreal Environ. Res. 15: 218–230.

O'Reilly, C. M., Sharma, S., Gray, D. K., Hampton, S. E., Read, J. S., Rowley, R. J., ... & Weyhenmeyer, G. A., 2015. Rapid and highly variable warming of lake surface waters around the globe. Geophysical Research Letters, 42, 24: 10-773.

Orru, K., Kangur, K., Kangur, P., Ginter, K., Kangur, A., 2014. Recreational ice fishing on the large lake peipsi: Socioeconomic importance, variability of ice-cover period, and possible implications for fish stocks, Estonian Journal of Ecology, 63,4:282–298.

634 Özkundakci, D., Gsell, A. S., Hintze, T., Täuscher, H., & Adrian, R. (2016). Winter severity
635 determines functional trait composition of phytoplankton in seasonally ice-covered lakes. Global
636 change biology, 22, 1: 284–298.
637
638 Preston, D.L., Caine, N., McKnight, D.M., Williams, M.W., Hell, K., Miller, M.P., Hart, S.J.,
639 Johnson, P.T.J., 2016. Climate regulates alpine lake ice cover phenology and aquatic ecosystem
640 structure, Geophysical Research Letters, 43, 10: 5353–5360.
641
642 Prowse, T.D., Brown, K., 2010. Hydro-ecological effects of changing Arctic river and lake ice
643 covers: a review. Hydrol. Res. 41, 6: 454–461.
644
645 Ptak M., Wrzesiński D., Choiński A. 2017. Long-term changes in the hydrological regime of
646 high mountain lake Morskie Oko (Tatra Mountains, Central Europe). Journal of Hydrology and
647 Hydromechanics, 65, 2: 146–153.
648
649 Ptak M., Sojka M., Choiński A., Nowak B. 2018. Effect of environmental conditions and
650 morphometric parameters on surface water temperature in Polish lakes. Water, 10: 580–590.
651
652 Ptak M., Tomczyk A., Wrzesiński D., Bednorz E., 2019a. Effect of teleconnection patterns on
653 ice conditions in lakes in lowland Poland, Theoretical and Applied Climatology, 138:1961–1969.
654
655 Ptak, M., Sojka M., Kozłowski, M., 2019b. The increasing of maximum lake water temperature
656 in lowland lakes of Central Europe: case study of the Polish Lakeland, Annales de Limnologie -
657 International Journal of Limnology, 55, 6: 1–11.
658

659 Rooney, G. G., Van Lipzig, N., & Thiery, W. 2018. Estimating the effect of rainfall on the surface
660 temperature of a tropical lake. Hydrology and Earth System Sciences, 22: 6357–6369.
661
662 Schmidt, D.F., Grise, K.M. & Pace, M.L. 2019. High-frequency climate oscillations drive ice-
663 off variability for Northern Hemisphere lakes and rivers. Climatic Change 152: 517–532
664 https://doi.org/10.1007/s10584-018-2361-5.
665 Sharma, S., Magnuson, J. J., Batt, R. D., Winslow, L. A., Korhonen, J., & Aono, Y., 2016. Direct
666 observations of ice seasonality reveal changes in climate over the past 320–570 years. Scientific
667 Reports, 6, 25061.
668
669 Sharma, S., Blagrave, K., Magnuson, J.J., O'Reilly, C.M., Oliver, S., Batt, R.D., Magee, M.R.,
670 Straile, D., Weyhenmeyer, G.A., Winslow, L., Woolway, R.I., 2019. Widespread loss of lake ice
671 around the Northern Hemisphere in a warming world, Nature Climate Change, 9, 3:227–231.
672
673 Soja, A.-M., Kutics, K., Maracek, K., Molnár, G., Soja, G., 2014. Changes in ice phenology
674 characteristics of two Central European steppe lakes from 1926 to 2012 - influences of local
675 weather and large-scale oscillation patterns, Climatic Change, 126, 1–2: 133.
676
677 Tan, Z., Yao, H., Zhuang, Q., 2018. A Small Temperate Lake in the 21st Century: Dynamics of
678 Water Temperature, Ice Phenology, Dissolved Oxygen, and Chlorophyll a, Water Resources
679 Research, 54, 7: 4681–4699.
680
681 Terzhevik, A., Golosov, S., Palshin, N., Mitrokhov, A., Zdorovennov, R., Zdorovennova, G., ...
682 & Zverev, I., 2009. Some features of the thermal and dissolved oxygen structure in boreal,
683 shallow ice-covered Lake Vendyurskoe, Russia. Aquatic ecology, 43, 3: 617–627.

Tomczyk A.M., Szyga-Pluta K. 2019. Variability of thermal and precipitation conditions in the growing season in Poland in the years 1966–2015. Theoretical and Applied Climatology 135, 3–4: 1517–1530.

Waples, J. T., & Klump, J. V. (2002). Biophysical effects of a decadal shift in summer wind direction over the Laurentian Great Lakes. Geophysical research letters, 29, 8: 43–51.

Weyhenmeyer, G.A., Meili, M., Livingstone, D.M. 2004. Nonlinear temperature response of lake ice breakup. Geophys. Res. Lett., 31, L07203.

Williams, G., Layman, K.L., Stefan, H.G., 2004. Dependence of lake ice covers on climatic, geographic and bathymetric variables, Cold Regions Science and Technology, 40, 3: 145–164.

Woolway, R.I., Simpson, J.H. 2017. Energy input and dissipation in a temperate lake during the spring transition. Ocean Dynamics 67: 959–971.

Woolway, R. I., & Merchant, C. J. (2019). Worldwide alteration of lake mixing regimes in response to climate change. Nature Geoscience, 12, 4: 271–276.

Woolway, R.I., Weyhenmeyer, G.A., Schmid, M., Dokulil, M.T., de Eyto, E., Maberly, S.C., May, L., Merchant, C.J., 2019. Substantial increase in minimum lake surface temperatures under climate change. Climatic Change, 155: 81–94.

Wu, T., Qin, B., Zhu, G., Huttula, T., Lindfors, A., Ventelä, A.-M., Sheng, Y., Ambrose, R.F., 2018. The contribution of wind wave changes on diminishing ice period in Lake Pyhäjärvi during the last half-century, Environmental Science and Pollution Research, 25, 25: 24895–24906.

Yao, H., Samal, N.R., Joehnk, K.D., Fang, X., Bruce, L.C., Pierson, D.C., Rusak, J.A., James, A., 2014. Comparing ice and temperature simulations by four dynamic lake models in Harp Lake: Past performance and future predictions, Hydrological Processes, 28, 16: 4587–4601.

Ye, X., E. J. Anderson, P. Y. Chu, C. Huang, Xue P., 2018. Impact of Water Mixing and Ice Formation on the Warming of Lake Superior: A Model-guided Mechanism Study. Limnology and Oceanography, 64, 2; 558–574. doi:10.1002/lno.11059.

719 **Tables**

720 Table. 1. Geographic coordinates (latitude – Lat.; longitude – Lon.) and limnological
721 characteristics of the five study lakes in northern Poland (after: Choiński 2006, Ptak et al. 2018;
722 arranged from west to east – A to E), including area, volume, mean and max depth, water
723 transparency (as Secchi depth, SD) and wind fetch (FT).

ID	Lake	Lat.	Lon.	Area (ha)	Volume ($10^3 \times m^3$)	Mean depth (m)	Max depth (m)	SD (m)	FT (km)
A	Gardno	54.64	17.16	2337.5	30951	1.3	2.6	0.7	6.8
B	Charzykowskie	53.73	17.50	1336.0	134533	9.8	30.5	2.1	9.5
C	Jeziorak	53.59	19.55	3152.5	141594	4.1	12.9	0.8	13.3
D	Mikołajskie	53.80	21.56	424.0	55740	11.2	25.9	1.3	5.8
E	Studzieniczne	53.86	23.09	244.0	22074	8.7	30.5	2.7	3.4

724

Table 2. Standardized coefficients (beta coefficients) from the multiple regression analysis of the relationship between seasonal effects of weather conditions, including air temperature (November to March), wind speeds (September to January) and rainfall (November to March), and ice conditions in temperate lakes in northern Poland. Values in bold are given for parameters that significantly ($p<0.05$) improved the predictability.

	Lake A		Lake B		Lake C		Lake D		Lake E		All Lakes	
	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick
Temp. Nov	0.01	**-0.16**	-0.02	0.11	-0.06	0.01	-0.06	-0.11	**-0.22**	-0.05	**-0.08**	-0.05
Dec	**-0.21**	-0.12	**-0.39**	0.15	**-0.20**	-0.05	**-0.43**	**-0.29**	**-0.30**	-0.14	**-0.30**	**-0.14**
Jan	**-0.35**	**-0.44**	**-0.38**	**-0.25**	**-0.29**	**-0.38**	**-0.30**	**-0.42**	-0.05	**-0.35**	**-0.23**	**-0.31**
Feb	**-0.33**	**-0.38**	0.05	-0.19	**-0.43**	**-0.42**	**-0.36**	**-0.41**	**-0.45**	**-0.52**	**-0.32**	**-0.43**
Mar	-0.05	0.04	**-0.22**	-0.15	-0.14	-0.01	-0.16	-0.09	**-0.23**	**-0.20**	**-0.20**	**-0.11**
Wind Sept	-0.02	0.05	0.09	0.02	-0.03	0.07	-0.10	0.05	-0.05	-0.11	0.05	0.02
Oct	**-0.09**	-0.06	-0.05	**-0.21**	-0.06	**-0.18**	-0.01	0.01	-0.15	**-0.18**	**-0.08**	**-0.08**
Nov	-0.02	**0.17**	**-0.13**	-0.14	0.06	**0.23**	0.08	**-0.23**	**0.31**	0.08	0.03	**0.15**
Dec	0.01	-0.14	-0.09	**-0.30**	0.12	-0.03	0.06	-0.11	0.07	0.10	0.01	-0.05
Jan	-0.12	-0.02	-0.02	0.07	0.01	0.04	0.12	**0.38**	-0.07	**0.35**	-0.02	0.04
Rain Nov	**0.13**	0.04	0.09	**-0.28**	0.04	0.02	-0.07	0.11	-0.03	-0.04	**0.12**	0.01
Dec	0.03	0.01	-0.08	0.07	**-0.16**	-0.06	0.04	-0.07	0.07	**-0.21**	0.06	-0.03
Jan	-0.08	0.08	**0.19**	-0.02	0.01	**-0.19**	-0.12	**-0.15**	**-0.13**	**-0.19**	-0.01	**-0.09**
Feb	-0.09	0.05	**-0.20**	-0.03	0.01	0.02	0.08	-0.02	-0.04	0.08	-0.01	0.01
Mar	**-0.12**	0.02	**-0.27**	**-0.22**	-0.08	0.03	-0.10	-0.09	0.01	0.06	**-0.12**	-0.04
R^2	**0.88**	**0.89**	**0.84**	**0.78**	**0.79**	**0.79**	**0.76**	**0.84**	**0.72**	**0.80**	**0.71**	**0.75**

732 Table 3. Goodness of fit statistics (MSE – mean square error; root-mean-square error – RMSE; MAPE – mean absolute percentage error and DW
733 – Durbin Watson Statistics) for the regression analyses (Cochrane-Orcutt method) between changes in air temperature, wind and rainfall conditions
734 on lake ice duration and thickness (1965-2015) in temperate lakes of Norther Poland.

	Lake A		Lake B		Lake C		Lake D		Lake E		All Lakes	
	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick	Dur	Thick
DF	39	39	38	38	40	40	38	38	40	40	259	259
F	27.4	21.4	14.5	8.5	10.1	9.8	12.2	18.6	6.9	15.6	41.3	54.5
MSE	188.0	21.75	174.2	40.9	333.2	51.8	303.7	37.1	290.8	38.8	340.1	46.2
RSME	13.7	4.7	13.2	6.4	18.3	7.2	17.5	6.1	17.0	6.2	18.4	6.8
MAPE	14.9	17.7	26.4	25.3	12.0	15.5	21.6	13.3	12.9	9.5	41.4	22.9
DW	1.96	1.96	1.90	1.92	2.08	2.04	1.97	2.03	2.06	2.00	2.01	1.98

735

736 Table 4. The relative increase in the probability (%) of ice-free conditions for the study lakes
737 between2020-2100 estimated using Flake model and an ensemble of four climate projections
738 (GFDL-ESM2M, HadGEM, IPSL and MIROC5) simulated under three relevant GHG concentration
739 trajectories – RCP 2.6 being the most conservative scenario (emissions declining by 2020 and
740 reaching zero by 2100), RCP 6.0 being moderate (emissions peak around 2080, then decline) and
741 RCP 8.5 being the least conservative (emissions continue to rise throughout the 21st century).

742

Lake	RCP2.6	RCP6.0	RCP8.5
			743
A	3.8	8.9	17.7
B	46.8	58.2	69.2
C	10.1	25.3	40.5
D	35.4	41.8	63.3
E	16.5	34.2	54.4

744 **Captions**

745 **Figure 1. Interactive effects of warming, changes in wind speed (stilling) and precipitation on**
746 **lake ice formation** (also as the Graphical Abstract) under three future greenhouse gas (GHG)
747 emission scenarios. Size and direction (downward-negative, upward-positive) of arrows indicate the
748 relative strength and direction of temperature, wind and rainfall on ice conditions in study lakes.

749

750 **Figure 2. Location, morphometry and bathymetry of the study lakes** as well as location of the
751 nearest meteorological station with continuous record between 1961 and 2017. Arrows indicate
752 prevalent wind directions (between 1980 and 2017).

753

754 **Figure 3. Surface meteorology** in the vicinity of temperate lakes in Northern Poland (A-
755 Charzykowskie, B-Gardno, C-Jeziorak, D- Mikolajskie, E-Studzieniczne, F-average for the entire
756 region) including air temperature (in red), wind speed (in black) and precipitation (in blue) as monthly
757 averages between 1961 and 2017 for the five lakes (n = 257). R is given for all trends but significant
758 ones ($p<0.05$) are shown in bold.

759

760 **Figure 4. Changes in the ice cover conditions** (thickness in grey and duration in black) for each
761 temperate lake in northern Poland (A-E) and for all lakes together (F). R indicates temporal trends
762 between 1961-2017, significant ($p<0.05$) ones are shown in bold.

763

764 **Figure 5. Results of the regression analysis (predicted duration and thickeness)** between air
765 temperature, wind speeds, rainfall) and ice conditions (ice cover thickness in grey and duration in
766 black) for each lake separately (panels A to E) and for all lakes together (panel F). All R^2 are
767 significant at $p<0.01$.

Figure. 6. Flake simulation of changing ice cover thickness for each lake separately (A to E) and for an average temperate lake in northern Poland (F) between 2020 and 2099. The model was forced through an ensemble of four climate models (i.e., GFDL-ESM2M, HadGEM, IPSL and MIROC5). Climate changes between 2020 and 2099 were simulated under three relevant GHG concentration trajectories with RCP 2.6 being the most conservative scenario (emissions declining by 2020 and reaching zero by 2100), RCP 6.0 being moderate (emissions peak around 2080, then decline) and RCP 8.5 being the least conservative (emissions continue to rise throughout the 21st century). Shaded area around means for each prediction represent a standard error in prediction for each individual lake (A-E, as a range between the four climate models). Shaded areas around mean in panel F (All lakes) represent variability between individual lakes in the region.

Figure 7. Flake simulation of changing ice cover duration for each temperate lake separately (A to E) and for an average temperate lake in northern Poland (F) between 2020 and 2099. The model was forced with an ensemble of four climate projection models (GFDL-ESM2M, HadGEM, IPSL and MIROC5). Climate changes between 2020 and 2099 were simulated under three relevant GHG concentration trajectories with RCP 2.6 being the most conservative scenario (emissions declining by 2020 and reaching zero by 2100), RCP 6.0 being moderate (emissions peak around 2080, then decline) and RCP 8.5 being the least conservative (emissions continue to rise throughout the 21st century). Shaded area around means for each prediction represent a standard error in prediction for each individual lake (A-E, as a range between the four climate models). Shaded areas around mean in panel F (All lakes) represent variability between individual lakes in the region.

Figures:



Fig. 1.


15°E
23 E
54 N
Baltic Sea
A
B
C
D
E
100 km
Lake monitoring station
Weather station (continuous record 1951-2017)
Lake A (Gardno)
1m
3m
Lake B (Charzykowskie)
5m
30m
20m
Lake C - Jeziorak
5m
Lake D (Mikolajskie)
30m
25m
15m
5m
1km
Lake E (Studzieniczne)
20m
35m
5m
15m

796

797 **Fig. 2.**


A
B
C
D
E
F
Wind
Air T
Rain
R=0.18
R=0.33
R=-0.02
R=0.05
R=0.39
R=0.36
R=-0.23
R=0.35
R=0.14
R=-0.30
R=0.35
R=0.14
R=-0.56
R=0.34
R=0.13
R=-0.12
R=0.30
R=0.15
Temperature °C
Wind (ms-1)
Rainfall (mm month-1)
1960
1970
1980
1990
2000
2010

Fig. 3.


R=-0.30
R=-0.24
Thickness
Duration
A
R=-0.42
R=-0.32
D
R=-0.27
R=-0.27
B
R=-0.46
R=-0.40
E
R=-0.37
R=-0.28
C
R=-0.34
R=-0.27
F
Thickness (cm)
Duration (days)
0
40
80
120
160
1960
1970
1980
1990
2000
2010

Fig. 4.

804



805
806 **Fig. 5.**
807


RPC 2.6
RPC 6.0
RPC 8.5
Ice Cover duration (days)
A
B
C
D
E
F
150
100
50
0
2020
2040
2060
2080
2100

Fig. 6.


RPC 2.6
RPC 6.0
RPC 8.5
Ice Cover Thickness (cm)
A
B
C
D
E
F
80
40
0
2020
2040
2060
2080
2100

Fig. 7.