

School environments and obesity: a systematic review of interventions and policies among school-age students in Latin America and the Caribbean

Article

Accepted Version

Vega-Salas, M. J., Murray, C., Nunes, R. ORCID: <https://orcid.org/0000-0003-0829-4130>, Hidalgo-Arestegui, A., Curi-Quinto, K., Penny, M. E., Cueto, S., Lovegrove, J. A. ORCID: <https://orcid.org/0000-0001-7633-9455>, Sanchez, A. and Vimalleswaran, K. S. ORCID: <https://orcid.org/0000-0002-8485-8930> (2023) School environments and obesity: a systematic review of interventions and policies among school-age students in Latin America and the Caribbean. *International Journal of Obesity*, 47. pp. 5-16. ISSN 1476-5497 doi: 10.1038/s41366-022-01226-9 Available at <https://centaur.reading.ac.uk/108349/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/s41366-022-01226-9>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

School environments and obesity: A systematic review of interventions and policies among school age students in Latin America and the Caribbean

María Jesús Vega-Salas^{1,2*}, Claudia Murray^{3*}, Richard Nunes³, Alessandra Hidalgo-Arestegui², Katherine Curi-Quinto⁴, Mary E. Penny⁴, Santiago Cueto^{5,6}, Julie Anne Lovegrove^{2,7}, Alan Sánchez⁵, Karani Santhanakrishnan Vimalleswaran^{2,7}

**Corresponding authors:* muvega@uc.cl / c.b.murray@henley.reading.ac.uk

Affiliations

¹ Carrera de Nutrición y Dietética, Departamento de Ciencias de la Salud, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago 7820436, Chile

² Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, University of Reading, Whiteknights, Reading RG6 6DZ, UK

³ Department of Real Estate and Planning, Henley Business School, University of Reading, Reading RG6 6UD, UK

⁴ Instituto de Investigación Nutricional (IIN), Lima 15024, Peru

⁵ Grupo de Análisis para el Desarrollo (GRADE), Lima 15063, Peru

⁶ Departamento de Psicología, Pontificia Universidad Católica del Peru, Lima 15088, Peru

⁷ Institute for Food, Nutrition and Health (IFNH), University of Reading RG6 6AH, UK

Competing Interests

The authors declare no competing interests.

Background: The rapid rise in obesity rates among schoolchildren in Latin America and the Caribbean (LAC) could have a direct impact on the region's physical and mental health, disability, and mortality. This review presents the available interventions likely to reduce, mitigate and/or prevent obesity among schoolchildren in LAC by modifying the food and built environments within and around schools.

Methods: Two independent reviewers searched five databases: MEDLINE, Web of Science, Cochrane Library, Scopus and LILACS for peer-reviewed literature published since 1st January 2000 to September 2021; searching and screening prospective studies published in English, Spanish and Portuguese. This was followed by data extraction and quality assessment using the Cochrane risk-of-bias tool (RoB 2) and the Risk of Bias in Non-Randomized Studies of Interventions (ROBINS-I), adopting also the PRISMA-2020 guidelines. Due to the heterogeneity of the intervention's characteristics and obesity-related measurements across studies, a narrative synthesis was conducted.

Results: 1 342 research papers were screened, and nine studies were included; four in Mexico, and one each in Argentina, Brazil, Chile, Colombia, and Ecuador. Four studies reported strategies for modifying food provision; four other targeted the built environment, (modifying school premises and providing materials for physical activity); a final study included both food and built environment intervention components. Overall, two studies reported that the intervention was significantly associated with a lower increase over time in BMI/obesity in the intervention against the control group. The remaining studies were non-significant.

Conclusions: Data suggests school environmental interventions, complementing nutritional and physical education, can contribute to reduce incremental childhood obesity trends. However, evidence of the extent to which food and built environment components factor into obesogenic environments, within and around school grounds is inconclusive. Insufficient data hindered any urban/rural comparisons. Further school environmental intervention studies to inform policies for preventing/reducing childhood obesity in LAC are needed.

Introduction

Globally, childhood overweight and obesity rates has increased substantially over recent decades (1). In Latin America and the Caribbean (LAC), three out of ten children aged 5 to 19 years-old are living with overweight or obesity (2). The consequences of childhood obesity have been well studied and include detrimental health (3), cognitive development and educational attainment (4), and increased risk of developing cardiovascular diseases and obesity in adulthood (5,6). The rapid nutritional transition in the LAC region due to urbanization, economic growth and transformation of broad food systems (7)(8), has had a direct effect on the rising childhood obesity rates. Dietary changes, including higher intakes of energy-dense and low-nutrient-density foods such as sugar-sweetened beverages as well as the lower intakes of vegetables and legumes, and higher physical inactivity and sedentary behaviours (SB) among children and adolescents in LAC have contributed to the rapid increase in obesity and overweight among children and adolescents (9).

Obesogenic environments, defined as “the sum of influences that the surroundings, opportunities, or conditions of life have on promoting obesity in individuals or populations” (10), have impacted children and adults across the world. Previous systematic reviews have focused primarily on assessing the association between the neighbourhood food and built environment (BE), and adiposity and/or weight status among children and adults (11–14). However, as children spend much of their weekday time at schools, and a large proportion of their energy intake and expenditure occurs in this setting (15); more information is needed about the role of schools in childhood obesity. This is the context for our LAC-focused systematic review that provides a valuable contribution, particularly given that several studies suggest that developing interventions at the school-level can contribute to prevention and/or reduction in overweight and/or obesity among children and adolescents (16,17).

School-based interventions have mostly focused on improving the nutritional education curriculum by delivering workshops and information (booklets, pamphlets, posters) for improving dietary behaviours, and increasing physical activity (PA) and/or reduce sedentary behaviours (SB) by

modifying physical education (PE) sessions (18). Several systematic reviews including mostly high-income countries, have reported inconsistent results of the effectiveness of only educational interventions at preventing increases in body weight status (19–21), but some reductions in adiposity or body composition measurements have been reported (18,22–24). Interventions combining diet and PA components, targeting the school and home settings and with longer follow-up, tend to be more successful in preventing or managing weight gain, compared to single component or setting and with a shorter intervention length (19,23,25). Most of the reviews assessing the effectiveness of school-based interventions do not analyse the results according to school level (22,24). A large review separated results between preschool and school-based (primary to secondary school) interventions, however, few studies were conducted among preschool settings to provide any conclusion (25). Two reviews including mostly primary school-aged children found some positive evidence for educational interventions at reducing but not preventing childhood obesity (18,23).

Systematic reviews focusing on school environments are more limited compared with those focusing only on educational components. A recent systematic review and meta-analysis assessed the effectiveness of the school food environment for preventing childhood obesity (26). Results showed that interventions including a food environment component had a significant and meaningful effect on adiposity (BMI z-score). This review included worldwide intervention studies published in English, identifying only one conducted in a LAC country and used a broad definition of food environment, including social marketing and changes to the schools' dietary guidelines, together with interventions targeting the food provision and the nutritional composition of food available at schools. A previous review assessing only isolated school food environment interventions (regulations and food provision) in the US and UK, concluded that the two interventions included, were successful in preventing increases in BMI in the treatment group (27).

The different definitions of school food environments provided by previous reviews included all food and drink available to students within the school (27), and all information influencing food choice

and physical aspects of the food environment, such as availability and accessibility of food within spaces, infrastructure and conditions within or around schools (26). Our review uses the International Network for Food and Obesity/non-communicable diseases Research, Monitoring and Action Support (INFORMAS) framework (28). We also identified dimensions from the WHO School Policy Framework (29) to define the BE within and around the educational premises affecting PA and/or SB to prevent/reduce childhood overweight or obesity. Both frameworks provide a comprehensive and internationally agreed definition for understanding the influence of school environments in childhood obesity. Furthermore, interventions targeting specific aspects of school's food and BE can provide low-cost and easily scaled-up strategies for tackling childhood obesity (27). Making our review particularly relevant for policymakers looking to capitalise on evidence from already available intervention studies.

To this end, our aim is to systematically assess the effectiveness of interventions and policies targeting the school environments for preventing/reducing overweight or obesity among schoolchildren in LAC. In particular, we aim to answer the question: Are school environment interventions/policies effective in the reduction/prevention of obesity and/or overweight among school-age students from LAC? When available, effectiveness will be compared according to the environmental intervention type (food and/or BE), intervention length, and participant's gender and age groups.

Methods

The protocol for this systematic review was registered in PROSPERO ([CRD42021285247](#)), and we followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (30) (Figure 1 and Supplementary Material – Table 1).

Eligibility

Given our focus in LAC, peer reviewed literature published in English, Spanish and Portuguese, from 1st January 2000 to September 2021 were eligible for inclusion. Prospective studies, including interventional study designs containing randomised/non-randomised controlled trials (RCTs and non-

RCTs) and, cohort studies comparing changes in overweight and/or obesity measurements, after a school environment intervention/policy had been implemented, were included.

Search strategy

The team conducted searches in duplicate in five electronic databases: MEDLINE (via PubMed), Web of Science, Cochrane Library, Scopus and the Latin American and Caribbean Health Sciences Literature (LILACS). Search terms and strategies for each database are in Supplementary Material – Tables 2-7. Retrieved reports were stored in EPPI-Reviewer (31) and duplicates identified and excluded. We hand-searched relevant systematic reviews' references and included research papers to identify and incorporate relevant additional studies.

Screening, data extraction and quality assessment

MJV-S and AH-A, both fluent in English and Spanish as well as a good level of Portuguese, conducted title and abstract screening and full-text selection in duplicate. They also pilot tested the first 200 titles and abstracts, obtaining a moderate inter-rate agreement between reviewers (Kappa=0.53) (32,33). Clarifications were made to the inclusion criteria with the whole team and the remaining title and abstract screening completed, obtaining an excellent agreement rate (Kappa=0.74). All discrepancies and full texts in Portuguese were discussed with a third reviewer fluent in this language (RN).

Data extraction was performed independently (MJV-S and AH-A) in EPPI-Reviewer using a piloted coding tool and included the following data: publication details (authors, title, journal, year of publication), study details (study design, RCT characteristics (grouping, randomization, allocation), sampling method, country, school setting, school area (urban/rural), data collection date (baseline and follow-up)), participant information (age, school level, gender/sex, ethnicity, and socioeconomic characteristics, number of participants at baseline and follow-up), intervention details (type of intervention, components, duration, theory), outcome data (measurement type, data collection tool, baseline and follow-up measurements), and effectiveness of intervention. Authors from five studies

were contacted for clarifications and one sent the required information (34,35). When results were presented in plots only, the software Plot Digitizer was used for extracting data (36).

Study quality assessment was undertaken independently (MJV-S and AH-A) by using the Cochrane Risk of Bias tool for cluster-RCT (RoB 2 C-RCT) (37,38), and the Risk of Bias in Non-Randomized Studies of Interventions (ROBINS-I) (39) for non-randomised trials. Studies were graded as low, unclear or high risk of bias. For C-RCTs, six domains were assessed: randomization, timing of identification and recruitment of participants, deviations from intended interventions, missing outcome data, measurement of outcomes and selection of the reported result. For non-RCTs, seven domains were assessed: confounding, selection of participants, classifications of interventions, deviations from intended interventions, missing data, measurement of outcomes and selection of the reported result. Risk of bias assessment by domains for each individual studies is then presented in plots (40).

Types of interventions

All interventions, including the introduction of policies, and/or regulations aiming at modifying obesity/overweight by changing food and/or BE within and around the schools were included. Food environment dimensions were defined by the INFORMAS framework (28): food composition, labelling, marketing, provision, retail, prices and, trade and investments. These dimensions can influence population health, diet and body weight, and can be modified by public and private sector policies. Additionally, we used the WHO School Policy Framework (29) to define two dimensions of the BE: educational buildings and facilities, and walking and cycling infrastructure from and to the educational establishment. Studies assessing interventions at the close proximity to schools were included if conducted within one-mile radius around the perimeter of the educational establishment. Interventions regardless length of follow-up were included. To avoid duplication of data analysis, only the most recent follow-up time including the population relevant to this review was included in the results.

Outcomes

All kind of overweight and/or obesity measurements, including those derived from weight and height (e.g., body mass index -BMI-, standard deviation scores -SDS-, Z-score, prevalence of overweight and obesity, ponderal index); waist circumference and body fat (e.g., body fat percentage, intra-abdominal fat, subcutaneous fat, visceral fat, skin-fold thickness), were included.

Data analysis

We performed a narrative synthesis containing the summary of findings over the effect of interventions on obesity-related measurements, reporting effectiveness of interventions either as mean difference, risk ratio or odds ratio, accordingly to the type of measurement reported in each individual study. We summarised data according to the intervention components reported by each study, classifying it either as a food or, a built and physical environmental intervention. Due to the large heterogeneity in intervention components and multiple outcomes measured across studies, a meta-analysis was not feasible.

Results

Figure 1 shows the PRISMA 2020 flow diagram (30) used for the process of the study selection. The search strategies retrieved 1 329 unique titles and abstracts, and 13 records were added from searching the reference list of relevant reviews and of the included research papers. In total, we assessed 40 full texts for eligibility and nine studies were included. One study was conducted in 2005 (41), another in 2008 (42), and the remaining seven were conducted after 2010. Four studies were in Mexico, while individual studies were in Argentina, Brazil, Chile, Colombia, and Ecuador. Seven studies included girls and boys from primary education, and two included adolescents from lower secondary education (34,43), classified according the International Standard Classification of Education (44). Sample sizes at baseline varied from 168 to 2 682 children, and 120 to 1 224 at follow-up.

Seven studies used a cluster-randomised controlled trial (C-RCT) design, and another two a longitudinal quasi-experimental design (LQE). Clusters selected (schools) varied from 1 per intervention and 1 per control, to 30 schools in each group. All studies compared changes in control and treatment groups between baseline and follow-up. Follow-up measurements varied from ten weeks to three years. Table 1 summarises the study characteristics of the seven studies.

In terms of bias, six out of the seven C-RCT presented a low risk of bias, and one presented some concerns due to the reporting of outcomes. The two LQE studies presented serious concerns related to baseline and time-varying confounding. Figures 2 and 3 summarise the risk of bias assessment.

Overall, four studies reported intervention components for modifying the food environments and four studies, the BE (Table 2). Only one study reported components for both food and BE interventions (45). Obesity-related outcomes were heterogeneous across the studies, including reports of BMI, BMI z-score and, overweight and/or obesity prevalence. The following sections present a detailed description of the design and results of included interventions targeting a) the food environment b) BEs, and c) a combination of food and BEs.

Food environments

Four studies included intervention components targeting the food environments of the schools. Following the INFORMAS dimensions, four studies targeted the food provision by increasing the availability of healthy products, while one limited the sales of high-energy and unhealthy foods (45). Only the study by Ramírez-López et al., (2005)(41) assessed one component intervention, targeting the food composition of free school breakfasts, while the remaining three studies had several other intervention components, including strategies around nutritional and PE. Some studies reported more than one obesity-related measurement; BMI outcomes and BMI z-score were both reported together by two studies, body fat percentage and fat-free body mass was presented in one study, and overweight and obesity prevalence was reported by one study. One study was conducted in urban areas (Rosario), one in a rural setting (Metropolitan region of Santiago), one in a semi-rural (State of

Mexico), and a final one comprising a large region (the State of Sonora) and including both rural and urban contexts.

The cluster RCT by Alvirde-García (2013) (46) included students aged 9-10 years-old at baseline attending five semi-rural schools from the State of Mexico. The food provision component included a modification to the food items offered in school canteens by increasing the availability of fruits and vegetables and products low in saturated fat and sugar. Additionally, the intervention included a nutritional and PA education component, delivering workshops with parents, school staff and school vendors, and booklets for students to complement their school curriculum. Results showed a similar (average) increase of BMI over time in the treatment group compared to the control group during the first two years, but a significantly lower rate of increase in (average) BMI among those in the treatment group, compared to the control group, for the third year of the intervention (1.6 ± 1.9 vs. 1.9 ± 1.7 Kg/m², $p < 0.01$). Despite both groups decreasing their energy intakes over time, on the third year this decrease was significantly higher among the intervention or treatment group compared with the control one (-756 kcal/d, $p < 0.05$).

The longitudinal quasi-experimental study by González et al., (2014) (47) included preschool to 8th grade students (4-15 years-old) from six schools located in rural areas of the Metropolitan region of Santiago, Chile. All students from five schools located in the same municipality received the year-long intervention, whereas students from one school at a different municipality were assigned to a control group. Students from the treatment group received fruits 3 times-per-week and a fruit basket was given to the family at the end of the year. This intervention also included workshops with nutritional education material for the students, their parents, and teachers. At the end of the year, results did not show any significant change in body weight status among participants in the control or treatment groups. However, the intervention was successful in increasing frequency of daily intakes of fruits, vegetables, dairy products, pulses and fish, but was ineffective for reducing consumption of unhealthy foods such as chips, hotdogs and pizza. Authors highlight the lack of

increase in overweight or obesity status in treatment groups, considering the higher intake of healthy products that might contribute to increasing energy intakes.

The year-long quasi-experimental study by Ramírez-López et al., (2005) (41) included 1st to 5th grade students (6-10 years-old) from urban and rural areas in the state of Sonora, Mexico. The intervention assessed the effect of a national-and-state-funded free school breakfast (FSB) programme on obesity, body composition and cardiovascular risk, compared to non-beneficiaries'. This was the only study assessing one intervention component (i.e., provision of a free school breakfast). Results showed that FSB beneficiaries did not differ in overweight or obesity prevalence, BMI, or in body fat percentage, to those in the control group at the end of the 9-month intervention. Similarly, no major differences between groups were reported for total cholesterol, triglycerides, and glucose.

The cluster RCT by Rausch Herscovici et al., (2013) (42) included students aged 9-11 years-old attending six schools from urban areas in Rosario, Argentina. The food provision intervention modified the school canteen options to include healthy food items (fruits, orange juice and low-sugar cereal). Additionally, the intervention included three nutritional and PE workshops for children and one for parents. Results after 6 months showed no significant difference in BMI between the intervention and control groups. However, girls in the experimental group (not the boys) increased consumption of some healthy foods targeted by the intervention (skim milk and orange juice), compared to their control group counterparts.

Built environments

Among the four interventions targeting the BE, one study intervened the school playground, and three studies provided materials for promoting PA within the school premises. Studies could report more than one obesity-related measurements; BMI was reported by one study, BMI z-score was reported in three studies, and overweight and obesity prevalence was reported in another study. Four studies included schools located in urban areas (in the secondary cities of Cuenca and Fortaleza, and the capital city of Bogota), while only one study covered both rural and urban areas (State of Sonora).

The 3-year cluster RCT by Andrade et al., (2014) (43) included 12 and 13 year-old adolescents attending 20 schools from urban areas of Cuenca, Ecuador, and involved a BE intervention with a walking trail drawn on the playground in the second year of the intervention. Other components included nutritional and physical education (PE) materials (booklets and posters), workshops for adolescents and their parents, and the organisation of social events with famous athletes. After 3 years of intervention, no effects were reported for mean BMI z-score or prevalence of overweight between control and treatment groups. However, students in the treatment group showed a positive effect on physical fitness parameters (vertical jump and speed shuttle run) and a higher percentage met the PA recommendations (60 min of MVPA/day), compared to students allocated to the control groups (6 vs. 18 percentage points, $p < 0.01$).

The 4-month cluster RCT by Barbosa Filho et al., (2017) (34,35) involved 11 to 13 year-old adolescents in six schools from urban areas of Fortaleza, Brazil. The BE intervention offered space and PA equipment (balls, rackets, mini courts) to promote PA during free time. Other components involved health and PE training and materials (booklets, interactive media, posters) for teachers to include in the school curriculum, pamphlets to students and parents. After four months, no significant effects were reported for BMI, overweight or obesity prevalence. However, the intervention was successful in increasing MVPA time, number of PA, and time spent in PA games per week (control= -75.15, -0.25, -28.30; intervention= 127.92 0.63, 92.01, respectively).

The cluster RCT by Gutiérrez-Martínez et al., (2018) (48) included 10-year-old students in three schools (two treated and one control) from urban areas in Bogotá, Colombia. Both treatment groups received PA equipment (ribbons, balls, hoops, stairs, parachute, and mats) to support PA during recess. Additionally, a PE instructor delivered 30 standardised PA activities lasting 20' each throughout the 10-week intervention period. Additionally, participants in one of the treatment groups received daily SMS messages to promote extra-curricular PA and healthy nutrition. Results suggested there were no effects on BMI z-score or body fat percentage over the 10-week intervention period.

Nevertheless, the intervention was successful in increasing MVPA and reducing SB minutes among participants in the treatment groups compared to those in the control one.

Finally, the 6-month cluster RCT by Shamah-Levy et al., (2012) (49) included 10 to 12 year-old students in six schools from both urban and rural areas in the State of Mexico. The treatment group received PA equipment (balls, ropes, and hoops) to support PA during recess over a 6-month intervention period. Other components included nutrition and PA education through workshops and materials (booklets, puppet show, advertising, banners) for students, parents, and school staff. Canteen personnel attended workshops aimed at promoting the daily sales of fruit, vegetables, and water. Results suggested a small but significant reduction in the probability of students in the treatment group to shift from the overweight to the obesity category after 6 months, compared to the ones in the control group (OR= 0.68; $p = 0.01$). However, no significant differences were reported for both groups (control and intervention) in the probabilities of shifting from the normal to overweight category after the intervention period. Overall, the intervention was relatively effective in maintaining BMI among children in the treatment group.

Food and built environments

The 18-month intervention reported by Safdie et al., (2013) (45) involved 4th and 5th grade children (9-10 years-old at baseline) attending 27 schools from urban areas of Mexico City. This study is the only one including food as well as BE strategies, among other intervention components. Additionally, the strategy was implemented in two treatment groups, basic and plus, with the latter having all the same activities than the first, plus extra components implemented with additional financial investment and human resources. This cluster RCT mixed different strategies, including the modification over the food provision in school canteens by limiting the availability of sugar-sweetened beverages (SSB) and the sales of energy-dense foods at the school canteens during the two-years for the plus group, and only during the second year for the basic one. It also included improvement of the school premises and provision of sports equipment for promoting the use of PA areas for two years in two different

treatment groups (basic and plus). Games and sports courts were drawn on the ground, and each school received PA equipment (balls, ropes, nets, and elastic bands) to support PE classes and PA during recess and free time. Other components included promoting the availability of healthy food (fruits, vegetables, and non-fried dishes) and beverages (water) within school premises, reducing the number of eating opportunities, while providing nutritional and PA education by delivering workshops and pamphlets to students, parents, school staff and vendors. The intervention also included strategies for promoting PA during recess, among other activities. A small, yet non-significant reduction in the prevalence of overweight and obesity was reported for children from control and interventions groups (basic and plus) during the first year (19.5 vs 17; 11.9 vs 11.3; 12 vs 11.2%, respectively). Conversely, a slight increase in the prevalence was reported at the beginning of year 2 for control and basic treatment groups, but not for the plus group (17.9, 12.1, 10.7%, respectively). Only children in the basic treatment groups reported a small but non-significant reduction in overweight and obesity prevalence during the second year (12.1% and 10.9%, respectively). In contrast, a small but significant BMI reduction was reported for control and plus groups in year 1 (19.9 to 18.4, and 20 to 18.5%, respectively). However, an inverse direction was reported for all in year 2, with small but non-significant increases in BMI across all groups (control =18.9 to 19.1; basic= 20.1 to 20.4; plus= 18.7 to 19%). Therefore, the small-in-magnitude changes presented in overweight and obesity prevalence and BMI across the intervention period cannot be associated with the study intervention as similar changes were reported in control and interventions groups between baseline and follow-up periods. Yet, the intervention was effective at increasing intakes in recommended food and beverages and decreasing unhealthy ones, together with significant increases in PA (e.g., increases in steps taken by the students), among both treatment groups, compared to the control.

Discussion

Our review found only nine studies assessing school-based interventions including components for modifying the food and BE within and outside primary and secondary schools in LAC. From this pool, we are unable to conclude that children's exposure to environmental interventions resulted in changes to obesity-related measurements. Albeit, two studies (46,49) showed some results related to the prevention of obesity. Both were implemented in the State of Mexico, the first one in semi-rural areas (46) while the second one targeting both rural and urban areas (49). It is not clear why this geographical concentration appears in our results; it could be speculated that the region has a higher obesity rate compared to other regions in our review –i.e., it already starts from a high rate of obesity and thus impact is easily detected. Notwithstanding, with such differences in the interventions' design, this cannot be evidenced and therefore must remain as a hypothesis for further studies.

The remaining seven studies did not present any significant changes in overweight or obesity-related measurements between control and treatment groups. However, all eight studies assessing intermediate outcomes contributing to prevent obesity on the long term reported some positive results, such as decreases in energy intakes (46) and in sedentary behaviour (48), increases in fruit and vegetable intake (47), healthy products (42,45), physical fitness (43), MVPA minutes (48,53), and steps taken (45). Our findings are similar to previous intervention studies reviews from the Global North, reporting improved dietary behaviours and increasing PA albeit inconclusive regarding the effects over obesity-related measures (54–56). Notwithstanding, a recent review and meta-analysis including studies worldwide and using a wider definition of school food environments reported a meaningful effect of interventions to reduce adiposity (-0.12 , 95% CI: $0.15-0.10$) (26). Overall, all but two studies were classified as showing a low risk of bias, with the remaining two as with moderate risk (41,46), and all but three studies (43,45,46) had 1-year or shorter follow-up measurements, which could have weakened or biased our results. However, these studies are examples of the relatively few

number of studies assessing interventions in LAC, providing valuable information concerning the study design and methodological implications for future research teams.

In terms of scientific research and evidence mapping, our systematic review revealed the low number of peer-review articles assessing the effectiveness of food and BE interventions in schools for preventing/reducing childhood overweight and obesity in LAC. Previous reviews (57,58) have primarily encountered interventions relying on educational components (e.g., nutritional education and modifications to PE sessions), and not environmental components. We also identified helpful methodological implications for future interventions in the region, for example, the need for a longer follow-up (beyond a 1-year horizon), and targeting both, the food and BEs. They should also assess mediating outcomes (changes in dietary and PA behaviours) and distal ones (obesity-related measures) when planning intervention strategies.

Our review has uncovered five studies targeting the BE, all within school boundaries, therefore not covering the 1-mile radius from the school as per our protocol. This is disappointing, particularly considering the positive impact that active commuting has in preventing obesity in schools (59–61). Studies promoting BE interventions outside schools, such as active commuting, requires organising multiple stakeholders (e.g., schools, councils, policymakers, and/or researchers), which might need more funding (62). Considering shortage of funding for research and development in LAC, with only 0.67% of its GPD allocated to it (63) and mostly from the public sector (64), interventions connecting different stakeholders and with a longer follow-up can face financial barriers. More research investment from governments and/or other funders could foster multi-stakeholder collaboration and design ambitious interventions, at the neighbourhood scale.

Moreover, most of the studies included here targeted urban areas, and even those targeting rural or semirural areas were in large metropolitan regions (Santiago de Chile and State of Mexico). Considering that food provision in rural areas in LAC is generally more expensive than in urban locations (due to transport and logistic costs) (65), we hoped to find interventions conducted in more

distinctive urban and rural settings. Expecting therefore to find contrasting results based on locations but the lack of published research on more typical rural landscapes has hampered any conclusive findings. However, this research gap does highlight the need for more interventions targeting the built and food environments in rural areas for preventing and reducing childhood obesity.

Our review has made a positive contribution to science and policymaking by updating the available evidence, even though included prospective studies only captured cluster-RCTs and LQE interventions excluding pre-and post-policy outcome evaluations related to childhood obesity. Only one LQE study in our pool assessed a state-wide school feeding programme, showing no difference between those receiving a free-school breakfast in any obesity-related measurement to those who did not. (41). It is in this area where our review also highlights a lack of policy evaluation studies reporting obesity-related outcomes. Indeed, 13 LAC countries have regulated the sale of food and beverages in schools (66), and four countries including Chile, Costa Rica, Ecuador and Uruguay, have implemented national policies aiming to restrict food marketing of unhealthy foods within school premises (67). Only two countries, Chile and Mexico (66), have performed policy evaluations regarding restrictions of unhealthy product sales in schools, reporting positive results for reducing their availability in school kiosks in Chile (68) and for decreasing energy intake in children who only consumed food purchased at school in Mexico (69).

The case of Chile is a unique example within the region for implementing, in 2016, a mandatory and comprehensive policy for reducing consumption of unhealthy products, and reducing and preventing obesity by including mandatory front-of-package warning labels, limiting advertising, and prohibiting school sales of products high in calories, sodium, sugar, or saturated fat (70). Recent policy evaluations have reported positive outcomes for reducing the consumption and exposure to television advertising of unhealthy products among pre-school children (71) and for households reducing purchases of unhealthy products (72). However, no peer-reviewed policy evaluation in Chile has yet assessed the effect over obesity-related outcomes. A good example of a pre-and post-evaluation of an

obesity prevention policy and its effectiveness in changing obesity-related measurements is the impact assessment of the sugar-sweetened beverages (SSB) tax in Mexico and its role in decreasing overweight or obesity prevalence among adolescents (73). Considering as stated above that research funding is scarce, future research should test the effectiveness of these policy-related interventions by conducting rigorous RCTs at a small-scale; and use this evidence to decide whether scaling-up is worthwhile. Scientists should exploit the opportunities presented by such policy changes and test their effect on changes in childhood obesity-related outcomes. The outcomes of such pre-and post-evaluations will take time but would at least inform governments if policy fixes are needed.

This lack of peer-reviewed policy evaluations suggests that there could be a disconnect between the scientific community and policymakers. A finding that can be attributed to a potential publication bias within our study based on the exclusion of grey literature (e.g., technical reports). Notwithstanding, the question is, are scientist producing sufficient and adequate evidence for policymakers? Some evidence from studies reviewed here shows positive results in intermediate outcomes, such as reduction of sedentary behaviour and increase in fruit and vegetable intake as reported above. Yet, it seems peer-reviewed studies are not assessing changes in obesity-related measurements before/after policies are implemented and therefore, policymakers do not seem to have the relevant evidence on the effectiveness of policies targeting childhood obesity in LAC. Decision-makers need evaluations of the short-term and long-term impact of childhood obesity prevention policies targeting school environments for reducing/preventing obesity, vis-à-vis the assessment of intermediate obesity determinants.

Considering that several countries in LAC are facing a double burden of obesity and undernutrition(74), this potential disconnect between the scientific community and policymakers is concerning, particularly considering the current COVID-19 pandemic. Indeed, the already large disparities in obesity rates and in behaviours contributing to obesity (diet, PA and sedentary behaviour) in LAC (75–78), predominantly affecting economically disadvantaged populations, has

placed a disproportionate burden on these groups during the pandemic (79). Due to disruptions in food supply chains, decreases in income and reductions in PA due to lockdowns (80,81), it is expected that obesity rates across the continent will be impacted. Furthermore, the pandemic also has highlighted this science and policy disconnect, particularly considering the emergency response measures coming from some LAC governments at the start of the pandemic (such as Brazil and Mexico) (82,83). Despite the large and conclusive scientific evidence suggesting effective measures for mitigating contagion (e.g., use of facemasks and social distancing), some countries simply ignored the science. Academics working on other pressing issues such as climate change, are already concluding that scientific evidence is more effective when academics and policymakers engage (84,85). LAC governments must take steps in bridging the science and policy gap, ensuring that policies are independently evaluated and peer-reviewed before upscaling.

Strengths and Limitations

The main methodological limitations arise from the different sources of heterogeneity we encountered among the included studies. We list below the sources and their effect on our review or the studies themselves.

The first source is the high heterogeneity in reporting outcome measurements and measures of error. Studies reported different cut-off points and operationalisations for obesity-related outcomes (e.g., BMI, BMI z-score, overweight and/or obesity prevalence). Some presented results as mean differences; others reported averages or prevalence and others compared the frequency of these changes. Additionally, only two reported straightforward measurements of variability for the effect changes (standard deviations, standard error, or confidence intervals). The second source is the high heterogeneity in study designs, age groups and types of intervention. A third source is the use of multiple intervention components used by the included studies, which might have influenced the lack of conclusive results. Together with modifications to the food and/or, BEs, interventions combined strategies by including nutrition and PA education, and/or changes to PE sessions. These components

are delivered by different strategies, such as providing materials within the school curriculum, presenting workshops for students, parents and school staff, and the use of social marketing strategies (e.g., pamphlets and posters), among others. This multiplicity prevents us from clearly identify if the intervention effects can be attributed to the inclusion of changes to the food or BEs in schools or to other types of intervention strategies. Only one study presented a food environmental strategy in isolation, and therefore, we cannot easily compare the effects of the different strategies. Altogether, these three sources of heterogeneity prevented us from quantitatively pooling data for a meta-analysis. Likewise, the variable duration of interventions may have had an impact on the extent to which obesity-related measurements were affected. Six interventions lasted less than an academic year (<9 months), one lasted two years (18 months) and two lasted more than 3 years (28 months). Despite most of the interventions reporting positive results on some intermediate outcomes (diet, PA and SB), most failed to find any significant difference in measurements of obesity between intervention and control groups. Furthermore, all studies had small sample sizes (i.e., a reduced number of treated and non-treated schools). It is possible that some of these interventions might have been successful but that the effects might not have been large enough to be detected. Future interventions should consider a larger number of schools (based on power size calculations) and longer follow-up periods in their design, ensuring more conclusive findings on long term obesity changes.

Conclusion

This review synthesised, for the first time, the effectiveness of interventions targeting the food and BEs in schools to prevent/reduce childhood obesity in LAC. Due to the high heterogeneity in study design and reporting outcomes, results were inconclusive. However, no study in our review reported a significant increase in BMI or obesity prevalence when interventions included modifications to the food and/or BEs.

In terms of evidence mapping, we revealed the low number of peer-review articles assessing the effectiveness of food and built and school environment interventions for preventing and reducing

489 childhood overweight and obesity in LAC. Furthermore, we also have detected a complete absence
490 of studies assessing the BE outside school buildings, for example encouraging of active school
491 commute.

492 Our conclusion is more concerning, as it points to a lack of policy evaluations from countries that
493 have implemented policies, vis-à-vis a lack of adequate policy-informing evidence in countries where
494 academics are active on obesity-related research, suggesting there is a potential disconnect between
495 science and policymaking. With three out of ten children aged 5 to 19 years-old living with
496 overweight or obesity in LAC countries, further funding to fund studies aiming to prevent and reduce
497 childhood obesity in school settings in the region is needed. Notwithstanding, the production of
498 evidence means little if science and policy operate in silos with little co-production of knowledge to
499 better understand the food and BE factors that underpin LAC's obesogenic environments where
500 children learn, play and grow.

Acknowledgements

We are grateful to all the authors who responded our request for data for inclusion in our review.

Funding

This review was funded by PROCIENCIA (CONCYTEC/FONDECYT) grant number 030-2019, the British Embassy, the MRC grant (MR/S024778/1) and Newton Fund; and the Old Dart Foundation

The funders had no role in the design, collection, analysis, or interpretation of the present study, or in the decision to publish.

Authorship

C.M., R.N. and V.K.S. conceived the review. C.M., R.N., V.K.S. and M.J.V-S. designed the methodology for the systematic review. M.J.V-S. led and conducted the literature search, screening, data extraction, risk of bias assessment, data analysis and interpretation, and writing of the first draft of the manuscript. A.H-A. was the second reviewer and contributed to searches, screening, data extraction, risk of bias assessment, and data analysis. R.N. was involved in resolving disagreements during screening processes. C.M. critically revised and edited the final draft of the manuscript. R.N., A.H-A, K.C-Q., M.E.P., S.C., J.A.L., A.S., and V.K.S., provided comments, revised the manuscript, and approved the final version. M.J.V-S. and C.M. confirm they had full access to the data in the study and final responsibility for the decision to submit for publication.

References

1. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014 Aug 30;384(9945):766–81.
2. Fondo de las Naciones Unidas para la Infancia. El sobrepeso en la niñez: Un llamado para la prevención en América Latina y el Caribe [Internet]. Ciudad de Panamá; 2021 [cited 2021 Oct 27]. Available from: www.unicef.org/lac
3. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002 Aug 10;360(9331):473–82.
4. Segal AB, Huerta MC, Aurino E, Sassi F. The impact of childhood obesity on human capital in high-income countries: A systematic review. *Obes Rev*. 2021 Jan 1;22(1):e13104.
5. Park MH, Falconer C, Viner RM, Kinra S. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obes Rev*. 2012 Nov 1;13(11):985–1000.
6. Singh AS, Mulder C, Twisk JWR, Mechelen W Van, Chinapaw MJM. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*. 2008 Sep 1;9(5):474–88.
7. Popkin BM, Reardon T. Obesity and the food system transformation in Latin America. *Obes Rev*. 2018;19(8):1028–64.
8. Popkin BM, Gordon-Larsen P. The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes*. 2004 Nov 15;28(S3):S2–9.
9. Corvalán C, Garmendia ML, Jones-Smith J, Lutter CK, Miranda JJ, Pedraza LS, et al. Nutrition status of children in Latin America. *Obes Rev*. 2017 Jul;18:7–18.

- 541 10. Swinburn B, Egger G, Raza F. Dissecting Obesogenic Environments: The Development and
542 Application of a Framework for Identifying and Prioritizing Environmental Interventions for
543 Obesity. *Prev Med (Baltim)*. 1999 Dec 1;29(6):563–70.
- 544 11. Townshend T, Lake A. Obesogenic environments: current evidence of the built and food
545 environments: *Perspect Public Health*. 2017 Jan 11;137(1):38–44.
- 546 12. Feng J, Glass TA, Curriero FC, Stewart WF, Schwartz BS. The built environment and obesity:
547 A systematic review of the epidemiologic evidence. *Health Place*. 2010 Mar 1;16(2):175–90.
- 548 13. Mackenbach JD, Rutter H, Compernelle S, Glonti K, Oppert JMM, Charreire H, et al.
549 Obesogenic environments: a systematic review of the association between the physical
550 environment and adult weight status, the SPOTLIGHT project. *BMC Public Health*. 2014 Mar
551 6;14(1):1–15.
- 552 14. Li Y, Luo M, Wu X, Xiao Q, Luo J, Jia P. Grocery store access and childhood obesity: A
553 systematic review and meta-analysis. *Obes Rev*. 2021 Feb 1;22(S1):e12945.
- 554 15. Story M, Kaphingst KM, French S. The Role of Schools in Obesity Prevention. *Futur Child*.
555 2006;16(1):109–42.
- 556 16. Fondo de las Naciones Unidas para la Infancia (UNICEF). El rol de la escuela en la prevención
557 del sobrepeso y la obesidad en estudiantes de América Latina y el Caribe [Internet]. 2021
558 [cited 2021 Oct 29]. Available from: [https://www.unicef.org/lac/media/29016/file/LACRO-](https://www.unicef.org/lac/media/29016/file/LACRO-El-rol-de-la-escuela-en-la-prevencion-del-sobrepeso.pdf)
559 [El-rol-de-la-escuela-en-la-prevencion-del-sobrepeso.pdf](https://www.unicef.org/lac/media/29016/file/LACRO-El-rol-de-la-escuela-en-la-prevencion-del-sobrepeso.pdf)
- 560 17. Foster GD, Sherman S, Borradaile KE, Grundy KM, Veur SS Vander, Nachmani J, et al. A
561 Policy-Based School Intervention to Prevent Overweight and Obesity. *Pediatrics*. 2008 Apr
562 1;121(4):e794–802.
- 563 18. Sbruzzi G, Eibel B, Barbiero SM, Petkowicz RO, Ribeiro RA, Cesa CC, et al. Educational

interventions in childhood obesity: A systematic review with meta-analysis of randomized clinical trials. *Prev Med (Baltim)*. 2013 May 1;56(5):254–64.

19. Wang Y, Cai L, Wu Y, Wilson RF, Weston C, Fawole O, et al. What childhood obesity prevention programmes work? A systematic review and meta-analysis. *Obes Rev*. 2015 Jul 1;16(7):547–65.

20. Kropiski JA, Keckley PH, Jensen GL. School-based Obesity Prevention Programs: An Evidence-based Review. *Obesity*. 2008 May 1;16(5):1009–18.

21. Guerra PH, Silveira JAC da, Salvador EP, da Silveira JAC, Salvador EP. Physical activity and nutrition education at the school environment aimed at preventing childhood obesity: evidence from systematic reviews. *J Pediatr (Rio J)* [Internet]. 2016 Jan 1 [cited 2021 Oct 29];92(1):15–23. Available from:

<http://www.scielo.br/j/jped/a/myYFxpVHnLB5V7HVX9Z7txm/abstract/?lang=en>

22. Lavelle H V., Mackay DF, Pell JP. Systematic review and meta-analysis of school-based interventions to reduce body mass index. *J Public Health (Bangkok)*. 2012 Aug 1;34(3):360–9.

23. Liu Z, Xu HM, Wen LM, Peng YZ, Lin LZ, Zhou S, et al. A systematic review and meta-analysis of the overall effects of school-based obesity prevention interventions and effect differences by intervention components. *Int J Behav Nutr Phys Act*. 2019 Oct 29;16(1):1–12.

24. Sobol-Goldberg S, Rabinowitz J, Gross R. School-based obesity prevention programs: A meta-analysis of randomized controlled trials. *Obesity* [Internet]. 2013 Dec 1 [cited 2021 Nov 1];21(12):2422–8. Available from:

<https://onlinelibrary.wiley.com/doi/full/10.1002/oby.20515>

25. Bleich SN, Vercammen KA, Zatz LY, Frelrier JM, Ebbeling CB, Peeters A. Interventions to

- 587 prevent global childhood overweight and obesity: a systematic review. *Lancet Diabetes*
588 *Endocrinol.* 2018 Apr 1;6(4):332–46.
- 589 26. Pineda E, Bascunan J, Sassi F. Improving the school food environment for the prevention of
590 childhood obesity: What works and what doesn't. *Obes Rev.* 2021 Feb 1;22(2).
- 591 27. Driessen CE, Cameron AJ, Thornton LE, Lai SK, Barnett LM. Effect of changes to the school
592 food environment on eating behaviours and/or body weight in children: a systematic review.
593 *Obes Rev.* 2014 Dec 1;15(12):968–82.
- 594 28. Swinburn B, Sacks G, Vandevijvere S, Kumanyika S, Lobstein T, Neal B, et al. INFORMAS
595 (International Network for Food and Obesity/non-communicable diseases Research,
596 Monitoring and Action Support): overview and key principles. *Obes Rev.* 2013 Oct;14(1,
597 SI):1–12.
- 598 29. World Health Organization. School policy framework: implementation of the WHO global
599 strategy on diet, physical activity and health [Internet]. 2008 [cited 2021 Aug 4]. Available
600 from: <https://apps.who.int/iris/handle/10665/43923>
- 601 30. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The
602 PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*
603 [Internet]. 2021 Mar 29 [cited 2021 Oct 12];372. Available from:
604 <https://www.bmj.com/content/372/bmj.n71>
- 605 31. Thomas J., Brunton J. EPPI-Reviewer: advanced software for systematic reviews, maps and
606 evidence synthesis [Internet]. London: EPPI-Centre Software, UCL Social Research Institute;
607 2020 [cited 2021 Nov 16]. Available from:
608 <https://eppi.ioe.ac.uk/cms/Default.aspx?tabid=1913>
- 609 32. Higgins JPT., Thomas J., Chandler J., Cumpston M., Li T., Page MJ., et al. *Cochrane*

- Handbook for Systematic Reviews of Interventions [Internet]. 2021 [cited 2021 Nov 16]. Report No.: 6.2. Available from: www.training.cochrane.org/handbook
33. Orwin R. Evaluating coding decisions. In: Cooper H, Hedges L, editors. The handbook of research synthesis. New York, NY: Russell Sage Foundation; 1994.
 34. Filho VCB, Da Silva KS, Mota J, Beck C, Da Silva Lopes A. A Physical Activity Intervention for Brazilian Students From Low Human Development Index Areas: A Cluster-Randomized Controlled Trial. *J Phys Act Heal*. 2016 Nov 1;13(11):1174–82.
 35. Barbosa Filho VC, da Silva KS, Mota J, Vieira NFC, Gubert F do A, Lopes A da S. “For whom was it effective?” Moderators of the effect of a school-based intervention on potential physical activity determinants among Brazilian students. *Prev Med (Baltim)*. 2017 Apr 1;97:80–5.
 36. Cochrane Training. Extracting data from figures using software [Internet]. 2016 [cited 2021 Oct 15]. Available from: <https://training.cochrane.org/resource/extracting-data-figures-using-software-webinar>
 37. Eldridge S, Campbell MK, Campbell MJ, Drahota AK, Giraudeau B, Reeves BC, et al. Revised Cochrane risk of bias tool for randomized trials (RoB 2) - Additional considerations for cluster-randomized trials (RoB 2 CRT) [Internet]. 2021 [cited 2021 Oct 26]. Available from: https://drive.google.com/file/d/1yDQtDkrp68_8kJiUdbongK99sx7RFI-/view
 38. Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration’s tool for assessing risk of bias in randomised trials. *BMJ*. 2011 Oct 18;343(7829).
 39. Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016 Oct

- 633 12;355:i4919.
- 634 40. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny
635 web app for visualizing risk-of-bias assessments. *Res Synth Methods*. 2021 Jan 1;12(1):55–
636 61.
- 637 41. Ramírez-López E, Grijalva-Haro MI, Valencia ME, Ponce JA, Artalejo E. Impacto de un
638 programa de desayunos escolares en la prevalencia de obesidad y factores de riesgo
639 cardiovascular en niños sonorenses. *Salud Publica Mex*. 2005;47(2):126–33.
- 640 42. Rausch Herscovici C, Kovalskys I, Jose De Gregorio M. Gender differences and a school-
641 based obesity prevention program in Argentina: a randomized trial. *Rev Panam SALUD*
642 *PUBLICA-PAN Am J PUBLIC Heal*. 2013;34(2):75–82.
- 643 43. Andrade S, Lachat C, Ochoa-Aviles A, Verstraeten R, Huybregts L, Roberfroid D, et al. A
644 school-based intervention improves physical fitness in Ecuadorian adolescents: a cluster-
645 randomized controlled trial. *Int J Behav Nutr Phys Act*. 2014 Dec 10;11(153):1–17.
- 646 44. UNESCO Institute for Statistics. International Standard Classification of Education - ISCED
647 2011 [Internet]. Montreal, Canada; 2012 [cited 2021 Dec 2]. Available from:
648 <http://www.uis.unesco.org>
- 649 45. Safdie M, Jennings-Aburto N, Lévesque L, Janssen I, Campirano-Núñez F, López-Olmedo N,
650 et al. Impact of a school-based intervention program on obesity risk factors in Mexican
651 children. *Salud Publica Mex*. 2013;55 Suppl 3:374–87.
- 652 46. Alvirde-García U, Rodríguez-Guerrero AJ, Henao-Morán S, Gómez-Pérez FJ, Aguilar-
653 Salinas CA. Resultados de un programa comunitario de intervención en el estilo de vida en
654 niños. *Salud Publica Mex*. 2013;55(supl 3):S406–14.
- 655 47. González G CG, Zacarías H I, Domper R A, Fonseca M L, Lera M L, Vio del R F. Evaluación

- 656 de un programa de entrega de frutas con educación nutricional en escuelas públicas rurales de
657 la Región Metropolitana, Chile. *Rev Chil Nutr.* 2014 Sep 1;41(3):228–35.
- 658 48. Gutiérrez-Martínez L, Martínez RG, González SA, Bolívar MA, Estupiñan OV, Sarmiento
659 OL. Effects of a strategy for the promotion of physical activity in students from Bogotá. *Rev*
660 *Saude Publica.* 2018 Jul 26;52.
- 661 49. Shamah Levy T, Morales Ruán C, Amaya Castellanos C, Salazar Coronel A, Jiménez Aguilar
662 A, Méndez Gómez Humarán I. Effectiveness of a diet and physical activity promotion strategy
663 on the prevention of obesity in Mexican school children. *BMC Public Heal* 2012 121. 2012
664 Mar 1;12(1):1–13.
- 665 50. Barlow SE, Committee and the E. Expert Committee Recommendations Regarding the
666 Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity:
667 Summary Report. *Pediatrics.* 2007 Dec 1;120(Supplement_4):S164–92.
- 668 51. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child
669 overweight and obesity worldwide: international survey. *BMJ.* 2000 May 6;320(7244):1240.
- 670 52. WHO MULTICENTRE GROWTH REFERENCE STUDY GROUP, de Onis M. WHO Child
671 Growth Standards based on length/height, weight and age. *Acta Paediatr.* 2006
672 Apr;450(S450):76–85.
- 673 53. Barbosa Filho VC, Minatto G, Mota J, Silva KS, de Campos W, Lopes A da S. Promoting
674 physical activity for children and adolescents in low- and middle-income countries: An
675 umbrella systematic review. A review on promoting physical activity in LMIC. Vol. 88,
676 Preventive Medicine. Academic Press Inc.; 2016. p. 115–26.
- 677 54. Micha R, Karageorgou D, Bakogianni I, Trichia E, Whitsel LP, Story M, et al. Effectiveness
678 of school food environment policies on children’s dietary behaviors: A systematic review and

679 meta-analysis. PLoS One. 2018 Mar 1;13(3):e0194555.

680 55. Williams AJ, Wyatt KM, Hurst AJ, Williams CA. A systematic review of associations
681 between the primary school built environment and childhood overweight and obesity. Health
682 Place. 2012 May 1;18(3):504–14.

683 56. Davison KK, Lawson CT. Do attributes in the physical environment influence children's
684 physical activity? A review of the literature. Int J Behav Nutr Phys Act. 2006 Jul 27;3(1):1–
685 17.

686 57. Lobelo F, Quevedo IG de, Holub CK, Nagle BJ, Arredondo EM, Barquera S, et al. School-
687 Based Programs Aimed at the Prevention and Treatment of Obesity: Evidence-Based
688 Interventions for Youth in Latin America. J Sch Health. 2013 Sep 1;83(9):668–77.

689 58. Mancipe Navarrete JA, Garcia Villamil SS, Correa Bautista JE, Meneses-Echávez JF,
690 González-Jiménez E, Schmidt-Riovalle J. Efectividad de las intervenciones educativas
691 realizadas en América Latina para la prevención del sobrepeso y obesidad infantil en niños
692 escolares de 6 a 17 años: una revisión sistemática. Nutr Hosp. 2015;31(1):102–14.

693 59. Lubans DR, Boreham CA, Kelly P, Foster CE. The relationship between active travel to school
694 and health-related fitness in children and adolescents: A systematic review. Int J Behav Nutr
695 Phys Act. 2011 Jan 26;8(1):1–12.

696 60. Larouche R, Mammen G, Rowe DA, Faulkner G. Effectiveness of active school transport
697 interventions: A systematic review and update. BMC Public Health. 2018 Feb 1;18(1):1–18.

698 61. Saunders LE, Green JM, Petticrew MP, Steinbach R, Roberts H. What Are the Health Benefits
699 of Active Travel? A Systematic Review of Trials and Cohort Studies. PLoS One. 2013 Aug
700 15;8(8):e69912.

701 62. Daly-Smith A, Quarmby T, Archbold VSJ, Corrigan N, Wilson D, Resaland GK, et al. Using


- 702 a multi-stakeholder experience-based design process to co-develop the Creating Active
703 Schools Framework. *Int J Behav Nutr Phys Act*. 2020 Feb 7;17(1):1–12.
- 704 63. UNESCO. Research and development expenditure (% of GDP) - Latin America & Caribbean,
705 European Union | Data [Internet]. Data Bank, World Bank. 2018 [cited 2022 Jun 20].
706 Available from: [https://data.worldbank.org/indicator/GB.XPD.RSDV.GD.ZS?locations=ZJ-](https://data.worldbank.org/indicator/GB.XPD.RSDV.GD.ZS?locations=ZJ-EU)
707 EU
- 708 64. ECLAC. Productive development in open economies [Internet]. San Juan, Puerto Rico; 2004
709 [cited 2022 Jun 20]. Available from:
710 [https://repositorio.cepal.org/bitstream/handle/11362/13092/S2004056_en.pdf?sequence=1&i](https://repositorio.cepal.org/bitstream/handle/11362/13092/S2004056_en.pdf?sequence=1&isAllowed=y)
711 [sAllowed=y](https://repositorio.cepal.org/bitstream/handle/11362/13092/S2004056_en.pdf?sequence=1&isAllowed=y)
- 712 65. FAO, IFAD, PAHO, WFP, UNICEF. Regional Overview of Food Security and Nutrition in
713 Latin America and the Caribbean 2020 – Food security and nutrition for lagging territories
714 [Internet]. Santiago; 2021 [cited 2022 Jun 24]. Available from:
715 <https://www.fao.org/3/cb2242en/cb2242en.pdf>
- 716 66. Molina M, Anderson LN, Guindon GE, Tarride JE. A review of implementation and
717 evaluation of Pan American Health Organization’s policies to prevent childhood obesity in
718 Latin America. *Obes Sci Pract*. 2021;
- 719 67. Taillie LS, Busey E, Stoltze FM, Dillman Carpentier FR. Governmental policies to reduce
720 unhealthy food marketing to children. *Nutr Rev*. 2019 Nov 1;77(11):787–816.
- 721 68. Massri C, Sutherland S, Källestål C, Peña S. Impact of the food-labeling and advertising law
722 banning competitive food and beverages in Chilean public schools, 2014–2016. *Am J Public*
723 *Health*. 2019 Aug 7;109(9):1249–54.
- 724 69. López-Olmedo N, Jiménez-Aguilar A, Morales-Ruan M del C, Hernández-Ávila M, Shamah-

- Levy T, Rivera-Dommarco JA. Consumption of foods and beverages in elementary schools: Results of the implementation of the general guidelines for foods and beverages sales in elementary schools in Mexico, stages II and III. *Eval Program Plann.* 2018 Feb 1;66:1–6.
70. Rodríguez Osiac L, Cofré C, Pizarro T, Mansilla C, Herrera CA, Burrows J, et al. Using evidence-informed policies to tackle overweight and obesity in Chile. *Rev Panam Salud Pública.* 2017;41(e156):1–5.
71. Jensen ML, Carpentier FD, Adair L, Corvalan C, Popkin BM, Taillie LS. Examining Chile’s unique food marketing policy: TV advertising and dietary intake in preschool children, a pre- and post- policy study. *Pediatr Obes.* 2021;16(4).
72. Taillie LS, Bercholz M, Popkin B, Reyes M, Colchero MA, Corvalán C. Changes in food purchases after the Chilean policies on food labelling, marketing, and sales in schools: a before and after study. *Lancet Planet Heal.* 2021 Aug 1;5(8):e526–33.
73. Gračner T, Marquez-Padilla F, Hernandez-Cortes D. Changes in Weight-Related Outcomes Among Adolescents Following Consumer Price Increases of Taxed Sugar-Sweetened Beverages. *JAMA Pediatr.* 2021 Dec 13;
74. Corvalán C, Garmendia ML, Jones-Smith J, Lutter CK, Miranda JJ, Pedraza LS, et al. Nutrition status of children in Latin America. *Obes Rev* [Internet]. 2017 Jul 1 [cited 2021 Dec 6];18:7–18. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1111/obr.12571>
75. Vega-Salas MJ, Caro P, Johnson L, Papadaki A. Socioeconomic Inequalities in Dietary Intake in Chile: A Systematic Review. *Public Health Nutr.* 2021 Jul 12;1–16.
76. Vega-Salas MJ, Caro P, Johnson L, Armstrong MEG, Papadaki A. Socioeconomic inequalities in physical activity and sedentary behaviour among the chilean population: A systematic review of observational studies. *Int J Environ Res Public Health.* 2021 Sep 1;18(18):9722.

- 748 77. Mayén AL, Marques-Vidal P, Paccaud F, Bovet P, Stringhini S. Socioeconomic determinants
749 of dietary patterns in low- and middle-income countries: a systematic review. *Am J Clin Nutr.*
750 2014 Dec 1;100(6):1520–31.
- 751 78. Mazariegos M, Auchincloss AH, Braverman-Bronstein A, Kroker-Lobos MF, Ramírez-Zea
752 M, Hessel P, et al. Educational inequalities in obesity: a multilevel analysis of survey data
753 from cities in Latin America. *Public Health Nutr.* 2021;1–9.
- 754 79. Halpern B, Louzada ML da C, Aschner P, Gerchman F, Brajkovich I, Faria-Neto JR, et al.
755 Obesity and COVID-19 in Latin America: A tragedy of two pandemics—Official document
756 of the Latin American Federation of Obesity Societies. *Obes Rev.* 2021 Mar 1;22(3):e13165.
- 757 80. Cortinez-O’Ryan A, Moran MR, Rios AP, Anza-Ramirez C, Slovic AD. Could severe
758 mobility and park use restrictions during the COVID-19 pandemic aggravate health
759 inequalities? Insights and challenges from Latin America. *Cad Saude Publica.* 2020 Oct
760 5;36(9):e00185820.
- 761 81. Ruíz-Roso MB, de Carvalho Padilha P, Matilla-Escalante DC, Brun P, Ulloa N, Acevedo-
762 Correa D, et al. Changes of Physical Activity and Ultra-Processed Food Consumption in
763 Adolescents from Different Countries during Covid-19 Pandemic: An Observational Study.
764 *Nutrients.* 2020 Jul 30;12(8):2289.
- 765 82. Garcia PJ, Alarcón A, Bayer A, Buss P, Guerra G, Ribeiro H, et al. COVID-19 Response in
766 Latin America. *Am J Trop Med Hyg [Internet].* 2020 Nov 1 [cited 2021 Dec 21];103(5):1765.
767 Available from: /pmc/articles/PMC7646820/
- 768 83. Knaul F, Arreola-Ornelas H, Porteny T, Touchton M, Sánchez-Talanquer M, Méndez Ó, et al.
769 Not far enough: Public health policies to combat COVID-19 in Mexico’s states. *PLoS One.*
770 2021 Jun 1;16(6):e0251722.

- 771 84. Norström A V., Cvitanovic C, Löff MF, West S, Wyborn C, Balvanera P, et al. Principles for
772 knowledge co-production in sustainability research. Nat Sustain 2020 33. 2020 Jan
773 20;3(3):182–90.
- 774 85. Irwin EG, Culligan PJ, Fischer-Kowalski M, Law KL, Murtugudde R, Pfirman S. Bridging
775 barriers to advance global sustainability. Nat Sustain 2018 17. 2018 Jul 16;1(7):324–6.
- 776
- 777

779 Figure 1. PRISMA 2020 flow diagram



780

781 *Main reasons for excluding records at title/abstract screening were: studies not including human participants, studies not conducted in LAC region, non-
782 prospective studies, studies not conducted within or around school settings, not-peer reviewed, among others.
783

Author	Country /State or City (Area)	Study design	School n (control/ treatment)	Students n (control/ treatment) [baseline]	Students n (control/ treatment) [follow-up]	Mean Age (SD) (control/ treatment) [grade/ education level]	Intervention length	Environment intervention type	Environmental intervention component	Other Intervention components
Alvirde-García (2013)(46)	Mexico/ State of Mexico (semi-rural)	C-RCT	2/3	755/1927	408/816	9.1(1.7)/ 9.0(1.7) [4 th and 5 th grade/ primary]	3 years (28 months)	Food provision	Increasing availability of fruits and vegetables and products low in saturated fat and in sugar in school canteens	(1) Nutritional education; (2) PA education. School curriculum (booklets and activity guide) and workshops for parents and school vendors
Andrade (2014)(43)	Ecuador/ Cuenca (urban)	C-RCT	10/10	740/700	533/550	12.9(0.8)/ 12.8(0.8) [8 th and 9 th grade /lower secondary]	3 years (28 months)	Built environment	Drawing of a walking trail on the school's playground	(1) PA education; (2) SB education. School curriculum (booklets), workshops for parents, social events, posters
Barbosa Filho (2017)(34,35)	Brazil/ Fortaleza (urban)	C-RCT	3/3	594/588	537/548	12-15 [7 th -9 th grade /lower secondary]	4 months	Built environment	PA equipment (balls, rackets, etc.)	(1) Health education; (2) PA education. School curriculum (booklets and interactive media), workshops for teachers, posters and pamphlets
González (2014)(47)	Chile/ Santiago metropolitan region (rural)	LQE	1/5	192/784	192/784	10 (2.9)/ 9.2(3.1) [preschool – 8 th grade /primary + lower secondary]	9 months	Food provision	Handout fruits 3 times-per-week to students and a fruit basket to the family at the end of the year	(1) Nutritional education. School curriculum (activities) and workshops for parents and teachers
Gutiérrez-Martínez (2018)(48)	Colombia/ Bogotá (urban)	C-RCT	1/1 (TG1)/1 (TG2)	60/60 (TG1)/68 (TG2)	45/34 (TG1)/44 (TG2)	10.6(0.8)/ 10.4(0.6)/ 10.4(0.7)	10 weeks	Built environment	PA equipment (ribbons, balls, hoops, stairs, parachute and mats)	(1) PA education Structured PA education during recess

						[5 th grade / primary]			to support PA during recess	(2) Daily SMS PA reminder
Rausch Herscovici (2013)(42)	Argentina/ Rosario (urban)	C-RCT	2/4	171/234	164/205	9.8(0.7)/ 9.6(0.8) [5 th and 6 th grade / primary]	6 months	Food provision	Provision of healthy food items in snack bar options	(1) Nutritional education; (2) PA education. Workshops for students and parents
Ramírez-López (2005)(41)	Mexico/ Sonora (urban and rural)	LQE	N/R	610	106/254	8.4(1.3)/ 8.6(1.3) [1 st to 5 th grade / primary]	9 months	Food provision	Provision of free school breakfast	
Safdie (2013)(45)	Mexico/ Mexico City (urban)	C-RCT	11/8 (TG1)/8 (TG2)	354/252 (TG1)/254 (TG2)	354/252 (TG1)/254 (TG2)	9.8(0.8)/ 9.7(0.7)/ 9.7(0.7) [4 th and 5 th grade / primary]	18 months	Food provision/ Built environment	Limiting the availability of SSB and energy-dense foods at school canteens Improve school premises and provide sports equipment	(1) Nutritional education (2) PA education School curriculum (activities and booklets), social marketing and workshops for teachers, school vendors and authorities. Structured PA activities during PE, recess, and free time.
Shamah Levy (2012)(49)	Mexico/ State of Mexico (urban and rural)	C-RCT	30/30	510/509	499/498	10 [5 th grade / primary]	6 months	Built environment	Provide sports equipment	(1) Nutritional education (2) PA education Workshops and materials for students, parents, school vendors and school staff. Social marketing (puppet show, audio spots, banners). Structured PA before the start of classes and during recess

BMI: Body Mass Index; C-RCT: Cluster RCT; CG: Control group; LQE: Longitudinal quasi-experimental design; n: number; PA: Physical activity; RCT: Randomised Controlled Trial; SB: Sedentary behaviour; SD: Standard deviation; SSB: Sugar-sweetened beverages; TG: Treatment group; % percentage

787 Figure 2. Risk of bias of cluster RCT (RoB2)

		Risk of bias domains						
		D1	D1b	D2	D3	D4	D5	Overall
Study	Alvirde-García (2013)	+	+	-	+	+	-	-
	Andrade (2014)	+	+	+	+	+	+	+
	Barbosa Filho (2017)	+	+	+	+	+	+	+
	Gutiérrez-Martínez (2018)	+	+	+	+	+	+	+
	Rausch Herscovici (2013)	+	+	+	+	+	+	+
	Safdie (2013)	+	+	+	+	+	+	+
	Shamah Levy (2013)	+	+	+	+	+	+	+
		Domains: D1 : Bias arising from the randomization process. D1b: Bias arising from the timing of identification and recruitment of Individual participants in relation to timing of randomization. D2 : Bias due to deviations from intended intervention. D3 : Bias due to missing outcome data. D4 : Bias in measurement of the outcome. D5 : Bias in selection of the reported result.						Judgement ⚠ High risk - Some concerns ⬤ Low

788

789 Figure 3. Risk of bias of non-randomised controlled trials (ROBINS-I)

		Risk of bias domains								
		D1	D2	D3	D4	D5	D6	D7	Overall	
Study	González (2014)	⊗	⊕	⊕	⊖	⊕	⊕	⊕	⊖	
	Ramírez-López (2005)	⊗	⊗	⊕	⊖	⊕	⊕	⊕	⊗	
		Domains: D1: Bias due to confounding. D2: Bias due to selection of participants. D3: Bias in classification of interventions. D4: Bias due to deviations from intended interventions. D5: Bias due to missing data. D6: Bias in measurement of outcomes. D7: Bias in selection of the reported result.							Judgement ⚡ Critical ⊗ Serious ⊖ Moderate ⊕ Low	

790

791 Table 2. Outcome effect summary of the included studies+

First author (year)	Outcome	Mean difference /OR*	Lower CI	Upper CI	Statistical test
Alvirde-García (2013)(46)	BMI-for-age percentile [CDC]	-0.07	-0.12	-0.02	ANOVA
Andrade (2014)(43)	BMI Z-Score	0.02	-0.02	0.06	Difference-in-difference
Barbosa Filho (2016)(34,35)	BMI-for-age Z-score [WHO 2007]	0.09	0.02	0.16	Generalized linear models
González (2014)(47)	Overweight (%) [WHO 2007]	0.89	0.48	1.64	T-test and two-sample Wilcoxon rank-sum test
	Obesity (%) [WHO 2007]	1.15	0.60	2.21	
Gutiérrez-Martínez (2018)(48)	BMI Z-Score (TG1) [WHO 2007]**	0.50	-4.56	5.56	Difference-in-difference
	BMI Z-Score (TG2) [WHO 2007]**	0.20	-6.58	6.98	
Rausch Herscovici (2013)(42)	BMI (kg/m ²) (F) [CDC]↓	-0.20	-1.18	0.78	ANOVA
	BMI (kg/m ²) (M) [CDC]↓	-0.34	-1.40	0.72	
	BMI Z-Score (F) [CDC]↓	-0.60	-9.95	8.75	
	BMI Z-Score (M) [CDC]↓	-1.40	-3.49	0.69	
Ramírez-López (2005)(41)	BMI (kg/m ²) [CDC]	0.30	-0.06	0.66	ANCOVA
	BMI Z-score [CDC]	0.08	-0.02	0.18	
	Body fat %	-0.30	-0.66	0.06	
	Fat-free body mass (kg)	0.10	0.03	0.17	
Safdie (2013)(45)	BMI (TG1) [IOTF]**	1.30	-0.25	2.85	Generalized linear models
	BMI (TG2) [IOTF]**	-0.10	-0.22	0.02	
Shamah Levy (2012)(49)	Overweight (%) [IOTF]	0.45	0.73	1.11	Generalized ordinal logistic regression
	Obesity (%) [IOTF]	0.34	0.51	0.91	

792 *Mean differences were estimated for continuous variables and Odds ratios (OR) for dichotomous outcomes + Values in
793 bold are significant results for the corresponding statistical tests (p<0.05) **Study presented 2 treatment groups ↓ Study
794 reports results for the subsample of girls and boys, respectively. BMI: Body Mass Index; CDC: Center for Disease Control
795 (50); F: Female; IOM: International Obesity Task Force (51); M: Male; MD: Mean difference; OR: Odds Ratio; SD:
796 Standard deviation; TG: Treatment group; WHO: World Health Organization (52).

