University of Reading

Subsistence behavior during the Initial Upper Paleolithic in Europe: site use, dietary practice, and carnivore exploitation at Bacho Kiro Cave (Bulgaria)

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Smith, G. M. ORCID: https://orcid.org/0000-0001-7155-5140, Spasov, R., Martisius, N. L., Sinet-Mathiot, V., Aldeias, V., Rezek, Z., Ruebens, K. ORCID: https://orcid.org/0000-0002-5621-5786, Pederzani, S., McPherron, S. P., Sirakova, S., Sirakov, N., Tsanova, T. and Hublin, J.-J. (2021) Subsistence behavior during the Initial Upper Paleolithic in Europe: site use, dietary practice, and carnivore exploitation at Bacho Kiro Cave (Bulgaria). Journal of Human Evolution, 161. 103074. ISSN 1095-8606 doi: 10.1016/j.jhevol.2021.103074 Available at https://centaur.reading.ac.uk/117104/

It is advisable to refer to the publisher's version if you intend to cite from the work. See Guidance on citing.

To link to this article DOI: http://dx.doi.org/10.1016/j.jhevol.2021.103074

Publisher: Elsevier

University of Reading

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Subsistence behaviour during the Initial Upper Palaeolithic in Europe: Site use, dietary practice and carnivore exploitation at Bacho Kiro Cave (Bulgaria).

Geoff M. Smith[a*], Rosen Spasov[b], Naomi L. Martisius[c,a], Virginie Sinet-Mathiot[a], Vera Aldeias[d], Zeljko Rezek[a,e], Karen Ruebens[a], Sarah Pederzani[a,f], Shannon P. McPherron[a], Svoboda Sirakova[g], Nikolay Sirakov[g], Tsenka Tsanova[a], Jean-Jacques Hublin[a,h]

[a] Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103, Leipzig, Germany.

[b] Archaeology Department, New Bulgarian University, 21 Montevideo Str., 1618 Sofia.

[c] Department of Anthropology, University of Tulsa, 800 South Tucker Drive, 74104, Tulsa, USA.

[d] Interdisciplinary Centre for Archaeology and the Evolution of Human Behaviour, Universidade do Algarve, FCHS, Universidade do Algarve, Campus de Gambelas 8005-139 Faro, Portugal.

[e] University of Pennsylvania Museum of Archaeology and Anthropology, University of Pennsylvania, 3260 South Street, Philadelphia, PA 19104, USA.

[f] Department of Archaeology, University of Aberdeen, Aberdeen, AB24 3FX, UK.

[g] National Institute of Archaeology with Museum, Bulgarian Academy of Sciences, 2 Saborna Str., 1000 Sofia, Bulgaria.

[h] Collège de France, 11, place Marcelin Berthelot, 75231, Paris Cedex 05, France.

[*] Corresponding author; email address: geoffrey_smith@eva.mpg.de

Abstract

The behavioural dynamics underlying the expansion of *Homo sapiens* into Europe remains a crucial topic in human evolution. Due to poor bone preservation, past studies have strongly focused on the Initial Upper Palaeolithic (IUP) stone tool record. Recent excavations and extensive radiocarbon dating at Bacho Kiro Cave (Bulgaria) pushed back the arrival of IUP *Homo sapiens* into Europe to ca. 45,000 years ago. This site has exceptional bone preservation and we present the study of 7431 faunal remains from across two IUP layers (I and J) and one Middle Palaeolithic layer (K). We identified a shift in site use and occupation intensity through time, marked by increased find density and human modifications in Layer I. Alongside a decrease in carnivore presence and seasonality data demonstrating human presence in all seasons, this indicates a more frequent or prolonged occupation of the site by IUP groups. Contrary, the dietary focus across the IUP and MP layers is similar, centred on the exploitation of species from a range of habitats including wild cattle, bison, deer, horse and caprines. While body parts of large herbivores were selectively transported into the site, the bears remains suggest that these animals died in the cave itself. A distinct aspect of the IUP occupation is an increase in carnivore remains with human modifications, including these cave bears but also smaller taxa (e.g.,wolf, fox). This can be correlated with their exploitation for pendants, and potentially for skins and furs. At a broader scale we identified similarities in subsistence behaviour across IUP sites in Europe and western Asia. It appears that the first IUP occupations were less intense with find densities and human modifications increasing in

succeeding IUP layers. Moreover, the exploitation of small game appears to be limited across IUP sites while carnivore exploitation seems a recurrent strategy.

Keywords: Bacho Kiro Cave; Initial Upper Palaeolithic; Zooarchaeology; Middle-Upper Palaeolithic transition; *Homo sapiens*; subsistence behaviour

Introduction

The Initial Upper Palaeolithic

A key question in human evolution relates to which mechanisms drove the global dispersal of groups of early *Homo sapiens* out of Africa and the subsequent replacement of local populations of other hominins (Hublin, 2013, 2015; Higham et al., 2014; Fu et al., 2015; Ruebens et al., 2015; Boaretto et al., 2021; Hajdinjak et al., 2021; Lalueza-Fox, 2021). In Europe and western Asia, this dispersal intensified after 50ka. with the emergence of the Initial Upper Palaeolithic (IUP) record . The IUP is defined based on a specific type of blade production that retains elements of Levallois technology but with a volumetric core exploitation and Upper Palaeolithic tool types (Kuhn et al., 2009; Kuhn and Zwyns, 2014; Zwyns et al., 2019). A small number of faunal assemblages associated with IUP contexts have been recovered, mainly from caves, with open-air contexts exhibiting poor or no bone preservation [e.g., Bohunician in central Europe; Svoboda and Bar-Yosef (2003); Skrdla (2014); Ruebens et al. (2015)]. At IUP sites with good organic preservation, ornaments and bone tools are also abundant (e.g., Üçağızlı cave in Turkey, Bacho Kiro Cave in Bulgaria, Ksar 'Akil in Lebanon; [Marks (1990); Kuhn et al. (1999); Hublin et al. (2020); Kuhn et al. (2001); Bosch et al. (2015); ;Zwyns et al. (2014); Bosch (2018)]. Overall, our understanding of

IUP subsistence practices, carcass exploitation and resource use is still limited, especially for Europe, and requires further investigation and comparison with local Neanderthal subsistence (Kuhn et al., 2009; Bosch et al., 2015; Bosch, 2018).

New excavations at Bacho Kiro Cave (Bulgaria) have provided early and substantial evidence of the dispersal of Upper Palaeolithic *Homo sapiens* into Europe dated around 45 ka BP (Fewlass et al., 2020; Hublin et al., 2020). This paper will present in-depth taphonomic and zooarchaeological analyses of the rich faunal assemblages recovered from the Middle and Initial Upper Palaeolithic layers of this cave, providing new and much needed insights into the subsistence strategies and dietary practices of these early groups of IUP *Homo sapiens*.

The site of Bacho Kiro Cave

Bacho Kiro Cave is a critical site for discussions relating to the transition from Neanderthal to modern human populations in Europe due to its stratigraphic sequence rich in well-preserved archaeological material (Kozłowski and Ginter, 1982; Tsanova, 2008; Hublin, 2015; Ruebens et al., 2015). The cave is at the mouth of a large karstic system located 5km west of the town of Dryanovo, on the northern slope of the Balkan Mountains range (Stara Planina) and about 70km south of the Danube River.

Dorothy Garrod (American School of Prehistoric Research) and Rafail Popov (Bulgarian Academy of Science) conducted one of the first excavations in 1938, though the scope of these excavations was relatively narrow and there is limited information regarding the stratigraphy and context (Garrod et al., 1939; Ginter and Kozlowski, 1982). Subsequent

excavations (1971-1975) led by Boleslaw Ginter and Janusz K. Kozlowski provided the most comprehensive record of the site, stratigraphy and archaeology (Kozłowski and Ginter, 1982). These excavations produced large assemblages of lithics and fauna (Kozłowski and Ginter, 1982), alongside fragmentary human remains that were subsequently lost. The cave is notable for the distinctive lithic assemblages from Layers 11 and 11a consisting of elongated Levallois-like blades, retouched points, end scrapers and splintered pieces (Kozłowski and Ginter, 1982; Tsanova, 2008, 2012) as well as pendants made of animal teeth (Guadelli, 2011). The lithic assemblage from Layers 11 and 11a was originally labelled 'Bachokirian,' and interpreted as being transitional between the Middle Palaeolithic (MP) and Upper Palaeolithic (UP; (Tsanova and Bordes, 2003; Tsanova, 2008). The assemblage is now recognized as part of the lUP, representing the earliest occurrence of Late Pleistocene *Homo sapiens* in Europe (Kuhn and Zwyns, 2014; Fewlass et al., 2020; Hublin et al., 2020). Kozłowski and Ginter (1982) described the faunal remains from these excavations in a series of detailed chapters within the final site report. While these provided detailed information about the palaeoenvironmental context and change throughout sequence, taphonomic data is limited and it has hindered our understanding of site formation and the role and importance of carnivores and humans in site formation.

a
Main Sector
Niche 1
1970s
excavation
Layer I
Stairs


b
Main sector
Niche 2
Niche 1
visitor`s walkway
Removed beton wall
Stairs
entrance
N
0
5 m
Excavated 2015-2018
Excavated 2015-2018;
faunal analysis ongoing
Not excavated
Excavated 1971-1975
Stratigraphic section
see figure 1c and 1d


c
CC8
BB8
G2(9b)
H(9c/10)
I(11)
J(11a)
K(12)
350 cm
400 cm
50 cm
Bedrock


d
G2 (9b)
H
I (11)
J (11a)
K (12)

102

Figure 1: a) Interior of Bacho Kiro Cave highlighting the location and proximity of the Main Sector and Niche 1 (modified from Fewlass et al., 2020: Supplementary Fig. 1); note the dark color of Layer I within Niche 1. The 1970s excavation is located beneath the concrete floor; b) Site plan and excavation grid at Bacho Kiro Cave (modified from Hublin et al., 2020: Extended Data Fig. 1). Fauna from the squares colored dark red have been analyzed fully while the light red squares indicate that excavation and faunal analysis is ongoing; c) Stratigraphic section from CC8-BB8 (orange line on site plan) illustrating position of Layers I, J and K in Niche 1 (numbers represent old layer assignments by Kozlowski and Ginter, 1982; letters represent layer assignments from new excavations as detailed in Hublin et al., 2020); d) Frontal view of stratigraphic section from Niche 1 sector showing layer subdivisions; note the distinctive dark color and composition of Layer I (modified from Hublin et al., 2020: Extended Data Fig. 1).

New excavations were initiated in 2015 led by the National Archaeological Institute with Museum-Bulgarian Academy of Science in Sofia and the Max Planck Institute for Evolutionary Anthropology. Fieldwork focused on two sectors that contained well-preserved sequences: the Main Sector (ca. 3m^2 excavated) and the previously unexcavated Niche 1 (ca. 8m^2 excavated; Figure 1). The recovered archaeological material illustrates that Layer I from the Main Sector (and its equivalent in Niche 1, N1-I) is consistent with Layer 11 and Layer J (and its equivalent in Niche 1, N1-J) is consistent with Layer 11a from Ginter and Kozlowski's excavations (Kozłowski and Ginter, 1982). This is supported by an extensive set of new radiocarbon samples that were dated with exceptional precision (Fewlass et al., 2020). The IUP at Bacho Kiro Cave starts within Layer J (ca. 46,000 years ago), and intensifies within Layer I which is dated between 45,040 and 43,280 cal BP (95.4% probability; ^{14}C dates recalibrated using IntCal20; Reimer et al. (2020)). The presence of several directly dated *Homo sapiens* bone fragments in Layer N1-I confirms the modern human association with the IUP record (Hublin et al., 2020). In Niche 1 these IUP layers are underlain by one layer of Middle Palaeolithic (Layer K). The vast majority of archaeological material from these new excavations come from the Niche 1 area where the find density is

also higher (see Figure 1-2). Densities are much lower in Layer K (Table 1), but the lithic assemblage, consisting mostly of Levallois flakes, is distinct from the overlying Initial Upper Palaeolithic in terms of technology, typology and raw material usage (Hublin et al., 2020). Lithic material from Layer K is produced, mainly, on coarser grained igneous rocks, while the IUP from Layers I and J is manufactured on fine grained imported flint and is often fragmented, reduced and reused.

Table 1: Fauna from Layers I, J and K in Niche 1 and Main Sector.

Layer	Archaeological Unit	Lithic	Bone	nbuckets	Volume (m^3)	Litres	bone/l	lithic/l
I	IUP	1,701	12,685	100	0.9	900	14.1	1.89
J	IUP	38	1,294	179	1.6	1,611	0.8	0.02
K	MP	81	714	159	1.4	1,431	0.5	0.06

Abbreviations: Lithic = number of lithics >20 mm, Bone = number of bones >20 mm, nbuckets = number of buckets of excavated sediment (a bucket is 9 liters), bone/l = number of bones per liter of excavated sediment, lithic/l = number of lithics per excavated liter of sediment.

Importantly, these new excavations have produced a large, well-contextualized faunal assemblage of individually piece-plotted faunal remains from throughout the stratigraphic sequence (total number of specimens = 18713). Most of these remains (number of specimens = 16402) were recovered from layers associated with the IUP (Layers I and J) and MP (Layer K; see Table 1-2) and several contact zones. During excavation, due to relatively high water content in the clay portions of the sediment it was often difficult to assign bone material to a specific layer at the interface between Layers I and J or J and K, so these bones are assigned into layer contacts I/J or J/K (for a detailed breakdown see Supplementary Online Material [SOM] Table S1). . All sediment was collected by 9 litre buckets and wet-screened on site through 6 and 1.2mm meshes to form two fractions (Hublin et al., 2020).

From both fractions all lithic and faunal remains were collected and will form part of a later study.

Figure 2 visualises the distribution of all piece plotted faunal specimens >20mm (Figure 2a) and the number of bones per litre of excavated sediment (Figure 2b) from Layers I-K within both Niche 1 (squares A to DD) and the Main Sector (squares G-E). This figure illustrates both the large quantity of faunal remains from Niche 1, compared to the Main Sector, and also the high density of remains within Layer I (see Table 1). Indeed, bone per litre of excavated sediment is 17.5 times greater in Layer I compared to Layer J and 28.2 times greater than in Layer K. Micromorphological analyses of Layers I and J show, at times, crude bedding of sands and silts, suggesting temporal hiatuses with exposed surfaces being affected by low energy sheet wash (Hublin et al., 2020). While these gaps have been identified in, at least, the top of Layer J and in Layer I, further micromorphological research is needed for the lower deposits of Layer K. Independently, however, sedimentation rates do not seem to be significantly high and the accumulation of these layers most probably relates to several visits to the site, hence to the normal time-average nature of archaeological deposits. Thus, previous analyses have illustrated the link between the IUP deposits (Layers I and J) in the Main Sector and Niche 1 both in terms of sedimentology and chronology (Fewlass et al., 2020; Hublin et al., 2020), so the fauna from these layers in both sectors are combined and analysed together along with Layer K from Niche 1.



Figure 2: a) Spatial distribution of piece plotted faunal specimens from Niche 1 and Main Sector subdivided by excavation layer (NSP = number of specimens); b) Number of bone specimens/liter of excavated sediment subdivided by layer. Squares are 1 x 1 meter , and the corresponding excavation numbers for each square can be obtained by joining the y-axis letter and the x-axis number.

Materials and methods

For this study, 7431 faunal remains from the 2015-2018 excavations at Bacho Kiro Cave were investigated (curated at History Museum in Dryanovo). This includes all of the fauna from the excavation squares indicated in dark red on Figure 1 alongside a sample of material from row 8 (light red excavation squares on Figure 1, for which excavation and analyses are ongoing), covering both the Niche 1 and Main sector and the IUP and MP layers. This includes all piece plotted material (bone fragments >20 mm [*n* = 6,808; 91.6] and morphologically identifiable remains <20 mm [*n* = 79; 1.1]) and a small portion of the piece plotted fauna that fell below the recording cut-off of 20mm (<20mm; *n* = 544; 7.3).

Table 2: Detailed breakdown of total piece plotted faunal material (NSP) excavated between 2015-2018 by sector and layer.

Sector	Layer	NSP	Analysed	%
	I	133	87	1
Main Sector	I/J	126	108	1
	J	137	98	1
	I	12,558	5,544	75
	I/J	1,392	193	3
Niche 1	J	1,157	678	9
	J/K	185	117	2
	K	714	606	8
Total		16,402	7,431	100

Abbreviations: : NSP = number of specimens.

The first section of this paper presents a new zooarchaeological and taphonomic analysis of faunal material recovered from Layers I, J and K (n = 7013; for data on the contact zones I/J and J/K see SOM Table S1). A specially designed database was constructed using the freely available E4 software (https://www.oldstoneage.com/). Faunal material was recorded using a text based zonal system based on previous methodologies (Smith, 2010, 2013, 2015)), where a zone was only recorded if >50% was present. This allowed for the comprehensive quantification and calculation of the number of identifiable specimens (NISP), minimum number of elements (MNE), minimum number of individuals (MNI) and associated indices such as minimum anatomical units (MAU). NISP was calculated as the number of specimens identified to species and element (Grayson, 1984; Lyman, 1994), though if species determination was unclear fragments were recorded to the family level (e.g. Ursidae sp.). MNE was calculated assessing which zone had the most representation of

195 >50% present and this was further combined with side and fusion data for each specific
196 element (Lyman, 1994; Lyman, 2008; Smith, 2010). The MNI was calculated for each specific
197 element (including left and right) and the final MNI value for each taxon chosen by selecting
198 the highest value. MAU values were normalized by dividing MNE values by the frequency of
199 this element within the skeleton. These values were then normed into %MAU by dividing
200 each MAU by the greatest MAU value in the assemblage [Lewis R. Binford (1981);L. R.
201 Binford (1981); ;Lyman (1994); Lyman (2008); Rendu et al. (2019)].In order to further
202 understand specific site formation processes at Bacho Kiro Cave, zooarchaeological data,
203 such as species and body part identification, were combined with a wide variety of
204 recorded taphonomic attributes.

205 Behrensmeyer's (1978) scheme was used to record the weathering stage of all bones and
206 provides a qualitative scale for understanding the exposure (short/long duration) of faunal
207 material prior to burial. Other biotic and abiotic modification agents recorded included root
208 etching and abrasion (expressed as a percentage of bone surface affected). The schemes
209 range from 0% (no visible modification observed), thorough to 100% (the whole bone
210 surface covered; Smith (2010); Smith (2015)]. Burnt remains were recorded using the
211 specific colour and surface changes as identified by Stiner (1995). Recording the presence,
212 intensity and distribution of these taphonomic agents across the site and throughout the
213 layers at Bacho Kiro Cave provides a detailed framework to contextualize carnivore and
214 human behaviour and site use.

All faunal material was studied under magnification (20x) using an oblique light source, which allowed for a detailed analysis and recording of bone surface conditions and modifications. The identification and distinction of human modifications (skinning, cut marks, deliberate marrow fractures) from other non-anthropogenic processes (trampling, carnivores) has generated considerable debate (Bunn, 1981; Gifford-Gonzalez, 1989a, 1989b, 1991; Fisher, 1995; Domínguez-Rodrigo, 1999; Galán et al., 2009). More recent approaches have attempted to provide quantitative data through the application of high resolution microscopy and the application of machine learning protocols (Cifuentes-Alcobendas and Domínguez-Rodrigo, 2019; Moclán et al., 2019; Pizarro-Monzo and Domínguez-Rodrigo, 2020). Despite much discussion about the criteria and methods for identifying and distinguishing cut marks on fauna (Domínguez-Rodrigo et al., 2017, 2019) the use of low-powered approaches has proven effective for differentiating human from natural bone surface modifications, especially within Middle and Upper Palaeolithic contexts (Blumenschine et al., 1996; Smith, 2015).

Specific carnivore modifications recorded included tooth pits, scratches, crenellation and damage from digestion. Human modifications included those related to butchery and carcass processing such as cut marks, skinning marks, deliberate marrow fractures(identification of impact point and/or percussion notch; see (Lyman, 1994; Fisher, 1995), alongside other secondary uses of organic material for informal bone tools ('retoucher'), formal bone tools (*lissoirs*, awls etc.) and ornaments. To understand fully the

behaviour of *Homo sapiens* at Bacho Kiro Cave, these new faunal analyses are further contextualized with data from sedimentology, micromorphology and lithic data.

To understand whether human groups occupied the site during specific seasons and focused on specific age classes, and if this varied across species, required the recording of different age and seasonality indicators from various species using both cranial (mainly teeth eruption and wear) and post-cranial bone fusion data (Reitz et al., 1999; Weinstock, 2009). Herbivore age was calculated using various methods depending on tooth type. For species with low-crowned teeth such as *Bos/Bison* and cervids the quadratic crown height measure (QCHM) was applied (Steele and Weaver, 2002, 2012, 2012; Steele, 2004, 2004) along with established wear stages (Grant, 1982). For equids, crown height was measured on juveniles and adults and calculated using established equations (Fernandez and Legendre, 2003; Bignon, 2006) and tooth wear stages documented (Levine, 1982). Bear dentition was scored according to the three stage scheme devised by Stiner (1998) and used extensively at numerous Pleistocene sites (Kindler, 2012; Abrams et al., 2014; Romandini et al., 2018). All analyses were undertaken in R v. 4.0.2 (R Core Team, 2020) using RStudio v. 1.4.1103 (RStudio Team, 2021), mainly by using the 'tidyverse' packages v. 1.3.1 (Wickham et al., 2019) and with statistics performed using the 'rstatix' package v. 0.7.0 (Kassambara, 2020). Figures were produced with the 'ggplot2' package v. 3.3.3 (Wickham, 2016) and the 'ggtern' package for ternary graphs (Hamilton and Ferry, 2018) with the exception of the maps that were produced using QGIS v. 3.18.3 (QGIS Development Team, 2009) and the manuscript was written in rmarkdown (v. 2.6) (Xie et al., 2018, 2020; Allaire et

al., 2021). The raw data and RMarkdown scripts to reproduce the article and its analyses along with the SOM are available at https://osf.io/62pbr/.

Results

Species representation and variation

Species representation

The species representation of the new collection is similar to that originally identified by Kozłowski and Ginter (1982), though the proportions vary (see Figure 3). Importantly, the identified macro faunal species are consistent between both the Niche 1 and Main Sector, which support the stratigraphic correspondence of layers between these two excavation areas (also see discussion and analysis in Hublin et al. (2020)).



Figure 3: Percentage of identifiable specimens (%NISP) per taxon by layer; the total percentage of identifiable remains by layer is indicated in the panel title.

Within all layers species identification rates for bone fragments is, relatively, high (I = 19.9% ;J = 31.6%; K = 26.1%) especially for sites of this age range in Europe (Morin, 2008, 2012; Morin et al., 2019; Rendu et al., 2019).

The most frequently identified species include *Bos/Bison* (I = %; J = %; K = %), cave bear (*Ursus spelaeus*; I = 16.6%; J = 45.4%; K = 20.1%) and red deer (*Cervus elaphus*; I = 11.9%; J = 1.7%; K = 3.2%; see Figure 3), though other carnivore (I = 4.6%; J = 8.8%; K = 7.1%) and herbivore (I = 18.2%; J = 9.6%; K = 27.3%) species are represented. Carnivores include cave lion (*Panthera leo spelaea*), leopard (*Panthera pardus*), cave hyaena (*Crocuta crocuta spelaea*), wolf (*Canis lupus*), red fox (*Vulpes vulpes*), brown bear (*Ursus arctos*) along with large to medium sized herbivore taxa (horse [*Equus ferus*], European ass [*Equus hydruntinus*], giant deer [*Megaloceros giganteus*], ibex [*Capra ibex*], chamois [*Rupicapra rupicapra*]). Megafaunal species including mammoth (*Mammuthus primigenius*; K = 1) and rhino (*Stephanorhinus hemiotechus*; I = 0.2%; J = 0.8%), are represented by very few specimens throughout all layers (see SOM Table S2). Overall, the fauna recovered throughout Layers K-I is characteristic of Marine Isotope Stage 3 (MIS 3) throughout south-east Europe, with a mix of cold and more temperate adapted species that have a broad range of climatic/ecological tolerances [Guérin (1982); Hublin et al. (2020); Made (2018); Figure 3).

This variation in species representation is also evident when examining the examining the examining the MNE and MNI values (see SOM Table S3). While these layers include a range of species (see Figure 3) most of these are represented by a low number of elements and individuals (), especially for carnivores (excluding Ursidae; MNI: Layer I = 8; Layer J = 8; Layer K = 6), *Leporidae* sp. (MNI: Layer I = 1; Layer J = 1), birds (*Aquila chrysaetos*; MNI: Layer I =1, *Aves* sp. MNI: Layer I =1) and fish (*Osteichthyes* sp. MNI: Layer I = 1; see SOM Table S3). The exception for carnivores are the Ursidae specimens, which includes *Ursus arctos* and Ursus

sp., but is dominated by cave bear (*Ursus spelaeus*), and represented by a relatively high number of elements and individuals (I: MNE =24; MNI = 11; J: MNE = 17; MNI = 10; K: MNE = 4; MNI = 3)

Within Layer I, herbivore species are most frequent and are, generally, represented by higher numbers of elements and individuals; for example, Bos/Bison, which includes *Bos primigenius*, *Bison priscus* and *Bos/Bison* sp.(MNE = 25; MNI = 12), *Cervus elaphus* which includes both *Cervus elaphus* and *Cervidae* sp. (MNE =21; MNI = 9) and Equidae sp. (MNE =10; MNI = 7), which includes *Equus ferus*, *Equus hydruntinus* and *Equidae* sp.. .

Species variation

To investigate further whether there is a significant change in species representation between layers, we assessed the NISP values for identifiable species from Layers K-I with a chi-square test. In order to observe changes in the representation, the most common species were grouped into broader categories of: *Bos/Bison* (*Bos primigenius*, *Bison priscus* and *Bos/Bison* sp.), Ursidae (*Ursus spelaeus*, *Ursus arctos* and Ursidae sp.) and *Cervus elaphus* (*Cervus elaphus* and Cervidae sp.) . The remaining identifiable species were grouped into broader categories of either herbivore or carnivore, irrespective of body size. The null hypothesis states that the variables of layer and taxon are independent of each other.

Table 3: Number of identified specimens (NISP) by layer and major taxon.

Taxon	NISP I	AR I	NISP J	AR J	NISP K
Bos/Bison	289	**6**	23	**-4.3**	40
carnivore	50	**-3**	21	0.6	11
Cervidae	285	**6**	18	**-2.2**	22

Taxon	NISP I	AR I	NISP J	AR J	NISP K
herbivore	199	**3**	23	**-4.6**	42
Ursidae	268	**-12**	154	**7.6**	39
chi-square		**167**		**66.9**	

Abbreviations: AR = adjusted residuals and composite chi-square values. Significant values in bold.

The change in the proportion of these NISP values was tested through the calculation of composite chi-square values and adjusted residuals (Table 3). These residuals refer to the layer to the left of the residuals column, respectively, and should be read as standard normal deviates (Grayson and Delpech, 2003; VanPool and Leonard, 2011). Absolute values over 1.96 indicate a significant change between layers (these are highlighted in bold in Table 3). Table 3 illustrates changes in the relative abundance of species within Layers K-I. This is driven by a significant decrease in herbivore species between K and J (χ^2 = 66.9, p = <.001). Likewise the abundance of all species within Layer I differs significantly from those within Layer J (χ^2 = 167, p = <.001). Within Layer I this results from a significant increase in all herbivore species relative to carnivore (and especially Ursidae remains) compared to the underlying Layer J. Thus, taxon is dependent on layer: there is an increase in proportion of herbivore and decrease in carnivore taxa within Layer I compared to the underlying layers (see Figure 3).

Bone fragmentation

Bone fragmentation, as expressed by the mean bone length, which was measured across the longest axis of each bone specimen, is similar within Layers I (41.5 ± 20.8mm), J (43.3 ± 23.6mm) and K (43.4 ± 25.3mm). Overall, the distribution of bone fragment lengths appears

similar across layers, with a large proportion of specimens <100mm, and few outliers with larger dimensions. This can be seen in Figure 4 where the mean for each layer is consistently higher than the median (SOM Table S4-S5).



Figure 4: Figure 4. a) Violin plot of the length of piece plotted bone fragments in main layers from Bacho Kiro Cave; b) Boxplot of the length of piece plotted bone fragments for major taxa from main layers (BB = Bos/Bison, Ce = Cervus elaphus, Us = Ursus spelaeus). Black line indicates median and diamonds indicates mean; box represents the interquartile range; box base is 25th percentile while box top represents the 75th percentile with potential outliers plotted. Y-axis on both figures is at the same scale.

A t-test illustrates no significant difference between the mean bone lengths between layers (Layer I versus Layer J: t = -1.7, df =628, p = 0.3;Layer I versus K: t = -1.6, df =557.8, p = 0.3 ; Layer J versus Layer K: t = -0.1, df =1009, p = 1; Figure 4a.).

Similarly, a t-test of mean bone length for the three major species (*Ursus spelaeus*, *Bos/Bison sp.* and *Cervus elaphus*) from Layer I (sample sizes in Layers J and K were too small) illustrates no significant difference in the overall fragment length distributions (*Bos/Bison* versus *Cervus elaphus*: t = 1.1, df =254.6, p = 0.5; *Bos/Bison* versus *Ursus spelaeus*: t = 2.2, df =168.7, p = 0.1; *Cervus elaphus* versus *Ursus spelaeus*: t = 1.2, df =201.6, p = 0.5). Therefore,

346 bone fragmentation does not appear to explain the observed pattern in species
347 representation within Layers K-I (see SOM Tables S2-S3).

348 Taken together, these statistics illustrate variation between the Middle Palaeolithic Layer K
349 at the base of the sequence and the overlying IUP Layers I and J in terms of changes in the
350 relative proportions of carnivore and herbivore taxa. Therefore, it is important to test
351 whether such changes in species representation are related to differences in depositional
352 history within Layers K-I as a result of either natural, carnivore or human action or a
353 combination of all of these.

354 *Species and bone element survival*
355 To understand specific deposit formation processes and the taphonomic history of the
356 Bacho Kiro Cave fauna, especially for Layers I, J and K, requires a detailed assessment of
357 skeletal element representation for the three major species. We merged again the
358 identifiable remains from these species into general categories of Ursidae, Cervidae and
359 *Bos/Bison*. Similarly, identifiable element were merged into more general body portions
360 (cranial: antler, horncore, cranium, mandible, teeth; axial: vertebrae, pelvis; forelimb:
361 humerus, radius, ulna; hindlimb: femur, tibia; foot: carpals, metacarpal, tarsals, metatarsal,
362 phalanges) to provide a broader overview of patterns of representation (modified from
363 Stiner, 1991a, 1991b).



Figure 5: Percentage of identifiable specimens (%NISP) for combined body part portions across three major taxon from Layer I; cranial: antler, horncore, cranium, mandible, teeth; axial: vertebrae, pelvis; forelimb: humerus, radius, ulna; hindlimb: femur, tibia; foot: carpals, metacarpal, tarsals, metatarsal, phalanges. Modified from Stiner (1991a, 1991b). BB = *Bos/Bison*; C = Cervidae; U = Ursidae.

Only Layer I contain sufficient numbers of specimens identifiable to taxon and element portion to discuss patterns of body part representation (Figure 5) . For Layer I, data illustrate differences in the representation of different body parts with both *Bos/Bison* and cervids showing higher proportions of limbs (*Bos/Bison* forelimb = % and hindlimb = %; Cervidae forelimb = % and hindlimb = %), while the ursid remains have a higher percentage of cranial elements (%) compared to limb bones (forelimb = %, hindlimb = %). We ran separate chi-square tests to see whether such differences between taxa and body portion were statistically significant. There is no significant difference between the element

representation of *Bos/Bison* and cervids within Layer I (χ^2 [df = 4, *n* = 528] = 8.5, p = 0.1). In contrast, comparing *Bos/Bison* and cervids with ursids illustrates significant difference between the element representation within Layer I (*Bos/Bison* vs. Ursidae: χ^2 [df = 4, *n* = 455] = 79.8, p = <.001; Cervidae vs. Ursidae: χ^2 [df = 4, *n* = 463] = 62.7, p = <.001). Such variation in element representation could represent different site formation histories, perhaps with the *Bos/Bison* and cervid remains being transported into the cave while the ursid remains indicate that the bears died in the cave itself.

This pattern is confirmed when looking at the minimum anatomical units (MAU) for these taxa from Layer I (see SOM Table S6). The skeletal profiles for *Bos/Bison* show an under-representation of cranial and axial parts (NISP = 29; MAU rib =5.4), with a similar pattern also identified for cervids (NISP = 23; MAU rib =2.1). Skeletal profiles from these species highlight a higher proportion of bone elements from forelimb (*Bos/Bison* NISP = 89; MAU = 22) and hindlimb (NISP = 87; MAU = 18.5). While ursids have high values for some long bone elements such as femur and radius, it is notable that their skeletal profile is dominated by cranial (especially teeth) and also foot elements (cranial NISP = 77; MAU cranial =29; foot NISP = 35; MAU foot =5.3).

Further, plotting relative bone density values (Kreutzer, 1992; Lyman, 1994; Lam et al., 1998, 1999, 2003; Lam and Pearson, 2005) for the most abundant herbivore species within Layers K-I (*Bos/Bison*, cervids and equids) does not illustrate a specific predominance of skeletal remains, though long bone remains appear common (see Figure 6).



Figure 6: Plots of the number of identifiable specimens (NISP) against relative bone density for three major herbivore species from Bacho Kiro Cave Layers I–K (a = Bos/Bison; b = Cervidae, c = Equidae) Note that Layer J is absent because no elements were present to correlate with bone density. Bone density values for Bos/Bison are based on volume density (VD) measures from Kreutzer (1992), and Equidae and Cervidae from BMD1 mean from Lam et al. (1999). Abbreviations on plots relate to specific scan zones on elements: AT = atlas; CA = calcaneus; CE = cervical vertebra; FE = femur; HU = humerus; IS = Ischium; LU = lumbar vertebra; MC = metacarpal; MR = metatarsal; RA = radius; TH = thoracic vertebra; TI = Tibia; UL = ulna.

Figure 6 illustrates element NISP plotted against relative bone density for *Bos/Bison*, cervids and equids. In Layer I there is no significant relationship between relative bone density and preservation. In Layers J and K there are some positive associations with cervids and equids, respectively, suggesting selective preservation of denser skeletal portions, but smaller samples sizes warrant caution in this interpretation. Therefore, in Layer I it appears

411 that there has been neither significant selective destruction of weaker skeletal portions nor
412 an over representation of the densest elements. Taken together with skeletal profile and
413 MAU data this suggests that there was a biased transport of skeletal elements from
414 herbivore species into the cave during the accumulation of Layer I with this transport not
415 including cranial, axial and, to a lesser extent, foot elements of those species.

416 *Bone surface visibility and weathering*

417 Overall, bones from Layers K-I show low proportions of both sub-aerial weathering (stages
418 0-2; Behrensmeyer (1978)] and abrasion (0-50%), which translates into high bone surface
419 visibility (>90%; Figure 7 and SOM Tables S7-S9 for detailed data breakdown).



Figure 7: Bone surface preservation and readability. a) Percentage of specimens displaying low weathering (stage 0–2) or high weathering (3–5) based on Behrensmeyer (1978); b) Percentage of specimens exhibiting low (0–50%) and high surface abrasion (50–100%); c) Percentage of specimens with low (0–50%) and high (50–100%) surface readability.

The extraordinary bone preservation at Bacho Kiro Cave has been recognized in the high collagen yields extracted for radiocarbon dating and ZooMS.

Carnivore and human bone surface modifications

The excellent preservation of the bone surfaces provides ample opportunity to identify and classify bone surface modifications from both carnivore and human action (see Figure 8; SOM Table S10).



Figure 8: Percentage of specimens (%NSP) with carnivore or human modifications within Layers I-K from Bacho Kiro Cave; this excludes burnt specimens (see SOM Tables S10-S15 for detailed breakdown of bone surface modifications including burnt remains).

Overall, Figure 8 illustrates a change in the proportion of bone surface modifications between the MP Layer K (carnivore modification = 11.4%; human modification = 7.4%) and the IUP Layers (J carnivore modification = 7.9%; J human modification = 7.9%; I carnivore modification = 1.3%; I human modification = 12.6%), showing a clear temporal trend of a decrease in carnivore modification and an increase in human modification. We tested the change in this proportion between layers by calculating the composite chi-square values and adjusted residuals (see Table 4). The null hypothesis states that modification types are independent of layers.

Table 4: Number of specimens (NSP) by layer with carnivore or human modifications.

Modification	NSP I	AR I	NSP J	AR J	NSP K
carnivore	76	**-12**	61	-2	69
human	709	**12**	61	2	45
chi-square		**131**		2	

Modification	NSP I	AR I	NSP J	AR J	NSP K

Abbreviations: AR = adjusted residuals and composite chi-square values. Significant values in bold.

There is no significant difference in %NSP of bones with carnivore and those with human modifications between Layers K and J (χ^2 = 2.2, p = 0.1) and thus the null hypothesis cannot be rejected (see Table 4). In contrast there is a strong relationship in the proportion of different modifications between Layers J and I (χ^2 = 130.7, p = <.001; see Table 4). The data for Layers I and J indicate that modification type is dependent upon layer. The main driving factor for this pattern appears to be the large increase in human modifications on bones of Layer I, which is consistent with geoarchaeological and micromorphological data that illustrates increased anthropogenic input in the formation of this deposit (Hublin et al., 2020) and also with the increase in density of lithic material (see Table 1). We also ran chi-square tests to assess whether different modification types (carnivore, human) illustrate significant differences between body size class and across the different layers. All test results were not significant (at 0.05 level). Overall, the large increase in human modifications within Layer I is statistically significant, but there is no relation to how these modifications are distributed across various species body size classes or between layers.

Carnivore bone surface modifications

Carnivore modifications are recorded in very small quantities, compared to human modifications, and distributed across a variety of taxa including common species such as *Bos/Bison* and cave bear.

462 Table 5: Percentage total number of specimens (%NSP) with different carnivore modifications in
463 Layers I-K.

Layer	NSP	%gast	%scratch	%scallop	%tp
I	5,631	0.8	0.1	0.2	0.5
J	776	7.1	0.3	0.6	0.8
K	606	11.1		0.7	0.5

Abbreviations: %gast = percentage of bones with gastric etching; %scratch = percentage of bones with tooth scratch; %scallop = percentage of bones with tooth scalloping; %tp = percentage of bones with tooth pit.

464 Table 5 illustrates that Layers J and K have the highest proportion of bones with recorded
465 carnivore modifications, and the most common modification recorded is digested and acid
466 etched bone (Layer K = 11.1%; Layer J = 7.1%). The higher proportions of carnivore
467 modifications in these layers combined with evidence for coprolites in the
468 micromorphology thin sections (Hublin et al., 2020), and a lower incidence of human
469 modifications, suggests a more substantial role for carnivores in bone accumulation and
470 modification (Figure 9 and SOM Tables S8-S9). The type and frequency of modifications,
471 especially gastric etched bone, suggests activities of cave hyaena though other small
472 carnivore taxa cannot be ruled out. However, the low incidence of carnivore modifications
473 within Layer I, and especially the dramatic reduction in digested bone, suggests a more
474 minimal role for carnivores in the formation of this faunal record. The predominance of
475 human action in the formation of Layer I and upper Layer J is illustrated, further, by the low
476 incidence of carnivore and human modifications on the same specimen (Layer I = 9; Layer J
477 =3; Layer K = 3). Bacho Kiro Cave offered a safe location for carnivore species to transport
478 and consume carcasses during periods when the site was unoccupied by humans.

However, such opportunities appear to have been more infrequent during the IUP compared to the MP.

Figure 9: Examples of carnivore bone surface modifications; a) *Ursus spelaeus* scapula (BB7-1384, I/J) with extensive gnawing and crenellation around glenoid and scapula spine; b) Unknown mammal bone fragment (BB8-1934, I/J) with extensive surface smoothing through carnivore digestion; c) Unknown mammal bone (BB7-1977, Layer K) with extensive surface damage caused by carnivore digestion; d) *Ursus spelaeus* femur (AA7-1214, Layer I) with carnivore gnawing and crenelation

Human bone surface modifications

The change in proportions of human modifications within the MP-IUP layers at Bacho Kiro Cave is statistically significant. Table 6 shows the numbers and proportions of different types of bone surface modifications by humans across the three layers. Layer I contains the

491 highest quantity of modifications for each category and the highest proportion of total
492 modifications (see SOM Table S12). Cut marks make up the largest quantity of
493 modifications in most layers, comprising of between 2-8% of bone fragments with the
494 highest proportion within Layer I. Within Layers J and K the frequency of bones with cut
495 and scrape marks is lower compared to Layer I, while there is a higher proportion of bones
496 that exhibit deliberate fractures related to marrow exploitation. Combined with higher
497 levels of carnivore presence (Figure 8) it seems that human groups were using the cave
498 more intermittently and over shorter periods of time.

499 Table 6: Number of specimens (NSP) and percentage of specimens with different human
500 modifications in Layers I-K.

Layer	NSP	%cm	%scrape	%marrow	%burn
I	5,631	8	2	4	0.7
J	776	3	1	4	0.4
K	606	2	1	5	0.5

Abbreviations: %cm = percentage of bones with cut marks; %scrape = percentage of bones with scraping marks; %marrow = percentage of bones showing marrow extraction (includes impact points and percussion notches); %burn = percentage of bones showing traces of burning.

501 Layer I is again distinct in terms of the quantity and variation of different modifications but
502 also in the increase in proportion of modifications across both carnivore and herbivore
503 taxa, across all body size classes, but especially an increased focus on smaller carnivores
504 and herbivores (see Figure 10 and 11; SOM Table S13). There also appears to be variation
505 in the distribution of different modification types across body size classes for both
506 herbivores and carnivores; for example, higher proportions of marrow fractured bones in
507 large compared to small herbivore taxa with medium sized animals having a value between

508 the two. Additionally, within Layer I the proportion of human bone surface modifications
509 increases across all carnivore body size classes (including wolf, wolverine, cave hyaena,
510 cave bear).

511

Figure 10: Examples of human bone surface modifications; a) *Rupicapra rupicapra* radius (A7-773, Layer I) with disarticulation marks (white arrows) overlain by carnivore tooth pit (black arrow); b) *Bos primigenius* proximal epiphysis of second phalanx (CC7-2120, Layer I) with disarticulation marks; c) *Ursus spelaeus* ulna fragment (BB7-1190, Layer I) with filleting and cut marks overlain by retouching damage; d) *Ursus spelaeus* phalanx (BB7-458, Layer I) with skinning and disarticulation marks; e) Mammal long bone fragment (BB8-283, Layer I) with two separate episodes of use as a retoucher; f) *Ursus spelaues* phalanx (AA7-2043, Layer I) with skinning and disarticulation marks; g) Carnivore rib fragment (CC8-1564, Layer I) with cut and disarticulation marks; h) *Crocuta crocuta spelaea* phalanx (BB7-1062, Layer I) with disarticulation marks; i) *Aquila chrysaetos* talon (BB7-721, Layer I) with cut mark suggesting disarticulation/removal of talon (compare with figures in Romandini et al. (2014) and Radovčić et al. (2015)); j) *Ursus spelaeus* phalanx (BB7-588, Layer I) with cut marks suggesting claw removal.

When comparing the distribution of human modifications across the major species for Layer I the emphasis on the limb bone elements for *Bos/Bison* and cervids is evident with high proportions of all three types of modification (scrape, cut and marrow fractures; Figure 12), supporting the selective transport of limb bone elements to the site and subsequent processing. *Ursidae* remains show a similar level of processing of limb bones but also increased modifications on cranium and foot portions, indicative of skinning and first stages of fur removal (Münzel and Conard, 2004; Kindler, 2012; Romandini et al., 2014). This could represent the direct procurement of such taxa as raw material for bone tools and personal ornaments (see Hublin et al. (2020), Fig. 3) but also potentially for fur (see Figure 11). Further modifications on small-medium sized carnivores (for example *Gulo gulo*, *Canis lupus*, *Crocuta spelaea*; modified from Morin (2012) are more limited but again distributed across cranium, limbs and foot elements, which suggests a more sporadic targeting of such species for fur. Nevertheless, Layer I illustrates clearly an increased and more systematic focus on carnivore species, relative to herbivores, which is distinct from the underlying Layers.



Figure 11: Bone surface modifications across carnivore (carn) and herbivore (herb) grouped by body size class; carnivores: small: wolf (*Canis lupus*), fox (*Vulpes vulpes*), dhole (*Cuon alpinus*); medium: hyaena (*Crocuta crocuta spelaea*); large: cave lion (*Panthera leo spelaea*), leopard (*Panthera pardus*), bear (*Ursus spelaeus*, *Ursus arctos*); herbivores: small: roe deer (*Capreolus capreolus*), goat (*Caprinae* sp.), fallow deer (*Dama dama*), ibex (*Capra ibex*), pig (*Sus scrofa*), ungulate small, ungulate small/med; medium: reindeer (*Rangifer tarandus*), wild ass (*Equus hydruntinus*), horse (*Equus ferus*), red deer (*Cervus elaphus*), *Cervidae* sp., ungulate med/large; large: mammoth (*Mammuthus primigenius*), rhino (*Stephanorhinus hemeiotechus*), giant deer (*Megaloceros giganteus*), *Bos/Bison* (*Bison priscus*, *Bos primigenius*), ungulate large. Size classes are modified from Morin (2012). Burning data are not included.

An interesting feature of the faunal assemblage from Bacho Kiro Cave is the low proportion of bone remains exhibiting evidence for being burnt (total remains from all layers=1.5%; see Table 6; SOM Tables S14-S15). The majority of burnt remains were identified from Layer I (0.7%) and this contrasts within comparatively high figures of burned lithics (Layer I= 13.3%; Layer J= 0%; Layer K= 2.6%). In particular, this low quantity of burnt faunal remains appears at odds with the micromorphological data, which suggests that the composition of Layer I, and especially the dark colour, is a direct result of burnt material, with common charcoal fragments (Hublin et al., 2020). Further analyses of the smaller screened fraction is ongoing and will assess the destruction, fragmentation and overall intensity of burning within the MP-UP sequence, especially from the IUP layers.



Figure 12: Percentage of specimens with human bone surface modifications across major taxa from Layer I at Bacho Kiro Cave separated by body portion; cranial: antler, horncore, cranium, mandible, teeth; axial: vertebrae, pelvis; forelimb: humerus, radius, ulna; hindlimb: femur, tibia; foot: carpals, metacarpal, tarsals, metatarsal, phalanges. Modified from Stiner, 1991a; 1991b. BB = *Bos/Bison*; C = Cervidae; U = Ursidae.

Alongside human butchery and processing modifications there is, potentially, direct evidence for hunting in the form of possible projectile impact marks (PIMs) from both Layers I and K (Fig. Figure 13). These two specimens, both large ungulates, exhibit damage consistent with projectile impact including notch formation with bevelling on the rib margin (BB8-1061) and a bending fracture on another rib (BB7-1870; Smith et al. (2020); O'Driscoll and Thompson (2014); Iovita et al. (2014); Rots and Plisson (2014); Smith (2003); Gaudzinski-

572 Windheuser et al. (2018)]. Further study of these modifications is required, as well as

573 correlation with ongoing lithic usewear studies, to be able to fully interpret these marks.

574

575 Figure 13: Potential projectile impact marks (PIMs) on ungulate rib fragments; a) BB8-1061 (N1-I)
576 exhibits a notch on the rib margin and bevelling (see expanded photo); b) BB7-1870 (N1-K) exhibits a
577 bending fracture and impact point illustrated by white arrow.

578 Finally, the faunal assemblage from Layers I-K illustrates the use of a range of animal

579 carcasses as raw material both for bone tools (*n* = 92; this includes bone retouchers) and

580 personal ornaments (*n* = 29). These were recovered across the site from all the squares

581 excavated in both the Niche 1 and Main Sector. Informal bone tool types, such as

582 retouchers (*n* = 48), are found throughout Layers K-I. Contrary, formal objects (*n* = 19; 0.3%)

583 are only identified within the IUP occupations of Layers I and J and include awls and *lissoirs*

584 (smoothers). Animal teeth pendants and pendant fragments (*n* = 27) were likewise
585 identified only within the IUP Layers and are produced mainly on cave bear teeth, along
586 with wolf and a few herbivore taxa (Hublin et al., 2020). Ongoing work will characterize and
587 contextualize the nature and manufacture of these objects against the broader
588 background of the MP-UP transition in Europe (Martisius et al., *in preparation*; NEWCOMER
589 and WATSON (1984); Newcomer (1974); Bosch et al. (2015); Kuhn et al. (2009); Soressi et al.
590 (2013); Gaudzinski (1999)).

591 *Mortality profiles, site use and seasonality*

592 Bone fusion

593 Overall, the faunal assemblage from Layers K-I produced 65 unfused bone elements
594 representing a small proportion of the total (0.9%) and a limited number of these were
595 assigned securely to both taxa and element (*n* = 12; Table 7).

596 Table 7: Bone elements from different species at Bacho Kiro Cave with first fusion date (FFD) and last
597 fusion date (LFD) in months and season of FFD and LFD (SFFD, SLFD)

Layer	Species	Element	*n*	FFD	LFD	SFFD	SLFD
I	*Bos primigenius*	Metatarsal-Distal	1	24	36		
I	*Bos primigenius*	Phalanx 2-Distal	1	18	18	Autumn	Autumn
I	*Bos primigenius*	Phalanx 2-Proximal	1	18	24	Autumn	Spring
I	*Capra caucasica*	Vertebra-Thoracic-Centrum	1	48	60		
I	Cervidae sp.	Metatarsal-Proximal	1	0	0	Spring	Spring
I	Cervidae sp.	Phalanx 2-Proximal	1	11	17	Spring	Autumn
I	Cervidae sp.	Radius-Proximal	1	5	8	Autumn	Winter
I	*Cervus elaphus*	Axis-Centrum	1	35	42	Spring	Autumn
I	*Cervus elaphus*	Humerus-Distal	1	12	20	Spring	Autumn

Layer	Species	Element	*n*	FFD	LFD	SFFD	SLFD
I	*Equus ferus*	Phalanx 2-Proximal	1	9	12	Winter	Spring
I/J	Cervidae sp.	Humerus-Proximal	1	42	42		
I/J	*Ursus spelaeus*	Phalanx 2-Distal	1	24	36		

Fusion data based on Reitz et al. (1999) and Weinstock (2009).

598 The unfused proximal metatarsal from a cervid, which is normally fully fused at birth,
599 suggests that this was a foetal individual. At least four of the elements identified have a
600 first fusion date (FFD) around 3 years (36 months) suggesting these individuals were, at
601 least dentally, prime adult and suggesting this individual died during the spring season.
602 Furthermore, additional elements (*n* = 4) of individuals less than 12 month old suggest a
603 season of death during spring/summer and winter/autumn, assuming an early May
604 birthing season. Overall, the available post-cranial fusion data suggests a focus on prime-
605 aged animals, at least for herbivore species, and the use of the cave by IUP groups during
606 all seasons.

607 Dental remains

608 Dental remains have been identified for a variety of species though ursid remains are most
609 common throughout all layers. Ursid teeth were grouped to combine teeth from cave bear,
610 brown bear and Ursidae sp. providing a larger sample for analysis. Bears have an unusual
611 dental development and eruption alongside a different season of birth (winter, January)
612 compared to other carnivores (hyaena and canids) and ungulates (generally spring time
613 late May; Discamps and Costamagno (2015); Stiner (1990)]. Much of the information on
614 ageing bear remains from teeth comes from modern brown bear (Stiner, 1990; Stiner,

615 1994, 1998). All bears are born during hibernation and are toothless, developing the full
616 deciduous dentition by the third month with the M1 usually by the fifth month. Bears
617 generally have all permanent dentition erupted by the end of the first year with the
618 eruption of the permanent canines starting during the second year and completion by the
619 end of the third year of life. Knowing the specific timing and eruption of deciduous and
620 permanent dentition allows for the development of a tooth eruption wear scheme that
621 includes nine stages, grouped into three age categories (Juvenile [I-III], Prime [IV-VII], Old
622 [VIII-IX]; Kindler (2012); Stiner (1994); Stiner (1998)]. While the scheme does not provide an
623 estimate for the age at death it provides the ability for inter and intra-site comparisons at
624 an ordinal scale (Figure 14; Stiner (1998): p.311).



Figure 14: Ternary plot with percentage of Ursidae teeth in the juvenile, prime and old categories within Layers I-K from Bacho Kiro Cave. Wear stages defined by Stiner (1994, 1998) and Kindler [(2012);]. Tooth position si represented by different symbols and layers represented by different colors. J = juvenile dominant mortality, NNVA = normal non-violent attrition, LS = living structure, P = prime dominant mortality, O = old dominant mortality.

Ursid deciduous dentition recovered from Layers I and J are all unworn or in very early wear, suggesting these individuals died during winter or early spring time possibly towards the end of hibernation period (March/April). Further, the presence of at least two individuals with permanent M1 not yet erupted suggests individuals under five months of age that died, similarly, during late winter-early spring. In total 74 teeth could be assigned to the wear stages developed by Stiner (1994). Despite low sample sizes for some teeth

637 Ursid mortality profiles for Layers K-I contain a mix of all three age classes (juvenile, prime
638 and old individuals) with perhaps slightly elevated proportions of juvenile and old animals
639 (Figure 14). Overall, these patterns are close to the natural non-violent attrition (NNVA)
640 profile, which Stiner (1998) attributes to natural hibernation death, which also correlates
641 well with the tooth eruption data and skeletal part representation patterns observed,
642 especially within Layers I and J.

643 The seasonal nature of bear occupation at caves, including at Bacho Kiro Cave, suggests
644 that there were extended periods of time where both humans and bears could safely
645 occupy the site. Indeed, Stiner (1998) notes that any disturbance at a denning location can
646 deter potential hibernating bears. While mortality data suggests bear remains at Bacho
647 Kiro Cave accumulated largely as a result of natural attrition, active exploitation of
648 hibernating bear individuals by humans cannot be completely excluded (Kindler, 2012;
649 Romandini et al., 2018). The high proportion of skinning and butchery modifications on
650 these remains suggests access to such remains soon after the individual died in order to
651 exploit skins and other carcass resources, which has been documented at other sites in
652 Europe (Münzel and Conard, 2004; Kindler, 2012; Abrams et al., 2014; Romandini et al.,
653 2018), and also to exploit abundant dental remains for pendants (Hublin et al., 2020).

654 Further seasonality data was obtained from additional carnivore taxa including red fox
655 (*Vulpes vulpes*) and cave hyaena (*Crocuta crocuta spelea*). As Layer I (and the contact I/J) is
656 accumulated largely as a result of human activity these remains likely represent the
657 deliberate accumulation by human groups. In Layer I, at least three adult cave hyaenas are

represented alongside a single deciduous third premolar (Stage I of hyaena wear scheme; (Stiner, 1994): p325). Previous work has documented that modern spotted hyaenas reproduce aseasonally, though in modern groups this tends to peak during the summer (June-August; Stiner (1994); Kruuk (1972)]. Undoubtedly Pleistocene hyaena populations in more northern latitudes, compared to their modern day distribution, had a more restricted breeding season (Stiner, 1994). Therefore, an approximate season of death is spring-summer time, based on modern data. Within Layer J there are no juvenile cave hyaena (all the teeth are permanent) though the crowns are unworn or in the early stages of wear perhaps suggesting the teeth have only just erupted (ca.9-12 months of age) again suggesting a spring-summer season of death. Red fox dentition from Layers I and J is limited (I = 3; J = 1) but contains erupted yet unworn premolars (P2) and molars (M1) in the early stages of wear (Stiner, 1994). Red foxes normally breed between March-April with the permanent premolars and molars erupting around 4 months (Stiner, 1994). The deciduous dentition is generally replaced by permanent teeth when the individual is ca. 6 months old, supporting a human occupation during the spring-summer at Bacho Kiro Cave.

Herbivore age and seasonality data is more limited due to an overall lack of cranial and dental fragments, presumably resulting from differential transport decisions by human groups. For *Bos/Bison*, crown height could be measured on seven teeth from Layers K-I and ages range between 3-10 years indicating prime age animals (juvenile: 0-3 years; prime: 3-12 years; old: 12-20 years; based on Discamps and Costamagno (2015)). There were no cervid remains suitable to measure crown height so age and seasonality cannot be

679 adequately assessed. Most of the teeth recorded in terms of wear and eruption indicate
680 prime aged individuals though in Layer J there are at least three juvenile individuals as
681 expressed by unworn or early wear dP2 and dP3. This indicate these individuals were
682 potentially no older than 6 months old suggesting they died between spring and autumn.

683 Finally, deciduous and permanent equid teeth were assessed in terms of eruption, tooth
684 wear and crown heights were measured, where possible. Individuals assigned to age
685 classes based on the revised schemes of Discamps and Costamagno [(2015); juvenile: 0-2
686 years; prime: 2-15 years; old: 15-25 years). Within Layer I three permanent teeth were
687 measured (ages 14-21 years) and assigned to prime and old age classes with at least one
688 juvenile deciduous premolar (dP4) in early wear suggesting a very young individual up to 1
689 month old (Levine, 1982) that died during spring. Layer J produced a similarly low number
690 of specimens (*n* = 3) with preserved crown heights again producing prime-old age
691 individuals (ages 7-25 years). A single deciduous incisor suggests at least one juvenile
692 individual from Layer J and the heavy wear suggests the animal died between spring and
693 autumn (Levine, 1982). Finally, Layer K preserved seven specimens with enough tooth
694 crown to measure and allowed all teeth to be assigned to prime-old age individuals (7-17
695 years) (see SOM).

696 Overall ageing and seasonality data illustrates that the IUP record at Bacho Kiro Cave
697 accumulated year round based on carnivore and herbivore bone fusion, dental eruption
698 and wear and tooth crown height. Most information comes from the extensive bear
699 remains that suggest a season of death during or soon after hibernation and most likely

represent natural attrition. Data from herbivore taxa is more limited. Juvenile remains from cervids and equids, along with some *Capra sp.* remains, suggest death during the spring-autumn seasons and quite soon after birth. The herbivore taxa illustrate mainly prime-old age individuals with occasional juvenile specimens though the numbers are insufficient to investigate more thoroughly the mortality profiles and hunting practices.

Overall ageing and seasonality data illustrates that there was no preferential season for accumulation during the IUP faunal record at Bacho Kiro Cave. Most information comes from the extensive bear remains that suggest a season of death during or soon after hibernation and most likely represent natural attrition. Data from herbivore taxa are more limited with mainly prime-old age individuals with occasional juvenile specimens though the numbers are insufficient to investigate more thoroughly the mortality profiles and hunting practices. Juvenile remains from cervids and equids, along with some Capra sp. remains, suggest death during the spring-autumn seasons and quite soon after birth. Seasonality data indicate that faunal material accumulated at Bacho Kiro Cave as a result of repeated use of the site by both human and carnivore groups throughout Layers K-I. The intensity of site use by humans within Layer I was substantially greater than earlier layers, either because more people were present, stayed longer, or both.

Discussion

Site use, diet and carnivore exploitation

This study represents the first in depth taphonomic and zooarchaeological analysis of fauna from Bacho Kiro Cave, characterizing the subsistence behaviour of humans during

721 the Initial Upper Palaeolithic. The faunal composition within Layers K-I illustrates climatic
722 shifts at a local and regional scale to a greater extent than previously suggested (Kozłowski
723 and Ginter, 1982; Spassov et al., 2007) and a diversity of habitats surrounding the site. The
724 identification of woolly mammoth (*Mammuthus primigenius*), reindeer (*Rangifer tarandus*)
725 and wolverine (*Gulo gulo*) from Layers I and J indicate a substantially colder climate than in
726 present day, which is also consistent with the, currently available, microfaunal data (Hublin
727 et al., 2020). Importantly, these species could have been more environmentally tolerant
728 compared to modern day populations. Species representation correlates well with high
729 resolution oxygen stable isotope data (Pederzani et al., 2021) suggesting mean annual air
730 temperatures 10-15°C below modern day (Layer J: mean annual temperature -1.1 ± 3.3°C;
731 Layer I: mean annual temperature -3.4 ± 2.5°C). Hence, palaeoclimate reconstructions for
732 Layer J suggest a continental climate similar to present day Russia or central Asia, while the
733 later IUP occupation in Layer I occurred during a climatic phase with a smaller difference in
734 summer-winter temperatures that is closer to modern-day northern Scandinavia. Despite
735 the considerably colder climate and environment around Bacho Kiro Cave, seasonality
736 indicators illustrate human occupation within Layer I throughout all seasons of the year.
737 Therefore, early groups of IUP *Homo sapiens* appear to have occupied south-eastern
738 Europe during colder climates than previously considered and were hunting a range of
739 herbivore taxa from a variety of local environments (see supplementary information in
740 Hublin et al., (2020)). Overall, such data have broader implications for our understanding of
741 the environmental and climatic conditions during the spread of Late Pleistocene *Homo*
742 *sapiens* (Müller et al., 2011; Hublin, 2013; Staubwasser et al., 2018).

743 Micromorphological and sedimentological analysis indicate increased anthropogenic input
744 towards the top of Layer J with human activities intensified within Layer I. This resulted in
745 the distinct dark colour of Layer I, coming from organic matter, including charcoal and, to a
746 lesser extent, burnt bone but also knapped chert, bone splinters, plant remains and some
747 coprolites (Hublin et al., 2020). Results from this study characterise the faunal assemblage
748 from Layer I as an accumulation primarily resulting from human action.

749 The richness of Layer I is exemplified by the high volume of excavated bone material from
750 Layer I (14.1/litre of sediment) compared to the underlying layers (Layer J = 0.8/litre of
751 sediment; Layer K = 0.5/litre of sediment). This is further emphasised by the excellent
752 identification rate (19.9%), which is high especially for Pleistocene localities (Morin et al.,
753 2017a, 2017b; Rendu et al., 2019). This is also the case for Layers J and K and the contact
754 between these layers (I/J and J/K). Faunal data from Bacho Kiro Cave illustrates that the diet
755 of the IUP groups within Layer I remained relatively consistent compared to the underlying
756 Middle Palaeolithic of Layer K with a focus on large-medium sized herbivores (especially
757 *Bos/Bison*, cervids and equids). However, within Layer I there appears to be a more
758 deliberate focus on carnivore species, especially cave bear, representing a specific
759 subsistence behaviour that is distinct from the underlying MP at Bacho Kiro Cave. Detailed
760 analysis reveals an increase in carnivore species alongside a diversification in other
761 herbivore species during the IUP occupation in Layer I and appearing to begin within Layer
762 J. The variation in identification and body portion for major species in Layer I is related to
763 differences in transport decisions (e.g., transport of herbivore limb elements), site

764 formation histories (cave bear, natural accumulation), butchery and processing behaviour
765 and the use and modification of bone for tools and personal ornaments by IUP groups at
766 Bacho Kiro Cave.

767 Bone surface modifications from Layers K-I shows an increase in the proportion of human
768 modifications in Layer I and a corresponding decline in carnivore modifications compared
769 with the underlying horizons. Throughout Layer K the intensity of anthropogenic input is
770 significantly lower in terms of frequency and intensity. Indeed, various proxies (carnivore
771 modified bones, digested elements) suggest that carnivore taxa had a more important role
772 in bone accumulation within this layer. This should in no way be seen to diminish the
773 capacity and hunting capabilities of Neanderthals (Stiner, 1994; Smith, 2010, 2015;
774 Discamps et al., 2011; Starkovich, 2017; Marin et al., 2020) but fits with other indications of
775 the low intensity of Neanderthal occupation signatures at some cave sites and the
776 frequency of large carnivore modifications within these assemblages (Rendu et al., 2019).
777 This could be related to more mobile lifeways resulting in shorter duration of site
778 occupation compared to the IUP occupation in Layer I. The distribution of human
779 modifications within Layer I illustrates a large increase across a range of herbivore and
780 carnivore species and includes unusual species such as raptors (golden eagle [*Aquila*
781 *chrysaetos*]) and Leporidae (Peresani et al., 2011; Morin and Laroulandie, 2012; Romandini
782 et al., 2014). Taken together, this suggests IUP subsistence at Bacho Kiro Cave focused on a
783 range of large to medium-sized herbivores along with the more opportunistic, perhaps
784 occasional hunting, of smaller and larger carnivores including cave bear. At Bacho Kiro

785 Cave the exploitation of carnivore taxa, especially cave bears, appears to be related to the
786 use of these species for fur as well as raw material for bone tools and personal ornaments.
787 At a broader scale, whether this increased focus on carnivores, and small numbers of other
788 lower ranked resources (leporids, birds and fish), represents the beginnings of dietary
789 diversification related to population or dietary pressures requires additional analyses and
790 comparisons to later Upper Palaeolithic sites and faunal assemblages (Morin, 2008, 2012;
791 Stiner, 2009; Starkovich, 2012; Morin et al., 2019). However, this is beyond the scope of this
792 paper which aims to characterise subsistence strategies across the IUP archaeological
793 record.

794 *IUP subsistence behaviour: A spatial and temporal review*

795 The spread, intensity and duration of IUP occupations remains a source of debate (Ginter
796 et al., 2000; Kuhn et al., 2009; Müller et al., 2011; Hublin, 2013, 2015; Kuhn and Zwyns,
797 2014; Ruebens et al., 2015; Hublin et al., 2020) especially since genetic evidence suggests
798 limited genetic transfer from these initial groups to later Upper Palaeolithic and even
799 modern day populations (Hublin et al., 2020; Hajdinjak et al., 2021; Lalueza-Fox, 2021).
800 Further the scarcity of IUP sites with well-preserved faunal remains has limited our
801 understanding of the environmental and climatic tolerances of these early *Homo sapiens*
802 groups, their subsistence behaviour and how this compares to local Neanderthal
803 populations. This is especially problematic for the European record (Figure 15) and
804 therefore Bacho Kiro Cave provides the first well-contextualized, comparative data point of
805 IUP subsistence behaviour in Europe. Nevertheless, in the regions surrounding south-

eastern Europe several extensive and well published sites offer the potential for comparison with new data from Bacho Kiro Cave to begin the process of categorizing and describing IUP subsistence behaviour.

Figure 15: Map showing location of Bacho Kiro Cave and other key Initial Upper Paleolithic (IUP) sites; white dots: sites with good bone preservation; black triangles: sites with poor bone preservation; black dots: sites with no bone preservation.

Figure 15 illustrates some of the major IUP sites, highlighting the problem of poor bone preservation. In Bulgaria the cave sites of Temnata and Kozarnika, which are situated relatively close to Bacho Kiro Cave, have produced layers containing IUP technology and fauna (Ginter et al., 2000; Guadelli and Delpech, 2000; Guadelli et al., 2005; Sirakov et al.,

2010; Tsanova et al., 2021). At Temnata I Layers 4 and VI illustrate a similar range of species including carnivores (cave hyaena [*Crocuta crocuta spelaea*], bear [*Ursus spelaeus*], fox [*Vulpes vulpes*]) alongside a similar range of herbivore taxa, indicative of various environmental types including open (horse [*Equus caballus cf. germanicus*], bison [*Bison priscus*], rhino [*Coelodonta antiquitatis*]), more closed (aurochs [*Bos primigenius*], cervids [*Cervus elaphus*]) and mountainous (ibex [*Capra ibex*], chamois [*Rupicapra rupicapra*]) settings (Guadelli and Delpech, 2000). Faunal representation suggests a similar mix of environments, characteristic of MIS 3, though the presence of the cold-adapted moose (*Alces alces*) within Layer VI in Temnata suggests human occupation, initially, during colder phases as in Bacho Kiro Cave Layer I and J. Bone surface modification data illustrate the use of the cave by human and carnivore groups. The early IUP (Temnata-II Layer VI) is characterized by high carnivore, especially hyaena, (n = 84; 6.8%) input compared to human (n = 34; 3.1%) suggesting a less intensive use of the site and, potentially, repeated periods of human absence. In contrast the later IUP, exemplified by Layer 4 from Temnata-I, has a higher proportion of human (n = 184; 12%) compared to carnivore (n = 6; 0.4%) modifications suggesting a more frequent and perhaps intensive use of the site by human groups. At Kozarnika, similar species representation has been noted (Guadelli, *pers comm*; Guadelli et al. (2005)) though further publication of taphonomic and lithic data is necessary for a more detailed comparison. Overall, these Bulgarian cave sites containing IUP layers illustrate similar species representation and high proportion of human modifications across a range of herbivore and carnivore taxa.

838 Table 8: Initial Upper Palaeolithic sites with published faunal data

ka cal BP	n	NISP	%h	%c	%sm	burn	%cm	%hm	bt	orn	refe
ca. 45-43	5,537	1,091	71	29.1	1.00	0.7	1.4	13	yes	yes	this
ca.46	760	239	27	73.2	0.40	0.4	7.9	8	yes	yes	
48-40	1,530	81	4	0.8	0.00		1.7	51			Gin 202
>45,000	3,683	1,823	50	1.2	0.60	37.4	0.2	11		yes	Bos
undated	272	170	62	1.1	0.02	1.3	0.0	16		yes	
ca. 45-38	3,822	1,264	95	3.4	2.00				yes	yes	Kuh 201
	1,813	654	95	2.1	2.60				yes	yes	
	3,581	1,139	97	2.4	1.10				yes	yes	
	1,114	360	94	2.8	3.00				no	no	

of identified specimens; %h = percentage of herbivore taxa; %c = percentage of carnivore taxa; %sm = percentage of small taxa; %burn = percentage of burnt faunal r
hm = percentage of human modified remains; bt = presence of bone tools; orn = presence of ornaments. Kozarnika Layer VI not included as data are currently unpub

839 Üçağızlı cave (Turkey) preserves a deep sequence spanning most of the Upper Palaeolithic
840 and contains extensive IUP layers (F-I and including sub-layers; Kuhn et al. (2009); Kuhn et
841 al. (1999)] with recent redating suggesting the beginning of the IUP ca. 45,000-43,000 years
842 ago (Kuhn et al., 2009; Douka, 2013). The IUP Layers Fa, Fb, Fc and H1-H3 have abundant
843 ash and charcoal remains and a high anthropogenic input suggesting a more intense or
844 continuous use of the site (Goldberg, 2003). Interestingly, the earliest IUP occupations at
845 Üçağızlı (Layers H, H1-3 and upper I) are represented by thin, sharply defined lenses within
846 these layers suggesting relatively brief events of accumulation (Goldberg, 2003; Kuhn et al.,
847 2009). Overall, the pattern of occupation at Üçağızlı cave during the IUP with more sporadic
848 site use at the beginning of the IUP occupation followed by a more intensive use is similar
849 to that identified at Bacho Kiro Cave in terms of the transition from Layer J into Layer I. IUP

subsistence at Üçağızlı is dominated by medium-large ungulates including fallow deer (*Dama mesopotamica*), bezoar goat (*Capra aegagrus*), and roe deer (*Capreolus capreolus*) with wild boar (*Sus scrofa)*, red deer (*Cervus elaphus*), and aurochs (*Bos primigenius*) present though less abundant. Small game forms a small proportion of the diet in the IUP layers, but is more diversified compared to the Middle Palaeolithic, though the importance of smaller taxa increases significantly in the later Upper Palaeolithic at the site (Kuhn et al., 2009). Again carnivores form a small but consistent portion of the faunal assemblage with human modifications on numerous species including bear and other large carnivores (see Table 8).

The recent redating of Ksâr 'Akil (Lebanon) has produced an age of ca. 45,000 years ago and is associated with IUP lithics and human remains (Bosch et al., 2015; but also see Douka et al., 2015). Detailed zooarchaeological analysis of fauna from Layers XXII and XXV has provided detailed information on the site formation history and shifts in human subsistence behaviour at the site (Bosch, 2018). The burnt bone recovered from the IUP layers illustrates a low proportion of burnt materials in the early IUP Layer XXV (*n* = 3; 1.3%) with a substantial increase within Layer XXII (*n* = 1289; 37.4%). This is a similar to both Bacho Kiro Cave and Üçağızlı, suggesting more intensive use of the cave during the formation of the later IUP in Layer XXII. Fallow deer (*Dama mesopotamica*) are the dominant species throughout the IUP layers, though there is a shift in species diversity. Bosch (2018) demonstrates a decrease in aurochs and wild boar (*Sus scrofa*), which is replaced by caprines (*Capra ibex*, *Capra aegagrus*) and roe deer (*Capreolus capreolus*). Evidence for

marine mollusc consumption occurs initially in Layer XXII though is rarely recorded in the IUP layers with a similarly low exploitation of smaller game, though this again expands in the later Upper Palaeolithic. Further, these layers contain various carnivore taxa, though the absence of associated modifications from the IUP layers (<1%) and the predominance of human butchery modifications (11.8%) suggests that Layers XXII and XXV accumulated as a result of human action.

Overall, this comparative review has highlighted some interesting similarities in IUP subsistence strategies. Comparing new faunal data from Bacho Kiro Cave with other IUP sites suggests that early phases of occupation were characterized by, less intensive or shorter term occupations (for example Layer J at Bacho Kiro, Layers H-I at Üçağızlı, Layers XXV at Ksâr 'Akil). Once established, occupation and site use appears more intensive and perhaps more continuous as exemplified by the large charcoal and ash deposits at sites such as Bacho Kiro Cave and Üçağızlı and the extensive evidence for burnt material in Ksâr 'Akil Layer XXII. This is also illustrated by the increase in density of both lithic and faunal material within later IUP occupations. . Across all sites, IUP subsistence appears to be focused mainly on large-medium sized herbivores with a limited exploitation of small game. The presence of humanly modified carnivore remains appears to be a consistent feature of these faunal assemblages suggesting, perhaps, a more specific focus on these individuals for, presumably, skins and furs but at Bacho Kiro Cave possibly also for dietary resources and raw material/ornaments. Despite the environment and climatic differences observed between these various IUP sites, the observed similarities in terms of human

892 occupation and subsistence behaviour are striking, though further work focused on
893 refining the chronology of the emergence and spread of the IUP is required. Overall, this
894 comparative analysis highlights the importance of faunal remains in understanding
895 similarities in subsistence practice (including diet and carnivore exploitation) across the
896 IUP.

897 **Conclusion**

898 Bacho Kiro Cave represents a key locality for understanding the subsistence behaviour
899 associated with the expansion of Late Pleistocene Homo sapiens into Europe. We identified
900 a change in site use and occupation intensity across the Middle to Upper Palaeolithic
901 transition. IUP Layer I is marked by a significant increase in find density and human
902 modifications on the faunal remains. Together with seasonality data indicating a use of the
903 cave in all seasons, this indicates more frequent and/or more long-term use of the site by
904 these early groups of Homo sapiens. Whether this represents continuous occupation
905 remains unclear, especially as micromorphological data (e.g., clay lenses) has identified
906 potential short periods of site abandonment (Hublin et al., 2020). Furthermore, the
907 presence of wolverine and reindeer suggests, especially within Layer I, that conditions were
908 considerably colder than had been previously considered (Kozłowski and Ginter, 1982;
909 Spassov et al., 2007) and that IUP groups inhabited Europe during varied climatic
910 conditions (Hublin, 2013, 2015; Staubwasser et al., 2018; Zwyns et al., 2019).

911 Across the IUP and the MP layers at Bacho Kiro Cave the dietary focus is similar, centred on
912 the exploitation of a range of large to medium herbivore taxa representing a diverse set of

habitats around the cave site. The large faunal dataset from Layer I allowed for the identification of the selective transport of large herbivore body parts, especially limb bones, into the site. This contrasts with the abundant cave bear remains, including cranial portions, indicating the bears died in the cave itself. Furthermore, a distinct aspect of the IUP at Bacho Kiro Cave is an increase in humanly modified carnivore remains, including both small (wolf, fox) and larger taxa (cave hyaena, cave bear). This relates to procurement of such taxa as raw material for personal ornaments (e.g., cave bear teeth pendants) but also potentially for skins and fur. This pattern of exploitation is distinct from the Neanderthal archaeological record at Bacho Kiro Cave.

At a broader scale, a distinct set of similarities was identified with other IUP sites with faunal preservation across Europe and western Asia. Interestingly, the earliest IUP occupations seem to have been shorter and less intense, as represented by lower proportions of ash and burnt bone, as well as lower proportions of human modifications. During subsequent IUP occupations site use was more intensive and, potentially, for a longer duration to the exclusion of other large carnivores. At all these sites small game exploitation by IUP *Homo sapiens* was limited. The consistent exploitation of carnivore taxa, of a variety of body size classes, for both skins and potentially other resources occurs at all these sites and could be viewed as something distinct for IUP *Homo sapiens*.

Future work should focus on further contextualising the distinct pattern of intense IUP site use and carnivore exploitation identified here. Firstly, within the IUP itself, with additional data coming from sites for which data publication is forthcoming or excavation still

ongoing. This would allow for the further assessment of the role of these subsistence strategies in the spread of *Homo sapiens* groups into Europe and further afield. Secondly, comparison with later Upper Palaeolithic (e.g., Proto-Aurignacian and Early Aurignacian) subsistence strategies would help to understand the specific nature of IUP subsistence and adaptation compared to later groups of incoming *Homo sapiens*. Thirdly, more in-depth comparisons with local MP sites are needed to fully understand the differences between Neanderthal and early *Homo sapiens* subsistence and their role, if any, in population replacements. Overall, with this paper we hope to have created a new data-driven characterisation of IUP subsistence which can now form the basis for a range of comparisons at various temporal and geographic scales.

Acknowledgements

Research and excavations at Bacho Kiro Cave were funded by the Max Planck Society. The re-excavation of Bacho Kiro Cave was jointly conducted by the National Institute of Archaeology with Museum, Bulgarian Academy of Sciences, Sofia (NAIM-BAS) and the Department of Human Evolution at the Max-Planck-Institute for Evolutionary Anthropology, Leipzig (MPI-EVA). We would like to thank all the students that participated in the excavations. GMS and RS would like to especially thank the lab manager Elena Endarova for the excellent curation and storage of the fauna. We would like to thank the Archaeology Department at the New Bulgarian University (Sofia), the Regional Museum of History in Gabrovo and the History Museum in Dryanovo for their ongoing assistance on the project. GMS and RS would especially like to thank Prof. Nikolay Spasov of the National Museum of

Natural History (Sofia) for providing working space and reference material and for his continued assistance and support with the study of the Bacho Kiro Cave fauna. We would also like to acknowledge and thank Jean-Luc Guadelli (PACEA/IPGQ-UMR5199 CNRS, Université Bordeaux I) whom was kind enough to share some of his unpublished faunal data from Kozarnika.NLM was funded through the Fulbright U.S. Scholar Grant (19-11-04) and the Bulgarian-American Commission for Educational Exchange. Karen Ruebens received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No 745662. Finally, we would like to extend our thanks to the Editor of the Journal of Human Evolution, Clément Zanolli, along with the Associate Editor and three anonymous reviewers. Your detailed comments and suggestions have helped to strengthen and improve our manuscript.

References

Abrams, G., Bello, S.M., Di Modica, K., Pirson, S., Bonjean, D., 2014. When Neanderthals used cave bear (Ursus spelaeus) remains: Bone retouchers from unit 5 of Scladina Cave (Belgium). Quaternary International: the journal of the International Union for Quaternary Research. 326-327, 274–287.

Allaire, J.J., Xie, Y., McPherson, J., Luraschi, J., Ushey, K., Atkins, A., Wickham, H., Cheng, J., Chang, W., Iannone, R., 2021. rmarkdown: Dynamic Documents for R.

Behrensmeyer, A.K., 1978. Taphonomic and ecologic information from bone weathering. Paleobiology. 4, 150–162.

Bignon, O., 2006. Approche morphométrique des dents jugales déciduales d'Equus caballus arcelini (sensu lato, Guadelli 1991): critères de détermination et estimation de l'âge d'abattage. Comptes rendus. Palevol. 5, 1005–1020.

Binford, Lewis R., 1981. 1978Nunamiut ethnoarchaeology.

Binford, L.R., 1981. Bones: Ancient Men and Modern Myths. Academic Press, New York.

Blumenschine, R.J., Marean, C.W., Capaldo, S.D., 1996. Blind Tests of Inter-analyst Correspondence and Accuracy in the Identification of Cut Marks, Percussion Marks, and Carnivore Tooth Marks on Bone Surfaces. Journal of Aarchaeological Science. 23, 493–507.

Boaretto, E., Hernandez, M., Goder-Goldberger, M., Aldeias, V., Regev, L., Caracuta, V., McPherron, S.P., Hublin, J.-J., Weiner, S., Barzilai, O., 2021. The absolute chronology of boker tachtit (israel) and implications for the middle to upper paleolithic transition in the levant. Proceedings of the National Academy of Sciences. 118.

Bosch, D.M., 2018. Molluscs in the levantine upper palaeolithic: Implications for modern human diets and subsistence behaviour (PhD thesis). Leiden University.

Bosch, M.D., Mannino, M.A., Prendergast, A.L., O'Connell, T.C., Demarchi, B., Taylor, S.M., Niven, L., Plicht, J. van der, Hublin, J.-J., 2015. New chronology for Ksâr 'Akil (Lebanon) supports Levantine route of modern human dispersal into Europe. Proceedings of the National Academy of Sciences of the United States of America. 112, 7683–7688.

Bunn, H.T., 1981. Archaeological evidence for meat-eating by Plio-Pleistocene hominids from Koobi Fora and Olduvai Gorge. Nature. 291, 574–577.

Cifuentes-Alcobendas, G., Domínguez-Rodrigo, M., 2019. Deep learning and taphonomy: high accuracy in the classification of cut marks made on fleshed and defleshed bones using convolutional neural networks. Scientific Reports. 9, 18933.

Discamps, E., Costamagno, S., 2015. Improving mortality profile analysis in zooarchaeology: a revised zoning for ternary diagrams. Journal of Archaeological Science. 58, 62–76.

1001 Discamps, E., Jaubert, J., Bachellerie, F., 2011. Human choices and environmental
1002 constraints: Deciphering the variability of large game procurement from mousterian to
1003 aurignacian times (MIS 5-3) in southwestern france. Quaternary Science Reviews. 30, 2755–
1004 2775.

1005 Domínguez-Rodrigo, M., 1999. Distinguishing between apples and oranges: the application
1006 of modern cut-mark studies to the plio-pleistocene (a reply to monahan). Journal of Human
1007 Evolution. 37, 793–800.

1008 Domínguez-Rodrigo, M., Saladié, P., Cáceres, I., Huguet, R., Yravedra, J., Rodríguez-Hidalgo,
1009 A., Martín, P., Pineda, A., Marín, J., Gené, C., Aramendi, J., Cobo-Sánchez, L., 2017. Use and
1010 abuse of cut mark analyses: The Rorschach effect. Journal of Archaeological Science. 86,
1011 14–23.

1012 Domínguez-Rodrigo, M., Saladié, P., Cáceres, I., Huguet, R., Yravedra, J., Rodríguez-Hidalgo,
1013 A., Patricia, M., Antonio, P., Juan, M., Clara, G., Aramendi, J., Cobo-Sánchez, L., 2019. Spilled
1014 ink blots the mind: A reply to Merrit et al. (2018) on subjectivity and bone surface
1015 modifications. Journal of Archaeological Science. 102, 80–86.

1016 Douka, K., 2013. Exploring “the great wilderness of prehistory”: The chronology of the
1017 middle to the upper paleolithic transition in the northern levant. Mitteilungen der
1018 Gesellschaft für Urgeschichte. 22.

1019 Douka, K., Higham, T.F., Bergman, C.A., 2015. Statistical and archaeological errors invalidate
1020 the proposed chronology for the site of ksar akil. Proceedings of the National Academy of
1021 Sciences. 112, E7034–E7034.

1022 Fernandez, P., Legendre, S., 2003. Mortality curves for horses from the Middle Palaeolithic
1023 site of Bau de l’Aubesier (Vaucluse, France): methodological, palaeo-ethnological, and
1024 palaeo-ecological approaches. Journal of Archaeological Science. 30, 1577–1598.

1025 Fewlass, H., Talamo, S., Wacker, L., Kromer, B., Tuna, T., Fagault, Y., Bard, E., McPherron,
1026 S.P., Aldeias, V., Maria, R., Martisius, N.L., Paskulin, L., Rezek, Z., Sinet-Mathiot, V., Sirakova,
1027 S., Smith, G.M., Spasov, R., Welker, F., Sirakov, N., Tsanova, T., Hublin, J.-J., 2020. A 14C
1028 chronology for the Middle to Upper Palaeolithic transition at Bacho Kiro Cave, Bulgaria.
1029 Nature Ecology & Evolution. 4, 794–801.

1030 Fisher, J.W., 1995. Bone surface modifications in zooarchaeology. Journal of Archaeological
1031 Method and Theory. 2, 7–68.

1032 Fu, Q., Hajdinjak, M., Moldovan, O.T., Constantin, S., Mallick, S., Skoglund, P., Patterson, N.,
1033 Rohland, N., Lazaridis, I., Nickel, B., others, 2015. An early modern human from romania
1034 with a recent neanderthal ancestor. Nature. 524, 216–219.

Galán, A.B., Rodŕiguez, M., Juana, S. de, Domínguez-Rodrigo, M., 2009. A new experimental study on percussion marks and notches and their bearing on the interpretation of hammerstone-broken faunal assemblages. Journal of Archaeological Science. 36, 776–784.

Garrod, D.A.E., Howe, B., Gaul, J.H., 1939. Excavations in the cave of Bacho Kiro, north-east Bulgaria. Bulletin of the American School of Prehistoric research. 15, 46–76.

Gaudzinski, S., 1999. Middle palaeolithic bone tools from the open-air site salzgitter-lebenstedt (germany). Journal of Archaeological Science. 26, 125–141.

Gaudzinski-Windheuser, S., Noack, E.S., Pop, E., Herbst, C., Pfleging, J., Buchli, J., Jacob, A., Enzmann, F., Kindler, L., Iovita, R., others, 2018. Evidence for close-range hunting by last interglacial neanderthals. Nature ecology & evolution. 2, 1087–1092.

Gifford-Gonzalez, D., 1989b. Shipman's Shaky Foundations. American Anthropologist. 91, 180–186.

Gifford-Gonzalez, D., 1989a. Overview- Modern Analogues: Developing an Interpretive Framework. In: Bonnichsen, R., Sorg, M. (Eds.), Bone Modification. Centre for the Study of the First Americans, Orono, pp. 43–53.

Gifford-Gonzalez, D., 1991. Bones are not enough: Analogues, Knowledge, and Interpretive strategies in zooarchaeology. Journal of Anthropological Archaeology. 10, 215–254.

Ginter, B., Kozlowski, J.K., 1982. Excavation and the stratigraphy of the cave. In: Kozlowski, J.K. (Ed.), Excavation in the Bacho Kiro Cave (Bulgaria): Final Report. pp. 7–12.

Ginter, B., Kozlowski, J.K., Guadelli, J.-L., Laville, H., 2000. Temnata Cave. Excavations in Karlukovo Karst Area, Bulgaria. 2.1. Jagellonian University, Cracow.

Goldberg, P., 2003. Some observations on middle and upper palaeolithic ashy cave and rockshelter deposits in the near east. More Than Meets the Eye: Studies on Upper Palaeolithic Diversity in the Near East. Oxbow, Oxford. 19–32.

Grant, A., 1982. The use of tooth wear as a guide to the age of domestic ungulates. In: Wilson, B., Grigson, C., Payne, S. (Eds.),. BAR, pp. 91–108.

Grayson, D., 1984. Quantitative zooarchaeology: Topics in the quantification of archaeofaunas. Orlando Acad.

Grayson, D.K., Delpech, F., 2003. Ungulates and the Middle-to-Upper Paleolithic transition at Grotte XVI (Dordogne, France). Journal of Archaeological Science. 30, 1633–1648.

Guadelli, A., 2011. Kostni Artefakti ot Paleolita v Bulgaria (Les artefacts en matière dure animale du Paléolithique Bulgare).

Guadelli, J.-L., Delpech, F., 2000. Les grands mammifères des couch VI de la grotte temnata.

Guadelli, J.-L., Sirakov, N., Ivanova, S., Sirakova, S., Anastassova, E., Courtaud, P., Dimitrova, I., Djabarska, N., Fernandez, P., Ferrier, C., others, 2005. Une séquence du paléolithique inférieur au paléolithique récent dans les balkans: La grotte kozarnika à orechets (nord-ouest de la bulgarie). In: Les Premiers Peuplements En Europe. British Archaeological Reports, pp. 87–103.

Guérin, C., 1982. Première biozonation du Pléistocène Européen,principal résultat biostratigraphique de l'étude des Rhinocerotidae (Mammalia, Perissodactyla) du Miocène terminal au Pléistocène supérieur d'Europe Occidentale. Geobios . Memoire special. 15, 593–598.

Hajdinjak, M., Mafessoni, F., Skov, L., Vernot, B., Hübner, A., Fu, Q., Essel, E., Nagel, S., Nickel, B., Richter, J., Moldovan, O.T., Constantin, S., Endarova, E., Zahariev, N., Spasov, R., Welker, F., Smith, G.M., Sinet-Mathiot, V., Paskulin, L., Fewlass, H., Talamo, S., Rezek, Z., Sirakova, S., Sirakov, N., McPherron, S.P., Tsanova, T., Hublin, J.-J., Peter, B.M., Meyer, M., Skoglund, P., Kelso, J., Pääbo, S., 2021. Initial Upper Palaeolithic humans in Europe had recent Neanderthal ancestry. Nature. 592, 253–257.

Hamilton, N.E., Ferry, M., 2018. ggtern: Ternary Diagrams Using ggplot2. Journal of Statistical Software, Code Snippets.

Higham, T., Douka, K., Wood, R., Ramsey, C.B., Brock, F., Basell, L., Camps, M., Arrizabalaga, A., Baena, J., Barroso-Ruíz, C., Bergman, C., Boitard, C., Boscato, P., Caparrós, M., Conard, N.J., Draily, C., Froment, A., Galván, B., Gambassini, P., Garcia-Moreno, A., Grimaldi, S., Haesaerts, P., Holt, B., Iriarte-Chiapusso, M.-J., Jelinek, A., Jordá Pardo, J.F., Maíllo-Fernández, J.-M., Marom, A., Maroto, J., Menéndez, M., Metz, L., Morin, E., Moroni, A., Negrino, F., Panagopoulou, E., Peresani, M., Pirson, S., Rasilla, M. de la, Riel-Salvatore, J., Ronchitelli, A., Santamaria, D., Semal, P., Slimak, L., Soler, J., Soler, N., Villaluenga, A., Pinhasi, R., Jacobi, R., 2014. The timing and spatiotemporal patterning of Neanderthal disappearance. Nature. 512, 306–309.

Hublin, J.-J., 2013. The Makers of the Early Upper Paleolithic in Western Eurasia. In: Smith, F.H., Ahern, J.C.M. (Eds.), The Origins of Modern Humans. John Wiley & Sons, Inc, Hoboken, NJ, pp. 223–252.

Hublin, J.-J., 2015. The modern human colonization of western Eurasia: when and where? Quaternary Science Reviews. 118, 194–210.

Hublin, J.-J., Sirakov, N., Aldeias, V., Bailey, S., Bard, E., Delvigne, V., Endarova, E., Fagault, Y., Fewlass, H., Hajdinjak, M., Kromer, B., Krumov, I., Marreiros, J., Martisius, N.L., Paskulin, L., Sinet-Mathiot, V., Meyer, M., Pääbo, S., Popov, V., Rezek, Z., Sirakova, S., Skinner, M.M., Smith, G.M., Spasov, R., Talamo, S., Tuna, T., Wacker, L., Welker, F., Wilcke, A., Zahariev, N., McPherron, S.P., Tsanova, T., 2020. Initial Upper Palaeolithic Homo sapiens from Bacho Kiro Cave, Bulgaria. Nature. 581, 299–302.

Iovita, R., Schönekeß, H., Gaudzinski-Windheuser, S., Jäger, F., 2014. Projectile impact fractures and launching mechanisms: results of a controlled ballistic experiment using replica Levallois points. Journal of Archaeological Science. 48, 73–83.

Kassambara, A., 2020. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.

Kindler, L., 2012. Die Rolle von Raubtieren in der Einnischung und Subsistenz jungpleistozäner Neandertaler. Archäozoologie und Taphonomie der mittelpaläolithischen Fauna aus der Balver Höhle (Westfalen). Monographien des Römisch-Germanischen Zentralmuseums. 99.

Kozłowski, J.K., Ginter, B., 1982. Excavation in the Bacho Kiro Cave (Bulgaria): Final Report. Państwowe Wydawnictwo Naukowe.

Kreutzer, L.A., 1992. Bison and deer bone mineral densities: Comparisons and implications for the interpretation of archaeological faunas. Journal of Archaeological Science. 19, 271–294.

Kruuk, H., 1972. The Spotted Hyena: a study of predation and social behavior.

Kuhn, S.L., Stiner, M.C., Güleç, E., 1999. Initial Upper Palaeolithic in south-central Turkey and its regional context: a preliminary report. Antiquity. 73, 505–517.

Kuhn, S.L., Stiner, M.C., Güleç, E., Ozer, I., Yilmaz, H., Baykara, I., Açikkol, A., Goldberg, P., Molina, K.M., Unay, E., Suata-Alpaslan, F., 2009. The early Upper Paleolithic occupations at Uçağizli Cave (Hatay, Turkey). Journal of Human Evolution. 56, 87–113.

Kuhn, S.L., Stiner, M.C., Reese, D.S., Güleç, E., 2001. Ornaments of the earliest Upper Paleolithic: new insights from the Levant. Proceedings of the National Academy of Sciences of the United States of America. 98, 7641–7646.

Kuhn, S.L., Zwyns, N., 2014. Rethinking the initial Upper Paleolithic. Quaternary international: the journal of the International Union for Quaternary Research. 347, 29–38.

Lalueza-Fox, C., 2021. Neanderthal assimilation? Nature Ecology & Evolution. 1–2.

Lam, Y.M., Chen, X., Marean, C.W., Frey, C.J., 1998. Bone Density and Long Bone Representation in Archaeological Faunas: Comparing Results from CT and Photon Densitometry. Journal of Archaeological Science. 25, 559.

Lam, Y.M., Chen, X., Pearson, O.M., 1999. Intertaxonomic Variability in Patterns of Bone Density and the Differential Representation of Bovid, Cervid, and Equid Elements in the Archaeological Record. American Antiquity. 64, 343–362.

Lam, Y.M., Pearson, O.M., 2005. Bone density studies and the interpretation of the faunal record. Evolutionary Anthropology: Issues, News, and Reviews. 14, 99–108.

Lam, Y.M., Pearson, O.M., Marean, C.W., Chen, X., 2003. Bone density studies in zooarchaeology. Journal of Archaeological Science. 30, 1701.

Levine, M.A., 1982. The use of crown height measurements and eruption-wear sequences to age horse teeth. In: Wilson, B., Grigson, C. (Eds.),. BAR.

Lyman, R., 1994. Vertebrate Taphonomy. Cambridge University Press, Cambridge.

Lyman, R.L., 2008. Quantitative paleozoology. Cambridge University Press.

Made, J. van der, 2018. Quaternary Large-Mammal Zones. In: López Varela, S.L. (Ed.), The Encyclopedia of Archaeological Sciences. John Wiley & Sons, Inc., Hoboken, NJ, USA, pp. 1–4.

Marin, J., Daujeard, C., Saladié, P., Rodŕiguez-Hidalgo, A., Vettese, D., Rivals, F., Boulbes, N., Crégut-Bonnoure, E., Lateur, N., Gallotti, R., others, 2020. Neanderthal faunal exploitation and settlement dynamics at the abri du maras, level 5 (south-eastern france). Quaternary Science Reviews. 243, 106472.

Marks, A.E., 1990. The Middle and Upper Palaeolithic of the Near East and the Nile Valley: the problem of cultural transformations. In: The emergence of modern humans: An archaeological perspective. pp. 56–80.

Moclán, A., Domínguez-Rodrigo, M., Yravedra, J., 2019. Classifying agency in bone breakage: an experimental analysis of fracture planes to differentiate between hominin and carnivore dynamic and static loading using machine learning (ML) algorithms. Archaeological and Anthropological Sciences.

Morin, E., 2008. Evidence for declines in human population densities during the early Upper Paleolithic in western Europe. Proceedings of the National Academy of Sciences of the United States of America. 105, 48–53.

Morin, E., 2012. Reassessing Paleolithic Subsistence: The Neandertal and Modern Human Foragers of Saint-Césaire. Cambridge University Press.

Morin, E., Laroulandie, V., 2012. Presumed symbolic use of diurnal raptors by neanderthals. PLoS One. 7, e32856.

Morin, E., Meier, J., El Guennouni, K., Moigne, A.-M., Lebreton, L., Rusch, L., Valensi, P., Conolly, J., Cochard, D., 2019. New evidence of broader diets for archaic Homo populations in the northwestern Mediterranean. Science Advances. 5, eaav9106.

Morin, E., Ready, E., Boileau, A., Beauval, C., Coumont, M.-P., 2017b. Problems of Identification and Quantification in Archaeozoological Analysis, Part II: Presentation of an Alternative Counting Method. Journal of Archaeological Method and Theory. 24, 938–973.

Morin, E., Ready, E., Boileau, A., Beauval, C., Coumont, M.-P., 2017a. Problems of Identification and Quantification in Archaeozoological Analysis, Part I: Insights from a Blind Test. Journal of Archaeological Method and Theory. 24, 886–937.

Müller, U.C., Pross, J., Tzedakis, P.C., Gamble, C., Kotthoff, U., Schmiedl, G., Wulf, S., Christanis, K., 2011. The role of climate in the spread of modern humans into Europe. Quaternary Science Reviews. 30, 273–279.

Münzel, S.C., Conard, N.J., 2004. Cave bear hunting in the Hohle Fels, a cave site in the Ach Valley, Swabian Jura. Revue de Paléobiologie. 23, 877–885.

Newcomer, M.H., 1974. Study and replication of bone tools from ksar akil (lebanon). World archaeology. 6, 138–153.

NEWCOMER, M., WATSON, J., 1984. Bone artifacts from ksar'aqil (lebanon). Paléorient. 143–147.

O'Driscoll, C.A., Thompson, J.C., 2014. Experimental projectile impact marks on bone: implications for identifying the origins of projectile technology. Journal of Archaeological Science. 49, 398–413.

Pederzani, S., Britton, K., Aldeias, V., Bourgon, N., Fewlass, H., McPherron, S.P., Rezek, Z., Sirakov, N., Smith, G.M., Spasov, R., Tran, N.-H., Tsanova, T., Hublin, J.-J., 2021. Subarctic climate for the earliest Homo sapiens in Europe. Science Advances. 7.

Peresani, M., Fiore, I., Gala, M., Romandini, M., Tagliacozzo, A., 2011. Late neandertals and the intentional removal of feathers as evidenced from bird bone taphonomy at fumane cave 44 ky BP, italy. Proceedings of the National Academy of Sciences. 108, 3888–3893.

Pizarro-Monzo, M., Domínguez-Rodrigo, M., 2020. Dynamic modification of cut marks by trampling: temporal assessment through the use of mixed-effect regressions and deep learning methods. Archaeological and Anthropological Sciences. 12, 4.

QGIS Development Team, 2009. QGIS Geographic Information System. Open Source Geospatial Foundation.

R Core Team, 2020. R: A Language and Environment for Statistical Computing.

Radovčić, D., Sršen, A.O., Radovčić, J., Frayer, D.W., 2015. Evidence for Neandertal jewelry: modified white-tailed eagle claws at Krapina. PloS One. 10, e0119802.

Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Ramsey, C.B., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Others, 2020. The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). Radiocarbon. 62, 725–757.

Reitz, E.J., Reitz, E., Wing, E.S., 1999. Zooarchaeology. Cambridge University Press.

Rendu, W., Renou, S., Soulier, M.-C., Rigaud, S., Roussel, M., Soressi, M., 2019. Subsistence strategy changes during the Middle to Upper Paleolithic transition reveals specific adaptations of Human Populations to their environment. Scientific Reports. 9, 15817.

Romandini, M., Peresani, M., Laroulandie, V., Metz, L., Pastoors, A., Vaquero, M., Slimak, L., 2014. Convergent evidence of eagle talons used by late Neanderthals in Europe: a further assessment on symbolism. PloS one. 9, e101278.

Romandini, M., Terlato, G., Nannini, N., Tagliacozzo, A., Benazzi, S., Peresani, M., 2018. Bears and humans, a Neanderthal tale. Reconstructing uncommon behaviors from zooarchaeological evidence in southern Europe. Journal of Archaeological Science. 90, 71–91.

Rots, V., Plisson, H., 2014. Projectiles and the abuse of the use-wear method in a search for impact. Journal of Archaeological Science. 48, 154–165.

RStudio Team, 2021. RStudio: Integrated Development Environment for R.

Ruebens, K., McPherron, S.J.P., Hublin, J.-J., 2015. On the local Mousterian origin of the Châtelperronian: Integrating typo-technological, chronostratigraphic and contextual data. Journal of Human Evolution. 86, 55–91.

Sirakov, N., Guadelli, J.-L., Ivanova, S., Sirakova, S., Boudadi-Maligne, M., Dimitrova, I., Ph, F., Ferrier, C., Guadelli, A., Iordanova, D., others, 2010. An ancient continuous human presence in the balkans and the beginnings of human settlement in western eurasia: A lower pleistocene example of the lower palaeolithic levels in kozarnika cave (north-western bulgaria). Quaternary International. 223, 94–106.

Skrdla, P., 2014. Moravian Bohunician. In: Otte, M. (Ed.), Neandertal/Cro-Magnon: La rencontre. Errance, Paris, pp. 123–147.

Smith, G.M., 2003. Damage inflicted on animal bone by wooden projectiles: experimental results and archaeological implications. Journal of Taphonomy. 1, 105–114.

Smith, G.M., 2010. A contextual approach to the study of faunal assemblages from Lower and Middle Palaeolithic sites in the UK (PhD thesis). UCL (University College London); discovery.ucl.ac.uk, London.

Smith, G.M., 2013. Taphonomic resolution and hominin subsistence behaviour in the Lower Palaeolithic: differing data scales and interpretive frameworks at Boxgrove and Swanscombe (UK). Journal of Archaeological Science. 40, 3754–3767.

Smith, G.M., 2015. Neanderthal megafaunal exploitation in Western Europe and its dietary implications: a contextual reassessment of La Cotte de St Brelade (Jersey). Journal of Human Evolution. 78, 181–201.

Smith, G.M., Noack, E.S., Behrens, N.M., Ruebens, K., Street, M., Iovita, R., Gaudzinski-Windheuser, S., 2020. When Lithics Hit Bones: Evaluating the Potential of a Multifaceted Experimental Protocol to Illuminate Middle Palaeolithic Weapon Technology. Journal of Paleolithic Archaeology.

Soressi, M., McPherron, S.P., Lenoir, M., Dogandžić, T., Goldberg, P., Jacobs, Z., Maigrot, Y., Martisius, N.L., Miller, C.E., Rendu, W., others, 2013. Neandertals made the first specialized bone tools in europe. Proceedings of the National Academy of Sciences. 110, 14186–14190.

Spassov, N., Popov, V., Miteva, S., Mihova, B., Georgiev, K., Petrov, B., Vansink, D., 2007. History of the formation of the Bulgarian mammal fauna. Miteva, S. Mihova, B. , Georgiev, K. , Petrov, B. , Vansink, D. (eds.), The mammals, important for conservation in Bulgaria. Dutch Mammal Society VZZ. Arnhem. 31–46.

Starkovich, B.M., 2012. Intensification of small game resources at klissoura cave 1 (peloponnese, greece) from the middle paleolithic to mesolithic. Quaternary International. 264, 17–31.

Starkovich, B.M., 2017. Paleolithic subsistence strategies and changes in site use at klissoura cave 1 (peloponnese, greece). Journal of human evolution. 111, 63–84.

Staubwasser, M., Drăgușin, V., Onac, B.P., Assonov, S., Ersek, V., Hoffmann, D.L., Veres, D., 2018. Impact of climate change on the transition of Neanderthals to modern humans in Europe. Proceedings of the National Academy of Sciences of the United States of America. 115, 9116–9121.

Steele, T.E., 2004. Variation in mortality profiles of red deer (Cervus elaphus) in Middle Palaeolithic assemblages from western Europe. International Journal of Osteoarchaeology. 14, 307–320.

Steele, T.E., Weaver, T.D., 2002. The Modified Triangular Graph: A Refined Method for Comparing Mortality Profiles in Archaeological Samples. Journal of Archaeological Science. 29, 317–322.

Steele, T.E., Weaver, T.D., 2012. Refining the Quadratic Crown Height Method of age estimation: do elk teeth wear quadratically with age? Journal of Archaeological Science. 39, 2329–2334.

Stiner, M., 2009. Prey choice, site occupation intensity & economic diversity in the middle–early upper palaeolithic at the Üçağizli caves, turkey. Before Farming. 2009, 1–21.

Stiner, M.C., 1990. The use of mortality patterns in archaeological studies of hominid predatory adaptations. Journal of Anthropological Archaeology. 9, 305–351.

Stiner, M.C., 1991a. The faunal remains from grotta guattari: A taphonomic perspective. Current Anthropology. 32, 103–117.

Stiner, M.C., 1991b. Food procurement and transport by human and non-human predators. Journal of Archaeological Science. 18, 455–482.

Stiner, M.C., 1994. Honor Among Thieves: a zooarchaeological study of Neanderthal ecology. Princeton University Press, Princeton.

Stiner, M.C., 1998. Mortality analysis of Pleistocene bears and its paleoanthropological relevance. Journal of Human Evolution. 34, 303–326.

Stiner, M.C., Kuhn, S.L., Weiner, S., Bar-Yosef, O., 1995. Differential Burning, Recrystallization, and Fragmentation of Archaeological Bone. Journal of Archaeological Science. 22, 223–237.

Svoboda, J., Bar-Yosef, O., 2003. Stránská skála. Origins of the upper paleolithic in the brno basin, moravia, czech republic. Peabody Museum of Archaeology; Ethnology, Harvard University, Cambridge.

Tsanova, T., 2008. Les débuts du Paléolithique supérieur dans l'Est des Balkans Réflexion à partir de l'étude taphonomique et techno-économique des ensembles lithiques des sites de Bacho Kiro (couche 11), Temnata (couches VI et 4) et Kozarnika (niveau VII), BAR international series 1752. BAR.

Tsanova, T., 2012. A diachronic view of flake production from the beginning of the Upper Palaeolithic in the eastern Balkans. Flakes not Blades: The role of flake production at the onset of the Upper Paleolithic in Europe. 215–237.

Tsanova, T., Bordes, J.-G., 2003. Contribution to the d é bat on the origin of the Aurignacian: main results of a technological study of the lithic industry of layer 11 of Bacho Kiro. na.

Tsanova, T., Veres, D., Hambach, U., Spasov, R., Dimitrova, I., Popov, P., Talamo, S., Sirakova, S., 2021. Upper palaeolithic layers and campanian ignimbrite/y-5 tephra in toplitsa cave, northern bulgaria. Journal of Archaeological Science: Reports. 37, 102912.

VanPool, T.L., Leonard, R.D., 2011. Quantitative Analysis in Archaeology. John Wiley & Sons.

Weinstock, J., 2009. Epiphyseal fusion in brown bears: a population study of grizzlies (Ursus arctos horribilis) from Montana and Wyoming. International Journal of Osteoarchaeology. 19, 416–423.

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. Journal of Open Source Software. 4, 1686.

Xie, Y., Allaire, J.J., Grolemund, G., 2018. R Markdown: The Definitive Guide.

Xie, Y., Dervieux, C., Riederer, E., 2020. R Markdown Cookbook.

Zwyns, N., Gladyshev, S.A., Gunchinsuren, B., Bolorbat, T., Flas, D., Dogandžić, T., Tabarev, A.V., Gillam, J.C., Khatsenovich, A.M., McPherron, S., Odsuren, D., Paine, C.H., Purevjal, K.-E., Stewart, J.R., 2014. The open-air site of Tolbor 16 (Northern Mongolia): Preliminary results and perspectives. Quaternary International: the journal of the International Union for Quaternary Research. 347, 53–65.

Zwyns, N., Paine, C.H., Tsedendorj, B., Talamo, S., Fitzsimmons, K.E., Gantumur, A., Guunii, L., Davakhuu, O., Flas, D., Dogandžić, T., Doerschner, N., Welker, F., Gillam, J.C., Noyer, J.B., Bakhtiary, R.S., Allshouse, A.F., Smith, K.N., Khatsenovich, A.M., Rybin, E.P., Byambaa, G., Hublin, J.-J., 2019. The Northern Route for Human dispersal in Central and Northeast Asia: New evidence from the site of Tolbor-16, Mongolia. Scientific Reports. 9, 11759.