

# *Generalized second order vectorial $\infty$ -eigenvalue problems*

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Clark, E. and Katzourakis, N. (2024) Generalized second order vectorial  $\infty$ -eigenvalue problems. *Proceedings of the Royal Society of Edinburgh A: Mathematics*. ISSN 1473-7124 doi: 10.1017/prm.2024.27 Available at <https://centaur.reading.ac.uk/119241/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/prm.2024.27>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading



# Generalized second order vectorial $\infty$ -eigenvalue problems

Ed Clark and Nikos Katzourakis

Department of Mathematics and Statistics, University of Reading,  
Whiteknights Campus, Pepper Lane, Reading RG6 6AX, United  
Kingdom ([e.d.clark@pgr.reading.ac.uk](mailto:e.d.clark@pgr.reading.ac.uk); [n.katzourakis@reading.ac.uk](mailto:n.katzourakis@reading.ac.uk))

(Received 28 March 2023; accepted 30 September 2023)

We consider the problem of minimizing the  $L^\infty$  norm of a function of the hessian over a class of maps, subject to a mass constraint involving the  $L^\infty$  norm of a function of the gradient and the map itself. We assume zeroth and first order Dirichlet boundary data, corresponding to the “hinged” and the “clamped” cases. By employing the method of  $L^p$  approximations, we establish the existence of a special  $L^\infty$  minimizer, which solves a divergence PDE system with measure coefficients as parameters. This is a counterpart of the Aronsson-Euler system corresponding to this constrained variational problem. Furthermore, we establish upper and lower bounds for the eigenvalue.

*Keywords:* calculus of variations in  $L^\infty$ ;  $\infty$ -eigenvalue problem; nonlinear eigenvalue problems; absolute minimizers; Lagrange multipliers

2020 *Mathematics Subject Classification* Primary: 35P30  
Secondary: 35D30, 35J94, 35P15

## 1. Introduction and main results

Let  $n, N \in \mathbb{N}$  with  $n \geq 2$ , and let  $\Omega \Subset \mathbb{R}^n$  be a bounded open set with Lipschitz boundary  $\partial\Omega$ . In this paper we are interested in studying nonlinear second order  $L^\infty$  eigenvalue problems. Specifically, we investigate the problem of finding a minimizing map  $u_\infty : \bar{\Omega} \rightarrow \mathbb{R}^N$ , that solves

$$\begin{aligned} \|f(D^2u_\infty)\|_{L^\infty(\Omega)} &= \inf \left\{ \|f(D^2v)\|_{L^\infty(\Omega)} : \right. \\ &\quad \left. v \in W_B^{2,\infty}(\Omega; \mathbb{R}^N), \|g(v, Dv)\|_{L^\infty(\Omega)} = 1 \right\}. \end{aligned} \quad (1.1)$$

Additionally, we pursue the necessary conditions that these constrained minimizers must satisfy, in the form of PDEs. In the above,  $f : \mathbb{R}_s^{N \times n^2} \rightarrow \mathbb{R}$  and  $g : \mathbb{R}^N \times \mathbb{R}^{N \times n} \rightarrow \mathbb{R}$  are given functions, that will be required to satisfy some natural assumptions, to be discussed later in this section. We note that  $\mathbb{R}_s^{N \times n^2}$  symbolizes the symmetric subspace of the tensor space  $\mathbb{R}^N \otimes (\mathbb{R}^n \otimes \mathbb{R}^n)$  wherein the Hessians of twice differentiable maps  $u : \Omega \rightarrow \mathbb{R}^N$  are valued. The functional

Sobolev space  $W_B^{2,\infty}(\Omega; \mathbb{R}^N)$  appearing above will be taken to be either of:

$$\begin{cases} W_C^{2,\infty}(\Omega; \mathbb{R}^N) : & = W_0^{2,\infty}(\Omega; \mathbb{R}^N), \\ W_H^{2,\infty}(\Omega; \mathbb{R}^N) : & = W^{2,\infty} \cap W_0^{1,\infty}(\Omega; \mathbb{R}^N). \end{cases} \quad (1.2)$$

The space  $W_C^{2,\infty}(\Omega; \mathbb{R}^N)$  encompasses the case of so-called clamped boundary conditions, which can be seen as first order Dirichlet or as coupled Dirichlet–Neumann conditions, requiring  $|u| = |\nabla u| = 0$  on  $\partial\Omega$ . On the other hand,  $W_H^{2,\infty}(\Omega; \mathbb{R}^N)$  encompasses the so-called hinged boundary conditions, which are zeroth order Dirichlet conditions, requiring  $u = 0$  on  $\partial\Omega$ . This is standard terminology for such problems, see e.g. [25].

Problem (1.1) lies within the Calculus of Variations in  $L^\infty$ , a modern area, initiated by Gunnar Aronsson in the 1960s. Since then this field has undergone a substantial transformation. There are some general complications one must be wary of when tackling  $L^\infty$  variational problems. For example, the  $L^\infty$  norm is generally not Gateaux differentiable, therefore the analogue of the Euler–Lagrange equations cannot be derived directly by considering variations. Any supremal functional also has issues with locality in terms of minimization on subdomains. Further, the space itself lacks some fundamental functional analytic properties, such as reflexivity and separability. Higher order problems and problems involving constraints present additional difficulties and have been studied even more sparsely, see e.g. [3, 4, 9, 10, 20–24, 26]. In fact, this paper is an extension of [23] to the second order case, and generalizes part of the results corresponding to the existence of minimizers and the satisfaction of PDEs from [25]. In turn, the paper [23] generalized results on the scalar case of eigenvalue problems for the  $\infty$ -Laplacian ([18, 19]). For various interesting results, see for instance [2, 3, 6, 8, 28–31].

The vectorial and higher order nature of the problem we are considering herein precludes the use of standard methods, such as viscosity solutions (see e.g. [1] for a pedagogical introduction). However, we overcome these difficulties by approximating by corresponding  $L^p$  problems for finite  $p$  case and let  $p \rightarrow \infty$ . The intuition for using this technique is based on the rudimentary idea that, for a fixed  $L^\infty$  function on a set of finite measure, its  $L^p$  norm tends to its  $L^\infty$  norm as  $p \rightarrow \infty$ . This technique is rather standard for  $L^\infty$  problems, and in the vectorial higher order case we consider herein is essentially the only method known. Even the very recent intrinsic duality method of [6] is limited to scalar-valued first order problems.

To state our main result, we now introduce the required hypotheses for the functions  $f$  and  $g$ :

$$\left\{ \begin{array}{l} (a) f \in C^1(\mathbb{R}_s^{N \times n^2}). \\ (b) f \text{ is (Morrey) 2-quasiconvex.} \\ (c) \text{There exist } 0 < C_1 \leq C_2 \text{ such that, for all } X \in \mathbb{R}_s^{N \times n^2} \setminus \{0\}, \\ \quad 0 < C_1 f(X) \leq \partial f(X) : X \leq C_2 f(X). \\ (d) \text{There exist } C_3, \dots, C_6 > 0, \alpha > 1 \text{ and } \beta \leq 1 : \text{for all } X \in \mathbb{R}_s^{N \times n^2}, \\ \quad -C_3 + C_4 |X|^\alpha \leq f(X) \leq C_5 |X|^\alpha + C_6, \\ \quad |\partial f(X)| \leq C_5 f(X)^\beta + C_6. \end{array} \right. \quad (1.3)$$

$$\left\{ \begin{array}{l} (a) \quad g \in C^1(\mathbb{R}^N \times \mathbb{R}^{N \times n}). \\ (b) \quad g \text{ is coercive, in the sense that} \\ \lim_{t \rightarrow \infty} \left( \inf_{(\eta, P) \in \mathbb{R}^N \times \mathbb{R}^{N \times n}, |(\eta, P)|=1} g(t\eta, tP) \right) = \infty. \\ (c) \quad \text{There exist } 0 < C_7 \leq C_8 : \text{ for all } (\eta, P) \in (\mathbb{R}^N \times \mathbb{R}^{N \times n}) \setminus \{(0, 0)\}, \\ 0 < C_7 g(\eta, P) \leq \partial_\eta g(\eta, P) \cdot \eta + \partial_P g(\eta, P) : P \leq C_8 g(\eta, P). \end{array} \right. \quad (1.4)$$

In the above,  $\partial f(X)$  denotes the the derivative of  $f$  whilst  $\partial_\eta g$  and  $\partial_P g$  signifies the respective partial derivatives. Additionally ‘:’ and ‘‘.’’ represent the Euclidean inner products. The terminology of (Morrey) 2-quasiconvex refers to the standard notion for integral functionals for higher order functionals (see e.g. [7, 11, 12, 33]), namely

$$F(X) \leq \int_{\Omega} F(X + D^2\phi) d\mathcal{L}^n, \quad \forall \phi \in W_0^{2,\infty}(\Omega; \mathbb{R}^N), \quad \forall X \in \mathbb{R}_s^{N \times n^2}.$$

We note that herein we will be using the following function space symbolizations:

$$C_B^2(\overline{\Omega}; \mathbb{R}^N) := C^2(\overline{\Omega}; \mathbb{R}^N) \cap W_B^{2,\infty}(\Omega; \mathbb{R}^N),$$

$$W_C^{2,p}(\Omega; \mathbb{R}^N) := W_0^{2,p}(\Omega; \mathbb{R}^N), \quad p \in [1, \infty),$$

$$W_H^{2,p}(\Omega; \mathbb{R}^N) := W^{2,p} \cap W_0^{1,p}(\Omega; \mathbb{R}^N), \quad p \in [1, \infty),$$

Further, we will be using the rescaled  $L^p$  norms for  $p \in [1, \infty)$ , given by

$$\|h\|_{L^p(\Omega)} := \left( \frac{1}{\mathcal{L}^n(\Omega)} \int_{\Omega} |h|^p d\mathcal{L}^n \right)^{\frac{1}{p}} = \left( \int_{\Omega} |h|^p d\mathcal{L}^n \right)^{\frac{1}{p}}.$$

Finally, we observe that (1.3)(c), implies that  $f > 0$  on  $\mathbb{R}_s^{N \times n^2} \setminus \{0\}$ ,  $f(0) = 0$  and  $f$  is radially increasing, meaning that  $t \mapsto f(tX)$  is increasing on  $(0, \infty)$  for any fixed  $X \in \mathbb{R}_s^{N \times n^2} \setminus \{0\}$ . Similarly, (1.4)(c) implies that  $g > 0$  on  $(\mathbb{R}^N \times \mathbb{R}^{N \times n}) \setminus \{(0, 0)\}$ ,  $g(0, 0) = 0$  and  $g$  is radially increasing on  $\mathbb{R}^N \times \mathbb{R}^{N \times n}$ , namely  $t \mapsto g(t\eta, tP)$  is increasing on  $(0, \infty)$  for any fixed  $(\eta, P) \in (\mathbb{R}^N \times \mathbb{R}^{N \times n}) \setminus \{(0, 0)\}$ .

Below is our main result, in which we consider both cases of boundary conditions simultaneously.

**THEOREM 1.1.** *Suppose that the assumptions (1.3) and (1.4) hold true. Then:*

(A) *The problem (1.1) has a solution  $u_\infty \in W_B^{2,\infty}(\Omega; \mathbb{R}^N)$ .*

(B) *There exist Radon measures*

$$M_\infty \in \mathcal{M}(\overline{\Omega}; \mathbb{R}_s^{N \times n^2}), \quad \nu_\infty \in \mathcal{M}(\overline{\Omega}),$$

*such that*

$$\int_{\overline{\Omega}} D^2\phi : dM_\infty = \Lambda_\infty \int_{\overline{\Omega}} \left( \partial_\eta g(u_\infty, Du_\infty) \cdot \phi + \partial_P g(u_\infty, Du_\infty) : D\phi \right) d\nu_\infty \quad (1.5)$$

for all test maps  $\phi \in C_B^2(\bar{\Omega}; \mathbb{R}^N)$ , where

$$\Lambda_\infty = \|f(D^2(u_\infty))\|_{L^\infty(\Omega)} > 0. \quad (1.6)$$

Additionally, we have the following a priori lower bound for the eigenvalue

$$\Lambda_\infty \geq \left( \frac{C_4}{\text{diam}(\Omega)^\alpha \left( C(\infty, \Omega) \|\partial_\eta g\|_{L^\infty(\{g \leq 1\})} + \|\partial_P g\|_{L^\infty(\{g \leq 1\})} \right)^\alpha} - C_3 \right)^+, \quad (1.7)$$

where  $(\cdot)^+$  symbolizes the positive part, and  $C(\infty, \Omega)$  equals either the constant of the Poincaré inequality (in the case of clamped boundary conditions), or the constant of the Poincaré-Wirtinger inequality (in the case of hinged boundary conditions), both taken for  $p = \infty$ .

If additionally the boundary  $\partial\Omega$  is  $C^2$ , we have the a priori upper bound

$$\begin{aligned} \Lambda_\infty &\leq C_6 + C_5 \frac{2^{5\alpha}}{(c\omega(n))^\alpha} \left( 1 + \sup_{0 \leq t \leq 1} R(t) \right)^\alpha \left( 2^{3n} + \max_{i=1, \dots, n-1} (\|\kappa_i\|_{C^0(\partial\Omega)})^n \right)^\alpha \cdot \\ &\quad \cdot \left\{ 1 + \left( 1 + \frac{C}{\varepsilon_0^{n+1}} \right) \mathcal{H}^{n-1}(\partial\Omega) + \sum_{i=1}^{n-1} \left\| \frac{\kappa_i \circ P_\Omega}{1 - (\kappa_i \circ P_\Omega) d_\Omega} \right\|_{L^\infty(\{d_\Omega < \varepsilon_0\} \cap \Omega)} \right\}^\alpha, \end{aligned} \quad (1.8)$$

where  $c, C > 0$  are dimensionless universal constants,  $\omega(n)$  is the volume of the unit ball in  $\mathbb{R}^n$ ,  $\mathcal{H}^{n-1}(\partial\Omega)$  is the perimeter of  $\Omega$ ,  $\{\kappa_1, \dots, \kappa_{n-1}\}$  are the principal curvatures of  $\partial\Omega$ ,  $P_\Omega$  is the orthogonal projection on  $\partial\Omega$ ,  $d_\Omega$  the distance function of  $\partial\Omega$ ,  $\varepsilon_0$  is the largest

$$\varepsilon \in \left( 0, \min \left\{ 1, \min_{i=1, \dots, n-1} \frac{1}{\|\kappa_i\|_{C^0(\partial\Omega)}} \right\} \right),$$

for which we have that  $d_\Omega \in C^2(\{d_\Omega \leq \varepsilon\} \cap \bar{\Omega})$  and  $R(t)$  is the smallest radius of the  $N$ -dimensional ball, for which the sublevel set  $\{g \leq t\}$  is contained into the cylinder  $\bar{\mathbb{B}}_{R(t)}^N(0) \times \mathbb{R}^{N \times n}$ , namely

$$R(t) := \inf \left\{ R > 0 : \{g \leq t\} \subseteq \bar{\mathbb{B}}_R^N(0) \times \mathbb{R}^{N \times n} \right\}.$$

(C) The quadruple  $(u_\infty, \Lambda_\infty, M_\infty, \nu_\infty)$  satisfies the following approximation properties: there exists a sequence of exponents  $(p_j)_1^\infty \subseteq (n/\alpha)$  where  $p_j \rightarrow \infty$  as  $j \rightarrow \infty$ , and for any  $p$ , a quadruple

$$(u_p, \Lambda_p, M_p, \nu_p) \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N) \times [0, \infty) \times \mathcal{M}(\bar{\Omega}; \mathbb{R}_s^{N \times n^2}) \times \mathcal{M}(\bar{\Omega}),$$

such that

$$\begin{cases} u_p \rightharpoonup u_\infty, & \text{in } C^1(\overline{\Omega}; \mathbb{R}^N), \\ D^2 u_p \rightharpoonup D^2 u_\infty, & \text{in } L^q(\Omega; \mathbb{R}_s^{N \times n^2}), \text{ for all } q \in (1, \infty), \\ \Lambda_p \rightharpoonup \Lambda_\infty, & \text{in } [0, \infty), \\ M_p \xrightarrow{*} M_\infty, & \text{in } \mathcal{M}(\overline{\Omega}; \mathbb{R}_s^{N \times n^2}), \\ \nu_p \xrightarrow{*} \nu_\infty, & \text{in } \mathcal{M}(\overline{\Omega}), \end{cases} \quad (1.9)$$

as  $p \rightarrow \infty$  along  $(p_j)_1^\infty$ . Further,  $u_p$  solves the constrained minimization problem

$$\|f(D^2 u_p)\|_{L^p(\Omega)} = \inf \left\{ \|f(D^2 v)\|_{L^p(\Omega)} : v \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N), \|g(v, Dv)\|_{L^p(\Omega)} = 1 \right\}, \quad (1.10)$$

and  $(u_p, \Lambda_p)$  satisfies

$$\begin{cases} \int_{\Omega} f(D^2 u_p)^{p-1} \partial f(D^2 u_p) : D^2 \phi \, d\mathcal{L}^n \\ = (\Lambda_p)^p \int_{\Omega} g(u_p, Du_p)^{p-1} \left( \partial_\eta g(u_p, Du_p) \cdot \phi + \partial_P g(u_p, Du_p) : D\phi \right) \, d\mathcal{L}^n \end{cases} \quad (1.11)$$

for all test maps  $\phi \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ . Finally, the measures  $M_p, \nu_p$  are given by

$$\begin{cases} M_p &= \frac{1}{\mathcal{L}^n(\Omega)} \left( \frac{f(D^2 u_p)}{\Lambda_p} \right)^{p-1} \partial f(D^2 u_p) \mathcal{L}^n \llcorner_{\Omega}, \\ \nu_p &= \frac{1}{\mathcal{L}^n(\Omega)} g(u_p, Du_p)^{p-1} \mathcal{L}^n \llcorner_{\Omega}. \end{cases} \quad (1.12)$$

We note that one could pursue optimality in theorem 1.1(A) by using  $L^\infty$  versions of quasiconvexity, as developed by Barron–Jensen–Wang [5] but adapted to this higher order case, in regards to the existence of  $L^\infty$  minimizers. However, for parts (B) and (C) of theorem 1.1 regarding the necessary PDE conditions, we do need Morrey 2-quasiconvexity, as we rely essentially on the existence of solutions to the corresponding Euler–Lagrange equations and the theory of Lagrange multipliers in the finite  $p$  case. Further, the measures  $M_\infty, \nu_\infty$  depend on the minimizer  $u_\infty$  in a non-linear fashion, hence one more could perhaps symbolize them more concisely as  $M_\infty(u_\infty), \nu_\infty(u_\infty)$ . Consequently, the significance of these equations is currently understood to be mostly of conceptual value, rather than of computational nature. However, it is possible to obtain further information about the underlying structure of these parametric measure coefficients. This requires techniques such as measure function pairs and mollifications up to the boundary as in [10, 17, 23], but to keep the presentation as simple as possible, we refrain from pursuing this -considerably more technical- endeavour, which also requires stronger assumptions.

## 2. Proofs

In this section we establish theorem 1.1. Its proof is not labelled explicitly, but will be completed by proving a combination of smaller subsidiary results, including a selection of lemmas and propositions.

Before introducing the approximating problem (the  $L^p$  case for finite  $p$ ), we need to establish a convergence result, which shows that the admissible classes of the  $p$ -problems are non-empty. It is required because the function  $g$  appearing in the constraint is not assumed to be homogeneous, therefore a standard scaling argument does not suffice.

LEMMA 2.1. *For any  $v \in W_B^{2,\infty}(\Omega; \mathbb{R}^N) \setminus \{0\}$ , there exists  $(t_p)_{p \in (n/\alpha, \infty]}$  with  $t_p \rightarrow t_\infty$  as  $p \rightarrow \infty$ , such that*

$$\|g(t_p v, t_p Dv)\|_{L^p(\Omega)} = 1,$$

for all  $p \in (n/\alpha, \infty]$ . Further, if  $\|g(v, Dv)\|_{L^\infty(\Omega)} = 1$ , then  $t_\infty = 1$ .

*Proof of lemma 2.1.* Fix  $v \in W_B^{2,\infty}(\Omega; \mathbb{R}^N) \setminus \{0\}$  and define

$$\rho_\infty(t) := \max_{x \in \bar{\Omega}} g(tv(x), tDv(x)), \quad t \geq 0.$$

It follows that  $\rho_\infty(0) = 0$  and  $\rho_\infty$  is continuous on  $[0, \infty)$ . We will now show that  $\rho_\infty$  is strictly increasing. We first show it is non-decreasing. For any  $s > 0$  and  $(\eta, P) \in \mathbb{R}^N \times \mathbb{R}^{N \times n} \setminus \{(0, 0)\}$ , our assumption (1.4)(c) implies

$$\begin{aligned} 0 &< \frac{C_7 g(s\eta, sP)}{s} \\ &\leq \partial_\eta g(s\eta, sP) \cdot \eta + \partial_P g(s\eta, sP) : P \\ &= \partial_{(\eta, P)} g(s\eta, sP) : (\eta, P) \\ &= \frac{d}{ds} (g(s\eta, sP)), \end{aligned}$$

thus  $s \mapsto g(s\eta, sP)$  is increasing on  $(0, \infty)$ . Hence, for any  $x \in \bar{\Omega}$  and  $t > s \geq 0$  we have  $g(tv(x), tDv(x)) \geq g(sv(x), sDv(x))$ , which yields,

$$\rho_\infty(s) = \max_{x \in \bar{\Omega}} g(sv(x), sDv(x)) \leq \max_{x \in \bar{\Omega}} g(tv(x), tDv(x)) = \rho_\infty(t).$$

We proceed to demonstrate that  $t \mapsto \rho_\infty(t)$  is actually strictly monotonic over  $(0, \infty)$ . Fix  $t_0 > 0$ . By Danskin's theorem [13], the derivative from the right  $\rho'(t_0^+)$  exists, and is given by the formula

$$\rho'_\infty(t_0^+) = \max_{x \in \Omega_{t_0}} \left\{ \partial_{(\eta, P)} g(t_0 v(x), t_0 Dv(x)) : (v(x), Dv(x)) \right\},$$

where

$$\Omega_{t_0} := \left\{ \bar{x} \in \bar{\Omega} : \rho_\infty(t_0) = g(t_0 v(\bar{x}), t_0 Dv(\bar{x})) \right\}.$$

Hence, by (1.4)(c) we estimate

$$\begin{aligned}
 \rho'_\infty(t_0^+) &= \frac{1}{t_0} \max_{x \in \Omega_{t_0}} \left\{ \partial_{(\eta, P)} g(t_0 v(x), t_0 Dv(x)) : (t_0 v(x), t_0 Dv(x)) \right\} \\
 &\geq \frac{C_7}{t_0} \max_{x \in \Omega_{t_0}} g(t_0 v(x), t_0 Dv(x)) \\
 &= \frac{C_7}{t_0} \rho_\infty(t_0) \\
 &> 0.
 \end{aligned}$$

This implies that  $\rho_\infty$  is strictly increasing on  $(0, \infty)$ . Next, recall that  $g$  is coercive by assumption (1.4)(b), namely  $g(s\eta, sP) \rightarrow \infty$  as  $s \rightarrow \infty$ , for fixed  $(\eta, P) \neq (0, 0)$ . Thus, for any fixed point  $\bar{x} \in \Omega$  with  $(v(\bar{x}), Dv(\bar{x})) \neq (0, 0)$ , which exists because by assumption  $v \not\equiv 0$ , we have

$$\lim_{t \rightarrow \infty} \rho_\infty(t) \geq \lim_{t \rightarrow \infty} g(tv(\bar{x}), tDv(\bar{x})) = \infty.$$

Since  $\rho_\infty(0) = 0$  and  $\rho_\infty(t) \rightarrow \infty$  as  $t \rightarrow \infty$ , by continuity and the intermediate value theorem, there exists a number  $t_\infty > 0$  such that  $\rho_\infty(t_\infty) = 1$ , that is

$$\|g(t_\infty v, t_\infty Dv)\|_{L^\infty(\Omega)} = 1.$$

If  $\|g(v, Dv)\|_{L^\infty(\Omega)} = 1$ , then  $t_\infty = 1$ , as a result of the strict monotonicity of  $\rho_\infty$ . Now let us fix  $p \in (n/\alpha, \infty)$  and define the continuous function

$$\rho_p(t) := \int_{\Omega} g(tv, tDv)^p d\mathcal{L}^n, \quad t \geq 0.$$

Since  $g(0, 0) = 0$ , it follows that  $\rho_p(0) = 0$  and that

$$\rho_p(t) = \frac{1}{\mathcal{L}^n(\Omega)} \int_{\{(v, Dv) \neq (0, 0)\}} g(tv, tDv)^p d\mathcal{L}^n.$$

By Morrey's theorem and our assumptions, we have that  $v \in C^1(\bar{\Omega}; \mathbb{R}^N) \setminus \{0\}$ , therefore  $\mathcal{L}^n(\{(v, Dv) \neq (0, 0)\}) > 0$ . Consider the family of functions  $\{g(tv, tDv)^p\}_{t>0}$ , defined on  $\{(v, Dv) \neq (0, 0)\} \subseteq \Omega$ . By the monotonicity of  $s \mapsto g(s\eta, sP)$  on  $(0, \infty)$  for  $(\eta, P) \neq (0, 0)$ , for  $s < t$  we have

$$g(sv, sDv)^p \leq g(tv, tDv)^p, \quad \text{on } \{(v, Dv) \neq (0, 0)\}.$$

Since  $g(tv, tDv)^p \rightarrow \infty$  pointwise on  $\{(v, Dv) \neq (0, 0)\}$  as  $t \rightarrow \infty$ , by the monotone convergence theorem, we infer that

$$\int_{\{(v, Dv) \neq (0, 0)\}} g(tv, tDv)^p d\mathcal{L}^n \rightarrow \infty,$$

as  $t \rightarrow \infty$ . As a consequence,  $\rho_p(t) \rightarrow \infty$  as  $t \rightarrow \infty$ . Since  $\rho_p(0) = 0$ , by the intermediate value theorem there exists  $t_p > 0$  such that  $\rho_p(t_p) = 1$ , namely

$$\|g(t_p v, t_p Dv)\|_{L^p(\Omega)} = 1.$$

For the sake of contradiction, suppose that  $t_p \neq t_\infty$ , as  $p \rightarrow \infty$ . In this case, there exists a subsequence  $(t_{p_j})_1^\infty \subseteq (n/\alpha, \infty)$  and  $t_0 \in [0, t_\infty) \cup (t_\infty, \infty]$  such that  $t_{p_j} \rightarrow$

$t_0$  as  $j \rightarrow \infty$ . Further,  $(t_{p_j})_1^\infty$  can be assumed to be either monotonically increasing or decreasing. We first prove that  $t_0$  is finite. If  $t_0 = \infty$ , then the sequence  $(t_{p_j})_1^\infty$  can be selected to be monotonically increasing. Therefore, by arguing as before,  $g(t_{p_j}v, t_{p_j}Dv) \nearrow \infty$  as  $j \rightarrow \infty$ , pointwise on  $\{(v, Dv) \neq (0, 0)\}$ , and the monotone convergence theorem provides the contradiction

$$1 = \lim_{j \rightarrow \infty} \int_{\Omega} g(t_{p_j}v, t_{p_j}Dv)^{p_j} d\mathcal{L}^n = \int_{\Omega} \lim_{j \rightarrow \infty} g(t_{p_j}v, t_{p_j}Dv)^{p_j} d\mathcal{L}^n = \infty.$$

Consequently, we have that  $t_0 \in [0, t_\infty) \cup (t_\infty, \infty)$ . Since  $(t_{p_j}v, t_{p_j}Dv) \rightarrow (t_0v, t_0Dv)$  uniformly on  $\bar{\Omega}$  as  $j \rightarrow \infty$ , we calculate

$$\begin{aligned} 1 &= \|g(t_{p_j}v, t_{p_j}Dv)\|_{L^{p_j}(\Omega)} \\ &= \|g(t_0v, t_0Dv)\|_{L^{p_j}(\Omega)} + o(1)_{j \rightarrow \infty} \\ &= \|g(t_0v, t_0Dv)\|_{L^\infty(\Omega)} + o(1)_{j \rightarrow \infty} \\ &= \rho_\infty(t_0) + o(1)_{j \rightarrow \infty}. \end{aligned}$$

By passing to the limit as  $j \rightarrow \infty$ , we obtain a contradiction if  $t_\infty \neq t_0$ , because  $\rho_\infty$  is a strictly increasing function and  $\rho_\infty(t_\infty) = 1$ . In conclusion,  $t_p \rightarrow t_\infty$  as  $p \rightarrow \infty$ .  $\square$

Utilizing the above result we can now show existence for the approximating minimization problem for  $p < \infty$ .

LEMMA 2.2. *For any  $p > n/\alpha$ , the minimization problem (1.10) has a solution  $u_p \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ .*

*Proof of lemma 2.2.* Let us fix  $p \in (n/\alpha, \infty)$  and  $v_0 \in W_B^{2,\infty}(\Omega; \mathbb{R}^N)$  where  $v_0 \not\equiv 0$ . By application of lemma 2.1, there exists  $t_p > 0$  such that  $\|g(t_p v_0, t_p Dv_0)\|_{L^p(\Omega)} = 1$  implying that  $t_p v_0$  is indeed an element of the admissible class of (1.10). Hence, we deduce that the admissible class is non empty. Further, by assumption (1.3)(b),  $f$  is (Morrey) 2-quasiconvex. We now confirm that  $f^p$  is also (Morrey) 2-quasiconvex function, as a consequence of Jensen's inequality: for any fixed  $X \in \mathbb{R}_s^{N \times n^2}$  and any  $\phi \in W_0^{2,\infty}(\Omega; \mathbb{R}^N)$ , we have

$$f^p(X) \leq \left( \int_{\Omega} f(X + D^2\phi) d\mathcal{L}^n \right)^p \leq \int_{\Omega} f(X + D^2\phi)^p d\mathcal{L}^n.$$

By assumption by assumption (1.3)(d), we have for some new  $C_5(p)$ ,  $C_6(p) > 0$  that

$$f(X)^p \leq C_5(p)|X|^{\alpha p} + C_6(p),$$

for any  $X \in \mathbb{R}_s^{N \times n^2}$ . Moreover, by [33, Theorem 3.6] we have that the functional  $v \mapsto \|f(D^2v)\|_{L^p(\Omega)}$  is weakly lower semi-continuous on  $W^{2,\alpha p}(\Omega; \mathbb{R}^N)$  and therefore the same is true over the closed subspace  $W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ . Let  $(u_i)_1^\infty$  be a minimizing sequence for (1.10). As  $f \geq 0$ , it is clear that  $\inf_{i \in \mathbb{N}} \|f(D^2u_i)\|_{L^p(\Omega)} \geq 0$ . Since the

admissible class is non-empty, the infimum is finite. Additionally, by (1.3)(d), we have the bound

$$\begin{aligned} \inf_{i \in \mathbb{N}} \|f(D^2 u_i)\|_{L^p(\Omega)} &\leq \|f(D^2(t_p v_0))\|_{L^p(\Omega)} \\ &\leq \|C_5 |t_p D^2 v_0|^\alpha + C_6\|_{L^\infty(\Omega)} \\ &\leq C_5(t_p)^\alpha \|D^2 v_0\|_{L^\infty(\Omega)}^\alpha + C_6 \\ &< \infty. \end{aligned}$$

Now we show that the functional is coercive in  $W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ , arguing separately for either case of boundary conditions. By assumption (1.3)(d) and the Poincaré inequality, for any  $u \in W_C^{2,\alpha p}(\Omega; \mathbb{R}^N)$  (satisfying  $|u| = |Du| = 0$  on  $\partial\Omega$ ), we have

$$\left( \int_{\Omega} |f(D^2 u) + C_3|^p d\mathcal{L}^n \right)^{\frac{1}{p}} \geq C_4 \left( \int_{\Omega} |D^2 u|^{\alpha p} d\mathcal{L}^n \right)^{\frac{1}{p}} \geq C'_4 \|u\|_{W^{1,\alpha p}(\Omega)}^\alpha,$$

for a new constant  $C'_4 = C_4(p) > 0$ . Hence, for any  $u \in W_C^{2,\alpha p}(\Omega; \mathbb{R}^N)$ ,

$$\|f(D^2 u)\|_{L^p(\Omega)} \geq C'_4 (\|u\|_{W^{2,\alpha p}(\Omega)})^\alpha - C_3. \quad (2.1)$$

The above estimate is also true when  $u \in W_H^{2,\alpha p}(\Omega; \mathbb{R}^N)$ , but since in this case we have only  $|u| = 0$  on  $\partial\Omega$ , it requires an additional justification. By the Poincaré-Wirtinger inequality involving averages, for any  $u \in W_H^{2,\alpha p}(\Omega; \mathbb{R}^N)$  we have

$$\left\| Du - \int_{\Omega} Du d\mathcal{L}^n \right\|_{L^{\alpha p}(\Omega)} \leq C \|D^2 u\|_{L^{\alpha p}(\Omega)},$$

where  $C = C(\alpha, p, \Omega) > 0$  is a constant. Since  $|u| = 0$  on  $\partial\Omega$ , by the Gauss-Green theorem we have

$$\int_{\Omega} Du d\mathcal{L}^n = \int_{\partial\Omega} u \otimes \hat{n} d\mathcal{H}^{n-1} = 0,$$

where  $\mathcal{H}^{n-1}$  denotes the  $(n-1)$ -dimensional Hausdorff measure. In conclusion,

$$\|Du\|_{L^{\alpha p}(\Omega)} \leq C \|D^2 u\|_{L^{\alpha p}(\Omega)},$$

for any  $u \in W_H^{2,\alpha p}(\Omega; \mathbb{R}^N)$ . The above estimate together with the standard Poincaré inequality applied to  $u$  itself allow to infer that (2.1) holds for any  $u \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$  in both cases of boundary conditions. Returning to our minimizing sequence, by standard compactness results, exists  $u_p \in W_H^{2,\alpha p}(\Omega; \mathbb{R}^N)$  such that  $u_i \rightharpoonup u_p$  in  $W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ , as  $i \rightarrow \infty$  along a subsequence of indices. Additionally, by the Morrey estimate we have that  $u_i \rightharpoonup u_p$  in  $C^1(\bar{\Omega}; \mathbb{R}^N)$  as  $i \rightarrow \infty$ , along perhaps a further subsequence. Since  $u \mapsto \|g(u, Du)\|_{L^p(\Omega)}$  is weakly continuous on  $W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ , the admissible class is weakly closed in  $W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$  and hence we may pass to the limit in the constraint. By weak lower semicontinuity of the functional, it follows that a minimizer  $u_p$  which satisfies (1.10) does indeed exist.  $\square$

Now we describe the necessary conditions (Euler–Lagrange equations) that approximating minimizer  $u_p$  must satisfy. These equations will involve a Lagrange multiplier, emerging from the constraint  $\|g(\cdot, D(\cdot))\|_{L^p(\Omega)} = 1$ .

LEMMA 2.3. *For any  $p > n/\alpha$ , let  $u_p$  be the minimizer of (1.10) procured by lemma 2.2. Then, there exists  $\lambda_p \in \mathbb{R}$  such that the pair  $(u_p, \lambda_p)$  satisfies the following PDE system*

$$\begin{aligned} & \int_{\Omega} f(D^2 u_p)^{p-1} \partial f(D^2 u_p) : D^2 \phi \, d\mathcal{L}^n \\ &= \lambda_p \int_{\Omega} g(u_p, Du_p)^{p-1} \left( \partial_{\eta} g(u_p, Du_p) \cdot \phi + \partial_P g(u_p, Du_p) : D\phi \right) \, d\mathcal{L}^n, \end{aligned}$$

for all test maps  $\phi \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$ .

In particular, it follows that in both cases  $u_p$  is a weak solution in  $W^{2,\alpha p}(\Omega; \mathbb{R}^N)$  to

$$\begin{cases} D^2 : \left( f(D^2 u_p)^{p-1} \partial f(D^2 u_p) \right) \\ = \lambda_p \left[ g(u_p, Du_p)^{p-1} \partial_{\eta} g(u_p, Du_p) - \operatorname{div} \left( g(u_p, Du_p)^{p-1} \partial_P g(u_p, Du_p) \right) \right], \end{cases} \quad (2.2)$$

where we have used the notation  $D^2 : F = \sum_{i,j=1}^n D_{ij}^2 F_{ij}$ , when  $F \in C^2(\Omega; \mathbb{R}^{n \times n})$ , which is equivalent to the double divergence (applied once column-wise and once row-wise). Note that in the case of hinged boundary data, we have an additional natural boundary condition arising (since  $Du$  is free on  $\partial\Omega$ ), we we will not make an particular use of this extra information in the sequel, therefore we refrain from discussing it explicitly.

*Proof of lemma 2.3.* The result follows by standard results on Lagrange multipliers in Banach spaces (see e.g. [32, p. 278]), by utilizing assumption (1.3)(d), which guarantees that the functional is Gateaux differentiable.  $\square$

Now we establish some further results regarding the family of eigenvalues.

LEMMA 2.4. *Consider the family of pairs of eigenvectors-eigenvalues  $\{(u_p, \lambda_p)\}_{p > n/\alpha}$ , given by lemma 2.3. Then, for any  $p > n/\alpha$ , there exists  $\Lambda_p > 0$  such that*

$$\lambda_p = (\Lambda_p)^p > 0.$$

Further, by setting

$$L_p := \|f(D^2 u_p)\|_{L^p(\Omega)},$$

we have the bounds

$$0 < \left( \frac{C_1}{C_8} \right)^{\frac{1}{p}} L_p \leq \Lambda_p \leq \left( \frac{C_2}{C_7} \right)^{\frac{1}{p}} L_p.$$

*Proof of lemma 2.4.* We begin by showing that  $L_p > 0$ , namely the infimum over the admissible class of the  $p$ -approximating minimization problem is strictly positive, owing to the constraint and our assumptions (1.3)-(1.4). Indeed, there is only one map  $u \in W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$  for which  $\|f(D^2u)\|_{L^p(\Omega)} = 0$ , namely  $u_0 \equiv 0$ , but this is not an element of the admissible class since  $\|g(u_0, Du_0)\|_{L^p(\Omega)} = 0$ . Now consider the Euler–Lagrange equations in lemma 2.3 and select  $\phi := u_p$ , to obtain

$$\begin{aligned} & \int_{\Omega} f(D^2u_p)^{p-1} \partial f(D^2u_p) : D^2u_p \, d\mathcal{L}^n \\ &= \lambda_p \int_{\Omega} g(u_p, Du_p)^{p-1} \left( \partial_{\eta} g(u_p, Du_p) \cdot u_p + \partial_P g(u_p, Du_p) : Du_p \right) \, d\mathcal{L}^n. \end{aligned}$$

As  $f, g \geq 0$  we can manipulate the respective assumptions (1.3)(c) and (1.4)(c) to produce the following bounds:

$$\begin{aligned} C_1 \int_{\Omega} f(D^2u_p)^p \, d\mathcal{L}^n &\leq \int_{\Omega} f(D^2u_p)^{p-1} \partial f(D^2u_p) : D^2u_p \, d\mathcal{L}^n \\ &\leq C_2 \int_{\Omega} f(D^2u_p)^p \, d\mathcal{L}^n, \\ C_7 \int_{\Omega} g(u_p, Du_p)^p \, d\mathcal{L}^n &\leq \int_{\Omega} g(u_p, Du_p)^{p-1} \left( \partial_{\eta} g(u_p, Du_p) \cdot u_p + \right. \\ &\quad \left. + \partial_P g(u_p, Du_p) : Du_p \right) \, d\mathcal{L}^n \\ &\leq C_8 \int_{\Omega} g(u_p, Du_p)^p \, d\mathcal{L}^n. \end{aligned}$$

The above two estimates, combined with the Euler–Lagrange equations, imply that  $\lambda_p > 0$ . Hence, we may therefore define  $\Lambda_p := (\lambda_p)^{\frac{1}{p}} > 0$ . We will now obtain the upper and lower bounds. We determine the lower bound as follows:

$$\begin{aligned} C_1(L_p)^p &= C_1 \int_{\Omega} f(D^2u_p)^p \, d\mathcal{L}^n \\ &\leq \int_{\Omega} f^{p-1}(D^2u_p) \partial f(D^2u_p) : D^2u_p \, d\mathcal{L}^n \\ &= \lambda_p \int_{\Omega} g(u_p, Du_p)^{p-1} \left( \partial_{\eta} g(u_p, Du_p) \cdot \phi + \partial_P g(u_p, Du_p) : Du_p \right) \, d\mathcal{L}^n \\ &\leq \lambda_p C_8. \end{aligned}$$

Hence,

$$\left( \frac{C_1}{C_8} \right)^{\frac{1}{p}} L_p \leq (\lambda_p)^{\frac{1}{p}} = \Lambda_p.$$

The upper bound is determined analogously, by reversing the direction of the inequalities. Combining both bounds, we obtain the desired estimate.  $\square$

PROPOSITION 2.5. *There exists  $(u_\infty, \Lambda_\infty) \in W_B^{2,\infty}(\Omega; \mathbb{R}^N) \times (0, \infty)$  such that, along a sequence  $(p_j)_1^\infty$  of exponents, we have*

$$\begin{cases} u_p \rightharpoonup u_\infty, & \text{in } C^1(\bar{\Omega}; \mathbb{R}^N), \\ D^2 u_p \rightharpoonup D^2 u_\infty, & \text{in } L^q(\Omega; \mathbb{R}_s^{N \times n^2}), \text{ for all } q \in (1, \infty), \\ \Lambda_p \rightharpoonup \Lambda_\infty, & \text{in } [0, \infty), \end{cases}$$

as  $p_j \rightarrow \infty$ . Additionally,  $u_\infty$  solves the minimization problem (1.1) and  $\Lambda_\infty$  is given by (1.6). Finally  $\Lambda_\infty$  satisfies the uniform bounds (1.7).

*Proof of proposition 2.5.* Fix  $p > n/\alpha$ ,  $q \leq p$  and a map  $v_0 \in W_B^{2,\infty}(\Omega; \mathbb{R}^N) \setminus \{0\}$ . Then, by lemma 2.1 there exists  $(t_p)_{p \in (n/\alpha, \infty]} \subseteq (0, \infty)$  such that  $t_p \rightarrow t_\infty$  as  $p \rightarrow \infty$  and satisfying  $\|g(t_p v_0, t_p Dv_0)\|_{L^p(\Omega)} = 1$  for all  $p \in (n/\alpha, \infty]$ . By Hölder's inequality and minimality, we have the following estimate

$$\begin{aligned} \|f(D^2 u_p)\|_{L^q(\Omega)} &\leq \|f(D^2 u_p)\|_{L^p(\Omega)} \\ &\leq \|f(t_p D^2 v_0)\|_{L^p(\Omega)} \\ &\leq \|f(t_p D^2 v_0)\|_{L^\infty(\Omega)} \\ &\leq K + \|f(t_\infty D^2 v_0)\|_{L^\infty(\Omega)} \\ &< \infty, \end{aligned}$$

for some  $K > 0$ . By (1.3)(d), we have the bound  $f^q(X) \geq C_4(q)|X|^{\alpha q} - C_3(q)$  for some constants  $C_3(q)$ ,  $C_4(q) > 0$  and all  $X \in \mathbb{R}_s^{N \times n^2}$ . By the previous bound, we conclude that

$$\sup_{q \geq p} \|D^2 u_p\|_{L^{\alpha q}(\Omega)} \leq C(q) < \infty,$$

for some  $q$ -dependent constant. By arguing as in the proof of lemma 2.2 through the use of Poincaré inequalities, we can conclude in both cases of boundary conditions with the bound

$$\sup_{q \geq p} \|u_p\|_{W^{2,\alpha q}(\Omega)} \leq C(q) < \infty,$$

for a new  $q$ -dependent constant  $C'(q) > 0$ . Standard compactness in Sobolev spaces and a diagonal sequence argument imply the existence of a mapping

$$u_\infty \in \bigcap_{n/\alpha < p < \infty} W_B^{2,\alpha p}(\Omega; \mathbb{R}^N)$$

and a subsequence  $(p_j)_1^\infty$  such that the desired modes of convergence hold true as  $p_j \rightarrow \infty$  along this subsequence of indices. Fix a map  $v \in W_B^{2,\infty}(\Omega; \mathbb{R}^N)$  satisfying the required constraint, namely  $\|g(v, Dv)\|_{L^\infty(\Omega)} = 1$ . In view of lemma 2.1, there exists  $(t_p)_{p \in (n/\alpha, \infty)} \subseteq (0, \infty)$  satisfying that  $t_p \rightarrow 1$  as  $p \rightarrow \infty$ , and additionally  $\|g(t_p v, t_p Dv)\|_{L^p(\Omega)} = 1$  for all  $p > n/\alpha$ . By Hölder's inequality, the definition of

$L_p$  and minimality, we have

$$\|f(D^2u_p)\|_{L^q(\Omega)} \leq \|f(D^2u_p)\|_{L^p(\Omega)} = L_p \leq \|f(t_p D^2v)\|_{L^p(\Omega)},$$

for any such  $v$ . By the weak lower semi-continuity of the functional on  $W_B^{2,\alpha q}(\Omega; \mathbb{R}^N)$ , we may let  $p_j \rightarrow \infty$  to obtain

$$\begin{aligned} \|f(D^2u_\infty)\|_{L^q(\Omega)} &\leq \liminf_{p_j \rightarrow \infty} L_p \\ &\leq \limsup_{p_j \rightarrow \infty} L_p \\ &\leq \limsup_{p_j \rightarrow \infty} \|f(t_{p_j} D^2v)\|_{L^p(\Omega)} \\ &= \|f(D^2v)\|_{L^\infty(\Omega)}. \end{aligned}$$

Now we may let  $q \rightarrow \infty$  in the above bound, hence producing

$$\|f(D^2u_\infty)\|_{L^\infty(\Omega)} \leq \liminf_{p_j \rightarrow \infty} L_p \leq \limsup_{p_j \rightarrow \infty} L_p \leq \|f(D^2v)\|_{L^\infty(\Omega)}.$$

for all mappings  $v \in W_B^{2,\infty}(\Omega; \mathbb{R}^N)$  satisfying the constraint  $\|g(v, Dv)\|_{L^\infty(\Omega)} = 1$ . If we additionally show that in fact  $u_\infty$  satisfies the constraint in (1.1), then the above estimate shows both that it is the desired minimizers (by choosing  $v := u_\infty$ ), and also that the sequence  $(L_{p_j})_1^\infty$  converges to the infimum. Now we show that this is indeed the case. In view of assumption (1.3)(d), the previous estimate implies also that  $D^2u_\infty \in L^\infty(\Omega; \mathbb{R}_s^{N \times n^2})$ , which together with Poincaré inequalities (as in the proof of lemma 2.2) shows that in fact  $u_\infty \in W_B^{2,\infty}(\Omega; \mathbb{R}^N)$ . By the continuity of the function  $g$  and the fact that  $u_p \rightarrow u_\infty$  in  $C^1(\bar{\Omega}; \mathbb{R}^N)$ , we have

$$\begin{aligned} 1 &= \|g(u_p, Du_p)\|_{L^p(\Omega)} \\ &= \|g(u_\infty, Du_\infty)\|_{L^p(\Omega)} + \|g(u_p, Du_p)\|_{L^p(\Omega)} - \|g(u_\infty, Du_\infty)\|_{L^p(\Omega)} \\ &= \|g(u_\infty, Du_\infty)\|_{L^p(\Omega)} + O\left(\|g(u_p, Du_p) - g(u_\infty, Du_\infty)\|_{L^\infty(\Omega)}\right) \\ &\longrightarrow \|g(u_\infty, Du_\infty)\|_{L^\infty(\Omega)}, \end{aligned}$$

as  $p_j \rightarrow \infty$ . Consequently,  $u_\infty$  satisfies the constraint, and therefore lies in the admissible class of (1.1). Since  $v$  was arbitrary in the energy bound, we conclude that  $u_\infty$  in fact solves (1.1). let us now define

$$\Lambda_\infty := \|f(D^2u_\infty)\|_{L^\infty(\Omega)}.$$

We now show that  $\Lambda_\infty > 0$ . Indeed, by our assumptions (1.3)–(1.4), there is only one map in  $W^{2,\infty}(\Omega; \mathbb{R}^N)$  satisfying  $\|f(D^2u_0)\|_{L^\infty(\Omega)} = 0$  and  $|u_0| \equiv 0$  on  $\partial\Omega$ , namely the trivial map  $u_0 \equiv 0$ , but  $u_0$  is not contained in the admissible class of (1.1) because  $\|g(u_0, Du_0)\|_{L^\infty(\Omega)} = 0$ . We now show that  $\Lambda_p \rightarrow \Lambda_\infty$  as  $p_j \rightarrow \infty$ . By our

earlier energy estimate, we have  $L_p \rightarrow \Lambda_\infty$  as  $p_j \rightarrow \infty$ . By lemma 2.4, we have

$$0 < \lim_{p_j \rightarrow \infty} \left( \frac{C_1}{C_8} \right)^{\frac{1}{p}} L_p \leq \lim_{p_j \rightarrow \infty} \Lambda_p \leq \lim_{p_j \rightarrow \infty} \left( \frac{C_2}{C_7} \right)^{\frac{1}{p}} L_p,$$

and therefore  $\Lambda_p \rightarrow \Lambda_\infty$  as  $p_j \rightarrow \infty$ . To complete the proof we must deduce the claimed bounds for  $\Lambda_\infty$ . We first establish the lower bound. By utilizing the Poincaré and Poincaré-Wirtinger inequalities (recall the proof of lemma 2.2) and that  $g(0, 0) = 0$ , we estimate

$$\begin{aligned} 1 &= \|g(u_\infty, Du_\infty)\|_{L^\infty(\Omega)} \\ &\leq \text{diam}(\Omega) \|D(g(u_\infty, Du_\infty))\|_{L^\infty(\Omega)} \\ &\leq \text{diam}(\Omega) \left( \|\partial_\eta g(u_\infty, Du_\infty) Du_\infty\|_{L^\infty(\Omega)} + \|\partial_P g(u_\infty, Du_\infty) D^2 u_\infty\|_{L^\infty(\Omega)} \right) \\ &\leq \text{diam}(\Omega) \left( \|\partial_\eta g\|_{L^\infty((u_\infty, Du_\infty)(\bar{\Omega}))} \|Du_\infty\|_{L^\infty(\Omega)} \right. \\ &\quad \left. + \|\partial_P g\|_{L^\infty((u_\infty, Du_\infty)(\bar{\Omega}))} \|D^2 u_\infty\|_{L^\infty(\Omega)} \right) \\ &\leq \|D^2 u_\infty\|_{L^\infty(\Omega)} \text{diam}(\Omega) \left( C(\infty, \Omega) \|\partial_\eta g\|_{L^\infty((u_\infty, Du_\infty)(\bar{\Omega}))} \right. \\ &\quad \left. + \|\partial_P g\|_{L^\infty((u_\infty, Du_\infty)(\bar{\Omega}))} \right), \end{aligned}$$

where  $C(\infty, \Omega) > 0$  is the maximum of the Poincaré and the Poincaré-Wirtinger inequality constants on  $\Omega$  for  $p = \infty$  (with the former being equal to  $\text{diam}(\Omega)$ ). As  $g \geq 0$  and  $\|g(u_\infty, Du_\infty)\|_{L^\infty(\Omega)} = 1$ , we conclude that  $0 \leq g(u_\infty, Du_\infty) \leq 1$  on  $\bar{\Omega}$ . Hence  $(u_\infty, Du_\infty)(\bar{\Omega}) \subseteq \{0 \leq g \leq 1\} = \{g \leq 1\}$ . Thus

$$1 \leq \|D^2 u_\infty\|_{L^\infty(\Omega)} \text{diam}(\Omega) \left( C(\infty, \Omega) \|\partial_\eta g\|_{L^\infty(\{g \leq 1\})} + \|\partial_P g\|_{L^\infty(\{g \leq 1\})} \right)$$

Rearranging assumption (1.3)(d), we may write  $|X| \leq C_4^{-\frac{1}{\alpha}} (f(X) + C_3)^{\frac{1}{\alpha}}$ , for any  $X \in \mathbb{R}_s^{N \times n^2}$ . Combining this inequality with the previous bound, we deduce

$$\begin{aligned} C_4^{\frac{1}{\alpha}} &\leq \left( \|f(D^2 u_\infty)\|_{L^\infty(\Omega)} + C_3 \right)^{\frac{1}{\alpha}} \text{diam}(\Omega) \left( C(\infty, \Omega) \|\partial_\eta g\|_{L^\infty(\{g \leq 1\})} \right. \\ &\quad \left. + \|\partial_P g\|_{L^\infty(\{g \leq 1\})} \right), \end{aligned}$$

which leads directly to the claimed lower bound for the eigenvalue.

Now we establish the upper bound for  $\Lambda_\infty$ . Since  $\Omega$  is by assumption a bounded domain with  $C^2$  boundary, by standard results (see e.g. [16, Sec. 14.6]), the distance function

$$d_\Omega \equiv \text{dist}(\cdot, \partial\Omega) : \mathbb{R}^n \rightarrow \mathbb{R},$$

which is in  $W_{\text{loc}}^{1,\infty}(\mathbb{R}^n)$ , is also  $C^2$  on an inner tubular neighbourhood of  $\partial\Omega$ , namely there exists  $\varepsilon_0 \in (0, 1)$  such that

$$d_\Omega \in C^2(\overline{\Omega^{\varepsilon_0}}), \quad \Omega^\varepsilon := \{d_\Omega < \varepsilon\} \cap \Omega.$$

Let us also for convenience symbolize  $\Omega_\varepsilon := \{d_\Omega > \varepsilon\} \cap \Omega$ . Let us also fix  $k \in \{1, 2\}$ , a unit vector  $e \in \mathbb{R}^N$  and  $\zeta \in C^2(\mathbb{R}^n)$  with  $\zeta \equiv 0$  on  $\Omega_{\varepsilon_0}$ . Then, for any  $t_0 > 0$ , the map  $\xi_0 := t_0(d_\Omega)^k \zeta e$  satisfies

$$\xi_0 \in C^2(\overline{\Omega}; \mathbb{R}^N).$$

Since  $d_\Omega = 0$  on  $\partial\Omega$  and also  $D(d_\Omega^2) = 0$  on  $\partial\Omega$ , it follows that  $\xi_0 \in W_H^{2,\infty}(\Omega; \mathbb{R}^N)$  if  $k = 1$ , whilst  $\xi_0 \in W_C^{2,\infty}(\Omega; \mathbb{R}^N)$  if  $k = 2$ . We will consider both cases simultaneously and declare this as

$$\xi_0 \in W_B^{2,\infty}(\Omega; \mathbb{R}^N).$$

By lemma 2.1, we can adjust the constant  $t_0 > 0$  to arrange

$$\|g(\xi_0, D\xi_0)\|_{L^\infty(\Omega)} = 1.$$

Hence,  $\xi_0$  is in the admissible class of the minimization problem (1.1). By minimality and assumption (1.3), we have the estimate

$$\Lambda_\infty \leq C_5(t_0)^\alpha \left( \|D^2(d_\Omega^k \zeta)\|_{L^\infty(\Omega^{\varepsilon_0})} \right)^\alpha + C_6. \quad (2.3)$$

By a direct computation, we have

$$\begin{cases} D^2(d_\Omega^k \zeta) = k[(k-1)Dd_\Omega \otimes Dd_\Omega + d_\Omega^{k-1} D^2 d_\Omega] \zeta + d_\Omega^k D^2 \zeta \\ \quad + kd_\Omega^{k-1} (Dd_\Omega \otimes D\zeta + D\zeta \otimes Dd_\Omega), \end{cases} \quad (2.4)$$

on  $\overline{\Omega}$ . For any  $x \in \overline{\Omega^{\varepsilon_0}}$ , let us set  $P_\Omega(x) := \text{Proj}_{\partial\Omega}(x)$ . Then, by [16, Sec. 14.6, L. 14.17], it follows that  $|x - P_\Omega(x)| = d_\Omega(x)$ , and we also have the next estimates

$$\begin{cases} \|d_\Omega\|_{L^\infty(\Omega^{\varepsilon_0})} \leq \varepsilon_0, \\ \|Dd_\Omega\|_{L^\infty(\Omega^{\varepsilon_0})} \leq 1, \\ \|D^2 d_\Omega\|_{L^\infty(\Omega^{\varepsilon_0})} \leq \sum_{i=1}^{n-1} \left\| \frac{\kappa_i \circ P_\Omega}{1 - (\kappa_i \circ P_\Omega) d_\Omega} \right\|_{L^\infty(\Omega^{\varepsilon_0})}, \end{cases} \quad (2.5)$$

where  $\{\kappa_1, \dots, \kappa_{n-1}\}$  are the principal curvatures of  $\partial\Omega$ . By (2.3)-(2.5) we have the estimate

$$\begin{aligned} \Lambda_\infty &\leq C_5(2t_0)^\alpha \left( \|D^2 \zeta\|_{L^\infty(\Omega)} + \|D\zeta\|_{L^\infty(\Omega)} \right. \\ &\quad \left. + \|\zeta\|_{L^\infty(\Omega)} \left( 1 + \sum_{i=1}^{n-1} \left\| \frac{\kappa_i \circ P_\Omega}{1 - (\kappa_i \circ P_\Omega) d_\Omega} \right\|_{L^\infty(\Omega^{\varepsilon_0})} \right)^\alpha \right) + C_6. \end{aligned} \quad (2.6)$$

It remains to select an appropriate function  $\zeta$  in order to estimate its derivatives in terms of the geometry of  $\Omega$ , and to obtain an estimate for  $t_0$ . For the former, we

argue as follows. Let  $(\eta^\delta)_{\delta>0}$  be the family of standard mollifying kernels, as e.g. in [27]. We select

$$\zeta := \eta^{\varepsilon_0} * (\chi_{\mathbb{R}^n \setminus \Omega}),$$

which is the regularization of the characteristic of the complement of  $\Omega$ . It follows that this function satisfies the initial requirements, and additionally

$$\begin{cases} D\zeta &= \eta^{\varepsilon_0} * (D\chi_{\mathbb{R}^n \setminus \Omega}) = \eta^{\varepsilon_0} * (\mathcal{H}^{n-1} \llcorner_{\partial\Omega} Dd_\Omega), \\ D^2\zeta &= D\eta^{\varepsilon_0} * (D\chi_{\mathbb{R}^n \setminus \Omega}) = \frac{1}{\varepsilon_0} (D\eta)^{\varepsilon_0} * (\mathcal{H}^{n-1} \llcorner_{\partial\Omega} Dd_\Omega), \end{cases}$$

by standard properties of convolutions and the differentiation of BV functions (see e.g. [15] and [14, Ch. 5, p. 198]). Then, by Young's inequality for convolutions, we have the estimates

$$\begin{cases} \|D^2\zeta\|_{L^\infty(\mathbb{R}^n)} &\leq \frac{C}{\varepsilon_0^{n+1}} \mathcal{H}^{n-1}(\partial\Omega), \\ \|D\zeta\|_{L^\infty(\mathbb{R}^n)} &\leq \mathcal{H}^{n-1}(\partial\Omega), \\ \|\zeta\|_{L^\infty(\mathbb{R}^n)} &\leq 1, \end{cases} \quad (2.7)$$

for some universal constant  $C > 0$ . Now we work towards an estimate for  $t_0$  appearing in (2.3). By assumption (1.4), we have that the sublevel sets  $\{g \leq t\}$  are compact in  $\mathbb{R}^N \times \mathbb{R}^{N \times n}$  for any  $t \geq 0$ . Let us define  $R(t)$  as the smallest radius of the  $N$ -dimensional ball, for which  $\{g \leq t\}$  is contained into the cylinder  $\bar{\mathbb{B}}_{R(t)}^N(0) \times \mathbb{R}^{N \times n}$ :

$$R(t) := \inf \left\{ R > 0 : \{g \leq t\} \subseteq \bar{\mathbb{B}}_R^N(0) \times \mathbb{R}^{N \times n} \right\}. \quad (2.8)$$

Then, we define a strictly increasing function  $\rho : [0, \infty) \rightarrow [0, \infty)$  by setting

$$\rho(t) := t + \sup_{0 \leq s \leq t} R(s). \quad (2.9)$$

Then,  $\rho$  satisfies  $\rho(0) = 0$ , and also that

$$\{g \leq t\} \subseteq \bar{\mathbb{B}}_{\rho(t)}^N(0) \times \mathbb{R}^{N \times n},$$

for any  $t \geq 0$ . Further, by construction,

$$\left\{ (\eta, P) \in \mathbb{R}^N \times \mathbb{R}^{N \times n} : \rho^{-1}(|\eta|) \leq t \right\} = \bar{\mathbb{B}}_{\rho(t)}^N(0) \times \mathbb{R}^{N \times n}.$$

The above imply

$$\rho^{-1}(|\eta|) \leq g(\eta, P), \quad (\eta, P) \in \mathbb{R}^N \times \mathbb{R}^{N \times n}.$$

Next, since  $d_\Omega^k \zeta$  vanishes on  $\partial\Omega \cup \overline{\Omega_{\varepsilon_0}}$  and  $g(0, 0) = 0$ , we have

$$\begin{aligned} 1 &= \|g(\xi_0, D\xi_0)\|_{L^\infty(\Omega)} \\ &= \sup_{\Omega^{\varepsilon_0}} g(\xi_0, D\xi_0) \\ &\geq \sup_{\Omega^{\varepsilon_0}} \rho^{-1}(|\xi_0|) \\ &\geq \sup_{\Omega^{\varepsilon_0}} \rho^{-1}(t_0 |d_\Omega^k \zeta|). \end{aligned}$$

Since  $d_\Omega \equiv \varepsilon_0/4$  on  $\partial\Omega^{\varepsilon_0/4}$ , and  $\rho^{-1}$  is strictly increasing, the above implies

$$\begin{aligned} 1 &\geq \max_{\partial\Omega^{\varepsilon_0/4}} \rho^{-1}(t_0 |d_\Omega^k \zeta|) \\ &= \max_{\partial\Omega^{\varepsilon_0/4}} \rho^{-1}\left(t_0 \left(\frac{\varepsilon_0}{4}\right)^k \zeta\right) \\ &= \rho^{-1}\left(t_0 \left(\frac{\varepsilon_0}{4}\right)^k \max_{\partial\Omega^{\varepsilon_0/4}} \zeta\right). \end{aligned} \tag{2.10}$$

Now we estimate  $\max_{\partial\Omega^{\varepsilon_0/4}} \zeta$  from below. Fix  $x \in \partial\Omega^{\varepsilon_0/4}$ . Then, since the standard mollifying kernel  $\eta$  is a radial function (see e.g. [27]), there exists a universal  $c > 0$  such that  $\eta \geq c$  on  $\mathbb{B}_{1/2}(0)$ . Therefore,

$$\begin{aligned} \zeta(x) &= \frac{1}{\varepsilon_0^n} \int_{\mathbb{B}_{\varepsilon_0}(x)} \chi_{\mathbb{R}^n \setminus \Omega} \eta\left(\frac{|y-x|}{\varepsilon_0}\right) dy \\ &\geq \frac{1}{\varepsilon_0^n} \int_{\mathbb{B}_{\varepsilon_0/2}(x) \setminus \Omega} \eta\left(\frac{|y-x|}{\varepsilon_0}\right) dy \\ &\geq \frac{c}{\varepsilon_0^n} \mathcal{L}^n(\mathbb{B}_{\varepsilon_0/2}(x) \setminus \Omega), \end{aligned}$$

for any  $x \in \partial\Omega^{\varepsilon_0/4}$ . Finally, since  $\partial\Omega$  satisfies the exterior sphere condition, the set  $\mathbb{B}_{\varepsilon_0/2}(x) \setminus \Omega$  contains a ball  $\mathbb{B}_{\bar{r}}(\bar{x})$  centred at some point  $\bar{x}$ , where the maximum possible radius  $\bar{r}$  is given by

$$\bar{r} = \min \left\{ \frac{\varepsilon_0}{8}, \min_{i=1, \dots, n-1} \frac{1}{\|\kappa_i\|_{C^0(\partial\Omega)}} \right\}.$$

Therefore, if  $\omega(n)$  is the volume of the unit ball in  $\mathbb{R}^n$ ,

$$\begin{aligned} \zeta(x) &\geq \frac{c}{\varepsilon_0^n} \mathcal{L}^n(\mathbb{B}_{\varepsilon_0/2}(x) \setminus \Omega) \\ &\geq \frac{c}{\varepsilon_0^n} \mathcal{L}^n(\mathbb{B}_{\bar{r}}(\bar{x})) \\ &= \frac{c}{\varepsilon_0^n} \omega(n) \bar{r}^n \end{aligned}$$

$$\begin{aligned}
&= \frac{c\omega(n)}{\varepsilon_0^n} \min \left\{ \left( \frac{\varepsilon_0}{8} \right)^n, \min_{i=1, \dots, n-1} \frac{1}{(\|\kappa_i\|_{C^0(\partial\Omega)})^n} \right\} \\
&= c\omega(n) \min \left\{ \frac{1}{2^{3n}}, \min_{i=1, \dots, n-1} \frac{1}{(\varepsilon_0 \|\kappa_i\|_{C^0(\partial\Omega)})^n} \right\},
\end{aligned}$$

for any  $x \in \partial\Omega^{\varepsilon_0/4}$ . Hence, we have established the lower bound

$$\max_{\partial\Omega^{\varepsilon_0/4}} \zeta \geq c\omega(n) \min \left\{ \frac{1}{2^{3n}}, \min_{i=1, \dots, n-1} \frac{1}{(\varepsilon_0 \|\kappa_i\|_{C^0(\partial\Omega)})^n} \right\}. \quad (2.11)$$

By (2.10) and (2.11), we infer (since  $\varepsilon_0 < 1$  and  $k \in \{1, 2\}$ ) that

$$\begin{aligned}
t_0 &\leq \frac{4^k \rho(1) \varepsilon_0^{n-k}}{c\omega(n)} \frac{1}{\min \left\{ \frac{1}{2^{3n}}, \min_{i=1, \dots, n-1} \frac{1}{(\varepsilon_0 \|\kappa_i\|_{C^0(\partial\Omega)})^n} \right\}} \\
&\leq \frac{32\rho(1)}{c\omega(n)} \left( 2^{3n} + \max_{i=1, \dots, n-1} (\|\kappa_i\|_{C^0(\partial\Omega)})^n \right).
\end{aligned} \quad (2.12)$$

By (2.6), (2.7), and (2.12), we conclude with the following upper bound for the eigenvalue:

$$\begin{aligned}
\Lambda_\infty &\leq C_6 + C_5 \left[ \frac{16\rho(1)}{c\omega(n)} \left( 2^{3n} + \max_{i=1, \dots, n-1} (\|\kappa_i\|_{C^0(\partial\Omega)})^n \right) \right]^\alpha \cdot \\
&\quad \cdot \left\{ 1 + \left( 1 + \frac{C}{\varepsilon_0^{n+1}} \right) \mathcal{H}^{n-1}(\partial\Omega) + \sum_{i=1}^{n-1} \left\| \frac{\kappa_i \circ P_\Omega}{1 - (\kappa_i \circ P_\Omega) d_\Omega} \right\|_{L^\infty(\Omega^{\varepsilon_0})} \right\}^\alpha. \quad (2.13)
\end{aligned}$$

The claimed estimate (1.8) follows from (2.13) above, by recalling that in view of (2.8)–(2.9), we have

$$\rho(1) = 1 + \sup_{0 \leq t \leq 1} R(t),$$

and also that the last term of (2.13) is finite at least when

$$\varepsilon_0 < \frac{1}{\max_{i=1, \dots, n-1} \|\kappa_i\|_{C^0(\partial\Omega)}}.$$

The result ensues.  $\square$

LEMMA 2.6. *For any  $p > (n/\alpha) + 2$ , there exist measures  $\nu_\infty \in \mathcal{M}(\overline{\Omega})$  and  $M_\infty \in \mathcal{M}(\overline{\Omega}; \mathbb{R}_s^{N \times n^2})$  such that, along perhaps a further sequence  $(p_j)_1^\infty$  of exponents, we have*

$$\begin{cases} \nu_p \xrightarrow{*} \nu_\infty, & \text{in } \mathcal{M}(\overline{\Omega}), \\ M_p \xrightarrow{*} M_\infty, & \text{in } \mathcal{M}(\overline{\Omega}; \mathbb{R}_s^{N \times n^2}), \end{cases}$$

as  $j \rightarrow \infty$ , where the approximating measures  $\nu_p, M_p$  are given by (1.12).

*Proof of lemma 2.6.* We begin by noting that since  $g \geq 0$  and  $\|g(u_p, Du_p)\|_{L^p(\Omega)} = 1$ , in view of (1.12) we have the bound

$$\|\nu_p\|(\bar{\Omega}) = \nu_p(\bar{\Omega}) = \int_{\Omega} g(u_p, Du_p)^{p-1} d\mathcal{L}^n \leq \left( \int_{\Omega} g(u_p, Du_p)^p d\mathcal{L}^n \right)^{\frac{p-1}{p}} = 1.$$

By the sequential weak\* compactness of the space of Radon measures we can conclude that  $\nu_p \xrightarrow{*} \nu_{\infty}$ , in  $\mathcal{M}(\bar{\Omega})$  up to the passage to a further subsequence. Now we establish appropriate total variation bounds for the measure  $M_p$ . Since  $f \geq 0$ , by the bounds of lemma 2.4 and assumption (1.3), we estimate (for sufficiently large  $p$ )

$$\begin{aligned} \|M_p\|(\bar{\Omega}) &= \int_{\Omega} \left( \frac{f(D^2 u_p)}{\Lambda_p} \right)^{p-1} |\partial f(D^2 u_p)| d\mathcal{L}^n \\ &\leq \frac{1}{\Lambda_p^{p-1}} \int_{\Omega} f(D^2 u_p)^{p-1} (C_5 f(D^2 u_p)^{\beta} + C_6) d\mathcal{L}^n \\ &= \frac{C_5}{\Lambda_p^{p-1}} \int_{\Omega} f(D^2 u_p)^{p-1+\beta} d\mathcal{L}^n + \frac{C_6}{\Lambda_p^{p-1}} \int_{\Omega} f(D^2 u_p)^{p-1} d\mathcal{L}^n. \end{aligned}$$

Hence,

$$\begin{aligned} \|M_p\|(\bar{\Omega}) &\leq \frac{C_5}{\Lambda_p^{p-1}} \left( \int_{\Omega} f(D^2 u_p)^p d\mathcal{L}^n \right)^{\frac{p-1+\beta}{p}} + \frac{C_6}{\Lambda_p^{p-1}} \left( \int_{\Omega} f(D^2 u_p)^p d\mathcal{L}^n \right)^{\frac{p-1}{p}} \\ &= C_5 \frac{(L_p)^{p-1+\beta}}{\Lambda_p^{p-1}} + C_6 \frac{(L_p)^{p-1}}{\Lambda_p^{p-1}} \\ &= \left( \frac{L_p}{\Lambda_p} \right)^{p-1} (C_5 L_p^{\beta} + C_6) \\ &\leq \left( \frac{C_8}{C_1} \right)^{1-\frac{1}{p}} (C_5 (\Lambda_{\infty} + 1)^{\beta} + C_6). \end{aligned}$$

The above bound allows to conclude that  $M_p \xrightarrow{*} M_{\infty}$  in  $\mathcal{M}(\bar{\Omega}; \mathbb{R}_s^{N \times n^2})$ , along perhaps a further subsequence of indices  $(p_j)_1^{\infty}$  as  $j \rightarrow \infty$ .  $\square$

To conclude the proof of theorem 1.1 we must ensure the PDE system (1.5) is indeed satisfied by the quadruple  $(u_{\infty}, \Lambda_{\infty}, M_{\infty}, \nu_{\infty})$ .

LEMMA 2.7. *If  $M_{\infty} \in \mathcal{M}(\bar{\Omega}; \mathbb{R}_s^{N \times n^2})$  and  $\nu_{\infty} \in \mathcal{M}(\bar{\Omega})$  are the measures obtained in lemma 2.6, then the pair  $(u_{\infty}, \Lambda_{\infty})$  satisfies (1.5) for all  $\phi \in C_B^2(\bar{\Omega}; \mathbb{R}^N)$ .*

*Proof of lemma 2.7.* Fix a test function  $\phi \in C_B^2(\bar{\Omega}; \mathbb{R}^N)$  and  $p > n/\alpha + 2$  by (1.12) we may rewrite the PDE system in (1.11) as follows

$$\int_{\Omega} D^2 \phi : dM_p = \Lambda_p \int_{\Omega} (\partial_{\eta} g(u_p, Du_p) \cdot \phi + \partial_P g(u_p, Du_p) : D\phi) d\nu_p,$$

Recall that, by proposition 2.5, we have  $\Lambda_p \rightarrow \Lambda_{\infty}$  and also  $(u_p, Du_p) \rightarrow (u_{\infty}, Du_{\infty})$  uniformly on  $\bar{\Omega}$ , as  $p_j \rightarrow \infty$ . By assumption (1.4)(a), we have that

$\partial_\eta g(u_p, Du_p) \rightarrow \partial_\eta g(u_\infty, Du_\infty)$  and also  $\partial_P g(u_p, Du_p) \rightarrow \partial_P g(u_\infty, Du_\infty)$ , both uniformly on  $\bar{\Omega}$ , as  $p_j \rightarrow \infty$ . The result ensues by invoking lemma 2.6, in conjunction with weak\*-strong continuity of the duality pairing  $\mathcal{M}(\bar{\Omega}) \times C(\bar{\Omega}) \rightarrow \mathbb{R}$ .  $\square$

### Acknowledgements

The authors would like to thank the referee of this paper for their careful reading of the manuscript, as well as their constructive suggestions.

E.C. has been financially supported through the UK EPSRC scholarship GS19-055

### References

- 1 An Introduction to Viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in  $L^\infty$ , (Springer Briefs in Mathematics, 2015). DOI: 10.1007/978-3-319-12829-0.
- 2 N. Ansini and F. Prinari. On the lower semicontinuity of supremal functional under differential constraints. *ESAIM - Control, Opt. Calc. Var.* **21** (2015), 1053–1075.
- 3 G. Aronsson and E. N. Barron.  $L^\infty$  variational problems with running costs and constraints. *Appl. Math. Optim.* **65** (2012), 53–90.
- 4 E. N. Barron and R. Jensen. Minimizing the  $L^\infty$  norm of the gradient with an energy constraint. *Comm. Partial Differ. Equ.* **30** (2005), 1741–1772.
- 5 E. N. Barron, R. Jensen and C. Wang. Lower semicontinuity of  $L^\infty$  functionals. *Ann. I. H. Poincaré AN.* **18** (2001), 495–517.
- 6 L. Bungert and Y. Korolev. Eigenvalue problems in  $L^\infty$ : optimality conditions, duality, and relations with optimal transport. *Commun. Am. Math. Soc.* **2** (2022), 345.
- 7 F. Cagnetti.  $k$ -quasi-convexity reduces to quasi-convexity. *Proc. Roy. Soc. Edinburgh Sect. A.* **141** (2011), 673–708.
- 8 T. Champion, L. De Pascale and F. Prinari.  $\Gamma$ -convergence and absolute minimizers for supremal functionals. *COCV ESAIM: Control, Optim. Calc. Var.* **10** (2004), 14–27.
- 9 E. Clark, N. Katzourakis and B. Muha. Data assimilation for the Navier-Stokes equations through PDE-constrained optimisation in  $L^\infty$ . *Nonlinearity* **35** (2021), 470.
- 10 E. Clark and N. Katzourakis, On isosupremic vectorial minimisation problems in  $L^\infty$  with general nonlinear constraints. *Adv. Calc. Var.* (in press).
- 11 B. Dacorogna. *Direct methods in the calculus of variations* (2nd ed., Vol. 78, Springer: Applied Mathematical Sciences, 2008).
- 12 G. Dal Maso, I. Fonseca, G. Leoni and M. Morini. Higher-order quasiconvexity reduces to quasiconvexity. *Arch. Ration. Mech. Anal.* **171** (2004), 55–81.
- 13 J. M. Danskin. The theory of min-max with application. *SIAM J. Appl. Math.* **14** (1966), 641–664.
- 14 L. C. Evans and R. Gariepy. *Measure theory and fine properties of functions* (CRC press: Studies in advanced mathematics, 1992).
- 15 G. B. Folland, *Real Analysis: Modern Techniques and Their Applications*, Pure and Applied Mathematics: A Wiley Series of Texts, Vol. 40 (Monographs and Tracts, 1999).
- 16 D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order* (2nd ed., Springer-Verlag GmbH Germany, part of Springer Nature, 2001).
- 17 J. Hutchinson. Second fundamental form for varifolds and the existence of surfaces minimising curvature. *Indiana J. Math.* **35** (1986), 45–71.
- 18 P. Juutinen and P. Lindqvist. On the higher eigenvalues for the  $\infty$ -eigenvalue problem. *Calc. Var.* **23** (2005), 169–192.
- 19 P. Juutinen, P. Lindqvist and J. Manfredi. The  $\infty$ -eigenvalue problem. *Arch. Ration. Mech. Anal.* **148** (1999), 89–105.
- 20 N. Katzourakis. An  $L^\infty$  regularisation strategy to the inverse source identification problem for elliptic equations. *SIAM J. Math. Anal.* **51** (2019), 1349–1370.

21 N. Katzourakis. Inverse optical tomography through PDE-constrained optimisation in  $L^\infty$ .  
*SIAM J. Control Optim.* **57** (2019), 4205–4233.

22 N. Katzourakis. A minimisation problem in  $L^\infty$  with PDE and unilateral constraints.  
*ESAIM: Control, Optim. Calc. Var.* **26** (2020), 60.

23 N. Katzourakis. Generalised vectorial  $\infty$ -eigenvalue nonlinear problems for  $L^\infty$  functionals.  
*Nonlinear Anal.* **219** (2022), 112806. <https://doi.org/10.1016/j.na.2022.112806>. (online).

24 N. Katzourakis and R. Moser, *Existence, uniqueness and structure of second order absolute minimisers*, Arch. Ration. Mech. Anal., published online 06/09/2018. 10.1007/s00205-018-1305-6.

25 N. Katzourakis and E. Parini. The eigenvalue problem for the  $\infty$ -Bilaplacian. *Nonlinear Differ. Equa. Appl. NoDEA* **24** (2017), 1–25.

26 N. Katzourakis and T. Pryer. 2nd order  $L^\infty$  variational problems and the  $\infty$ -polyLaplacian.  
*Adv. Calc. Var.* **13** (2020), 115–140.

27 N. Katzourakis and E. Varvaruca, *An illustrative introduction to modern analysis* (CRC Press / Taylor & Francis, 2018), p. 560.

28 C. Kreisbeck and E. Zappale. Lower semicontinuity and relaxation of nonlocal  $L^\infty$ -functionals. *Calc. Var. PDE* **59** (2020), 1–36.

29 Q. Miao, C. Wang and Y. Zhou. Uniqueness of absolute minimizers for  $L^\infty$ -functionals involving Hamiltonians  $H(x, p)$ . *Arch. Ration. Mech. Anal.* **223** (2017), 141–198.

30 F. Prinari and E. Zappale. A relaxation result in the vectorial setting and power law approximation for supremal functionals. *J. Optim. Theory Appl.* **186** (2020), 412–452.

31 A. N. Ribeiro and E. Zappale. Existence of minimisers for nonlevel convex functionals.  
*SIAM J. Control Opt.* **52** (2014), 3341–3370.

32 E. Zeidler. *Nonlinear functional analysis and its application III: variational methods and optimization* (Berlin: Springer-Verlag, 1985).

33 X. P. Zhou. Weak lower semicontinuity of a functional with any order. *J. Math. Anal. Appl.* **221** (1998), 217–237.