

Maize monoculture supported pre-Columbian urbanism in southwestern Amazonia

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Lombardo, U., Hilbert, L., Bentley, M., Ramsey, C. B., Dudgeon, K., Gaitan-Roca, A., Iriarte, J., Mejía Ramón, A. G., Quezada, S., Raczka, M. ORCID: <https://orcid.org/0000-0002-6602-8087>, Watling, J. G., Neves, E. and Mayle, F. ORCID: <https://orcid.org/0000-0001-9208-0519> (2025) Maize monoculture supported pre-Columbian urbanism in southwestern Amazonia. *Nature*, 639. pp. 119-123. ISSN 1476-4687 doi: 10.1038/s41586-024-08473-y Available at <https://centaur.reading.ac.uk/119851/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/s41586-024-08473-y>

Publisher: Nature Publishing

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Maize monoculture supported pre-Columbian urbanism in southwestern Amazonia

<https://doi.org/10.1038/s41586-024-08473-y>

Received: 26 April 2024

Accepted: 2 December 2024

Published online: 29 January 2025

Open access

 Check for updates

Umberto Lombardo^{1,2}✉, Lautaro Hilbert^{1,3}, McKenzie Bentley^{4,5}, Christopher Bronk Ramsey⁴, Kate Dudgeon¹, Albert Gaitan-Roca², José Iriarte⁶, Andrés G. Mejía Ramón¹, Sergio Quezada⁷, Marco Raczká^{8,9}, Jennifer G. Watling³, Eduardo Neves³ & Francis Mayle⁹

The Casarabe culture (500–1400 CE), spreading over roughly 4,500 km² of the monumental mounds region of the Llanos de Moxos, Bolivia, is one of the clearest examples of urbanism in pre-Columbian (pre-1492 CE) Amazonia. It exhibits a four-tier hierarchical settlement pattern, with hundreds of monumental mounds interconnected by canals and causeways^{1,2}. Despite archaeological evidence indicating that maize was cultivated by this society³, it is unknown whether it was the staple crop and which type of agricultural farming system was used to support this urban-scale society. Here, we address this issue by integration of remote sensing, field survey and microbotanical analyses, which shows that the Casarabe culture invested heavily in landscape engineering, constructing a complex system of drainage canals (to drain excess water during the rainy season) and newly documented savannah farm ponds (to retain water in the dry season). Phytolith analyses of 178 samples from 18 soil profiles in drained fields, farm ponds and forested settings record the singular and ubiquitous presence of maize (*Zea mays*) in pre-Columbian fields and farm ponds, and an absence of evidence for agricultural practices in the forest. Collectively, our findings show how the Casarabe culture managed the savannah landscape for intensive year-round maize monoculture that probably sustained its relatively large population. Our results have implications for how we conceive agricultural systems in Amazonia, and show an example of a Neolithic-like, grain-based agrarian economy in the Amazon.

The role of grain agriculture as the subsistence base of prehistoric complex societies in both the Old and New World has been a matter of sustained debate for many decades (see, for example, refs. 4–8). In Mesoamerica, the earliest evidence of maize as a staple crop dates to 4,000 calendar years before the present⁹. The timing and nature of maize's role as the staple crop of Andean civilizations, as seen in early historical accounts, is controversial (see, for example, refs. 6,10). In Amazonia it is well established, from both archaeological and palaeoecological data, that maize has been cultivated since at least 6,850 calendar years before the present¹¹; however, to date there is no evidence of it being a staple crop. Most societies had mixed economies relying on multiple cultigens^{12–16}. Roosevelt¹⁷ proposes that the rise of social complexity in the Amazon was based on maize agriculture. However, current archaeological evidence has not been conclusive of maize cultivation being the staple crop of complex societies of the Amazon¹⁵. Current archaeobotanical and palaeoecological data from Late Holocene complex societies in Amazonia indicate polyculture (mixed-cropping) agroforestry, not maize monoculture, as the basis of a subsistence economy^{15,18–22}.

Recent archaeological research has revealed evidence for low-density urbanism, social complexity and large populations in the Andean foothills of the Upano River region of Ecuador²³, and in the monumental

mounds region (MMR) in the seasonally flooded savannahs of the Bolivian Amazon¹. Here in the MMR, the Casarabe people built hundreds of monumental mounds interconnected by canals and causeways across a flat forest–savannah mosaic landscape dominated by seasonally flooded savannahs, with forests restricted to non-flooded palaeo-river levées. Whereas drained fields and terraces, built on extremely fertile volcanic soils, were clearly integral to low-density agrarian urbanism of the Upano region²³, the type of farming system needed to sustain the Casarabe culture is still unknown. It has been proposed that the construction of drainage canals permitted cultivation of the relatively fertile sediments of the seasonally flooded savannahs of the MMR²⁴ without the need for deforestation²⁵. However, no agricultural fields or other food production systems have hitherto been found in connection with such canals, leaving unanswered the question of how the Casarabe people managed to feed its relatively large population. To address this issue, we combine remote-sensing imagery with a programme of coring, test pits, radiocarbon dating and pollen and phytolith analyses on both seasonally flooded savannahs and forest.

We have identified two unreported and complementary agrotechnologies in the savannahs of the MMR: dense drainage networks and artificial farm ponds (Fig. 1 and Extended Data Fig. 1), in which different portions of a savannah (Fig. 1, top left inset), or different savannahs within the same

¹Institute of Environmental Science and Technology, Universitat Autònoma de Barcelona, Barcelona, Spain. ²Department of Prehistory, Universitat Autònoma de Barcelona, Barcelona, Spain.

³Museum of Archaeology and Ethnology, University of São Paulo, São Paulo, Brazil. ⁴School of Archaeology, University of Oxford, Oxford, UK. ⁵School of Ocean and Earth Science, University of Southampton, Southampton, UK. ⁶Department of Archaeology and History, University of Exeter, Exeter, UK. ⁷Trinidad, Bolivia. ⁸School of Geography, University of Nottingham, Nottingham, UK.

⁹Department of Geography and Environmental Science, University of Reading, Reading, UK. [✉]e-mail: Umberto.Lombardo@uab.cat




Fig. 1 | Two examples of engineered landscape in the MMR. Triangles represent monumental mounds; black dots, ponds; thin black lines, canals; light grey areas, forest; white areas, savannah; and dark grey areas, lakes. Inset top left, the northern side of a savannah is crisscrossed by canals but the southern side is dotted with ponds. Inset bottom right, one savannah (to the east) has been modified by the excavation of a densely packed network of

drainage canals, and two other savannahs, to the west, are dotted with ponds; in the former, the network of canals drains into Lake Francia located about 4 km to the north. 690 and 695 indicate the locations of the two farm ponds sampled for this study. Inset top right, boundaries of Amazonia as defined in ref. 43 and the image of World Countries Generalized provided by ESRI under the ArcGIS Pro licence. Scale bar, 5 km. Credit: European Commission JRC.

area (Fig. 1, bottom right inset), have been heavily modified—into either intricate arrangements of canals or clusters of circular depressions.

The drainage network

In one of the savannahs under study (Fig. 2), the small canals converge into larger canals that drain the whole savannah toward Lake Francia to the north (Fig. 1b). We identified three orders of drainage canals: the first order (1 in Fig. 2b), the smallest, are around 4 m wide and 25 cm deep, the second order (2 in Fig. 2b) are around 8 m wide and 70 cm deep and the main canal (third order) that drains into the lake is 14 m wide and 1.8 m deep (3 in Fig. 2b), becoming 3.2 m deep about 1.5 km before reaching the lake. Overall, the drainage network drains towards the north, becoming ever deeper with respect to the general topography. Several stratigraphic profiles of the canals show that the original depth of the canal network was around 80 cm deeper than at present for the second-order canals (see profiles 667 and 671 in Extended Data Fig. 2) and around 45 cm deeper for the first-order canals (for example, profile 674 in Extended Data Fig. 2). The drainage network is associated with circular elevated platforms roughly 50 cm in height, resembling pre-Columbian forest islands¹¹, and with small mounds of around 2–3 m in diameter. The elevated platforms are surrounded by deep canals (profiles 666 and 677 in Extended Data Fig. 2).

Soil cores were collected from several locations both inside the canals and between them (Fig. 2b). Phytolith analysis shows a high abundance of phytoliths derived from the cob glumes and leaves of *Zea mays* in almost all canal soil profiles (Extended Data Fig. 2), with sporadic presence of *Cucurbita* spp. (666 and 677), *Manihot* sp. (677), *Calathea* sp. (674) and *Lagenaria* sp. (667) phytoliths. We cannot exclude the possibility that *Cucurbita* was cultivated in greater amounts than implied by the phytolith assemblage, because some domesticated *Cucurbita* varieties may lack scalloped phytoliths²⁶.

The majority of *Oryza* phytoliths are concentrated in the upper levels of all the profiles. Most upper-level glume phytoliths were from domesticated plants (Methods), whereas those from lower levels in the soil profiles (30–50 cm depth) were classified as wild species. These results are not surprising, because some of these fields are currently being used to grow modern Asian rice²⁷. The low production of diagnostic wavy-top rondel phytoliths in maize²⁸ and the high abundance of these, relative to the sporadic presence of phytoliths of other cultivars, indicate that maize was by far the principal cultivar in these savannahs. We attribute the absence of maize phytoliths in the uppermost 20–25 cm of the canal soil profiles to sedimentary fill from adjacent fields over recent decades or centuries—an inference corroborated by our soil phytolith data from fields between the canals (Extended Data Fig. 3). Here, maize phytoliths appear in only one of the three profiles, suggesting that the cultivated area is likely to have been established along the canals, probably on elevated rims that have since eroded into the canals. It is probable that, while in use, the original depth of the canals was maintained by their periodic re-excavation and redistribution of canal sediment fill along the canal margins, where maize was then planted (Extended Data Fig. 4), mimicking what has been proposed for raised-field agriculture in other regions of the Llanos de Moxos²⁹.

The forest

Forest in the study region grows on elevated surfaces, mostly fluvial levees, that remain above the water level during the rainy season. Four soil profiles were dug and sampled across the forest, along a transect from the savannah to a large 15-ha monumental mound (Fig. 2a and Extended Data Fig. 3), to reveal to what extent the forest was cleared for agriculture. No charcoal or any other evidence of fire was visible in any of the profiles. Phytolith profiles are all similar and do not show any obvious stratigraphic change, apart from a slight reduction in Areceaceae (palms)




Fig. 2 | Drainage canals and digital elevation model. **a**, The complete drainage network. Numbered dots indicate the location of phytolith profiles. **b**, Detail of the digital elevation model of the northern part of the drainage network. Dashed lines represent the forest–savannah boundary, showing that a great deal of the drainage network is presently covered by forest. See Fig. 1b for location. Scale bars, 1 km (a), 500m (b).

and an increase in Poaceae (grass) phytoliths, which could indicate a slight opening of the forest canopy. No cultivar phytoliths were found in any of the profiles (Extended Data Fig. 3). Although our data cannot show the extent to which the forest was used for agroforestry, wood harvesting, hunting or cultivation of medicinal plants, the absence of charcoal does show that slash-and-burn agriculture did not take place here.

The clusters of farm ponds

A large portion of the savannahs in the MMR contains clusters of circular depressions of 10–100 m in diameter. They are often connected, either by canals or directly by adjacency. These ponds are similar to

natural depressions called *gilgais*, an Aboriginal Australian name for water holes, that form in vertisols because of repeated expansion and contractions of the clay³⁰. To understand their genesis and use, we sampled two ponds in two different savannahs. Profiles 690 (Fig. 3) and 689 (Extended Data Fig. 3) were excavated and cored, respectively, in a large pond of roughly 100 m in diameter, with its central depression about 60 cm below the surroundings. Profile 695 (Extended Data Fig. 3) was excavated in a pond of roughly 30 m in diameter, and with a central depression currently 40 cm below the surrounding savannah. Pond profile 690 exhibits a very irregular, sharp contact between the organic sediment fill and the grey, inorganic clay below (Extended Data Fig. 5), and shows no evidence of shear surfaces (slickensides), suggesting that the pond was excavated and is not a *gilgai*. The anthropogenic origin of these depressions is further supported by their size, which is far larger than the 15–20-m-diameter *gilgais*^{31,32}, and by their clustered linear distribution (Extended Data Fig. 6). Sediment profiles from both ponds show the continuous presence of maize phytoliths and pollen (Fig. 3 and Extended Data Fig. 7) throughout, with phytoliths of *Cucurbita* sp. present in only two adjacent samples in profile 690 at around 40 cm depth, and a pollen grain of *Manihot* at approximately 50 cm depth (Extended Data Figs. 3, 7 and 8). No other cultivars were detected. The chronology of pond profile 690 indicates that this system was in use around 1250–1550 calendar years CE (Extended Data Fig. 5).

Today, the majority of these ponds hold water for most of the year, maintaining wet soil until the very end of the dry season. If this hydrological balance is representative of the past, these ponds would have provided sufficient water for maize cultivation around their margins throughout the dry season. This is not dissimilar to the *k' hochas* in the Bolivian altiplano³³, where yields are up to four times higher than for regional rainfed production. Similar pond-based farming systems have been described in Bangladesh and India^{34,35}, where ponds provide an integrated production system that includes fish farming, poultry and cultivation of pond dykes. A very similar system could have been in place in the MMR, where swamp eels (*Synbranchus marmoratus*) were an important part of the diet of the Casarabe culture³⁶; the Muscovy duck (*Cairina moschata*), the only known domesticated animal in the Amazon³⁷, was probably kept^{38,39}. Bones of *S. marmoratus* were found in pond 695 at a depth of 70 cm (Extended Data Fig. 9). During the dry season, these farm ponds would have served as ‘watering holes’, attracting game.

A pre-Columbian green revolution

The combination of these two types of landscape engineering—drainage canals and farm ponds—is unique to the MMR. We argue that it was a highly innovative agricultural strategy that enabled the Casarabe culture to substantially increase the cultivation period for maize, as well as providing easy access to fish, birds and game. Through the sophisticated system of drainage canals, some savannah wetlands were converted into drained fields suitable for intensive maize monoculture in the wet season, whereas the construction of clusters of farm ponds in other savannahs provided a reservoir of water that allowed pot irrigation, which enabled the continuation of maize agriculture throughout the dry season. The combination of these two water management systems would have allowed at least two harvests of maize per year. The lack of any evidence of cultivation and fire in the nearby forested areas suggests that slash-and-burn agriculture was unlikely to have been practised. Instead, this pre-Columbian Casarabe culture probably preserved the spatially limited, and hence highly valuable, forest resource for other key ecosystem services, such as firewood, building materials, medicinal plants and probably polyculture agroforestry. These data are corroborated by palaeoecological studies in the MMR that show no substantial change in forest cover²⁵ during the Casarabe culture period. There are at least seven monumental mounds surrounding the drainage network and five surrounding the savannah




Fig. 3 | Relative frequencies of phytoliths recovered from farm pond, profile 690. Horizontal bars represent percentages, circles correspond to the presence of plant taxa lower than 1% in abundance. Vertical axis is the depth of

the sample in cm. Source data are provided in Supplementary Data 1. non-diagn., non-diagnostic.

with the pond cluster, which includes profile 690. These form part of the four-tier settlement pattern belonging to a political structure that guaranteed food production and agricultural infrastructure maintenance for hundreds of years^{1,2}. Even though micro- and macrobotanical remains from monumental mounds show the presence of a variety of food and industrial crops, including maize, manioc, lerén, squash, peanuts, cotton, yams and palms^{40,41}, our data suggest that maize was the staple crop for the Casarabe culture when the drainage and pond agricultural system was in operation. Our data show that the absence of other cultigen pollen in MMR lake cores²⁵, and the greater abundance of maize macroremains, phytoliths and starch grains in the sediments and ceramics of Mendoza and Salvatierra monumental mounds^{40,41}, is not due to low pollen productivity/preservation or taphonomic bias, but instead reflects a real phenomenon of greater reliance on maize in the diet compared with other cultigens.

Our results overturn the assumption that the seasonally flooded savannah of southwestern Amazonia is suitable only for cattle ranching and intensive Asian rice agriculture, and unsuitable for nutrient-demanding crops such as maize. These findings have implications for our understanding of pre-Columbian subsistence economies across Amazonia and beyond. They indicate that, during the late Holocene, alongside intensive polyculture (mixed-cropping) agroforestry on Amazonian Dark Earths¹⁵, other agricultural systems such as drained fields and farm ponds in the Llanos de Moxos were primarily focused on the cultivation of maize. These practices bear similarities to agricultural strategies observed in later Andean states and chiefdoms⁴². Collectively, as long argued, intensive cultivation of maize has had a major role in supporting the economy of some of the most complex societies in the Americas. The Casarabe people demonstrated the ability to establish a highly intensive monoculture farming system on the savannahs based on maize, maintaining the surrounding forest cover and supporting one of the most complex pre-Columbian societies in lowland South America. The Casarabe culture of the MMR provides a clear example of when the rise of social complexity is linked to intensive food production and, more specifically, to maize monoculture. It also confirms the role of grain agriculture as the main driver for increasing social complexity and, probably, inequality⁷.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information,

acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-024-08473-y>.

1. Prümmer, H., Betancourt, C. J., Iriarte, J., Robinson, M. & Schaich, M. Lidar reveals pre-Hispanic low-density urbanism in the Bolivian Amazon. *Nature* **606**, 325–328 (2022).
2. Lombardo, U. & Prümmer, H. Pre-Columbian human occupation patterns in the eastern plains of the Llanos de Moxos, Bolivian Amazonia. *J. Archaeol. Sci.* **37**, 1875–1885 (2010).
3. Hermenegildo, T. *Fields and Forests: A Stable Isotope Perspective on the Subsistence Strategies of Past Amazonian Peoples*. Doctoral dissertation thesis, Univ. of Cambridge (2022).
4. d'Alpoim Guedes, J. An archaeobotanical perspective on the relationship between grain crops, non-grain crops and states. *Camb. Archaeol. J.* **29**, 694–696 (2019).
5. Lathrap, D. W. *The Upper Amazon*, Vol. 5 (Thames and Hudson, 1970).
6. Piperno, D. R. & Pearsall, D. M. *The Origins of Agriculture in the Lowland Neotropics* (Academic Press, 1998).
7. Scott, J. C. *Against the Grain: A Deep History of the Earliest States* (Yale Univ. Press, 2017).
8. Staller, J. E., Tykot, R. H. & Benz, B. F. *Histories of Maize in Mesoamerica* (Routledge, 2010).
9. Kennett, D. J. et al. Early isotopic evidence for maize as a staple grain in the Americas. *Sci. Adv.* **6**, eaba3245 (2020).
10. Burger, R. L. & Van Der Merwe, N. J. Maize and the origin of highland Chavín civilization: an isotopic perspective. *Am. Anthropol.* **92**, 85–95 (1990).
11. Lombardo, U. et al. Early Holocene crop cultivation and landscape modification in Amazonia. *Nature* **581**, 190–193 (2020).
12. Aceituno, F. J. & Loaiza, N. The origins and early development of plant food production and farming in Colombian tropical forests. *J. Anthropol. Archaeol.* **49**, 161–172 (2018).
13. Morell-Hart, S., Dussol, L. & Fedick, S. L. Agriculture in the Ancient Maya Lowlands (Part 1): paleoethnobotanical residues and new perspectives on plant management. *J. Archaeol. Res.* **31**, 561–615 (2023).
14. Fedick, S. L., Morell-Hart, S. & Dussol, L. Agriculture in the Ancient Maya Lowlands (Part 2): landesque capital and long-term resource management strategies. *J. Archaeol. Res.* **32**, 103–154 (2024).
15. Iriarte, J. et al. The origins of Amazonian landscapes: plant cultivation, domestication and the spread of food production in tropical South America. *Quat. Sci. Rev.* **248**, 106582 (2020).
16. Watling, J. et al. Direct archaeological evidence for Southwestern Amazonia as an early plant domestication and food production centre. *PLoS ONE* **13**, e0199868 (2018).
17. Roosevelt, A. C. *Parman: Prehistoric Maize and Manioc Subsistence Along the Amazon and Orinoco* (Academic Press, 1987).
18. Fausto, C. & Neves, E. G. Was there ever a Neolithic in the Neotropics? Plant familiarisation and biodiversity in the Amazon. *Antiquity* **92**, 1604–1618 (2018).
19. Neves, E. G. In *Human-Environment Interactions. Current and Future Directions* (eds Brondizio, E. S. & Moran, E. F.) 371–388 (Springer, 2013).
20. Schaan, D. In *Handbook of South American Archaeology* (eds Silverman, H. & Isbell, W. H.) 339–357 (Springer, 2008).
21. Maezumi, S. Y. et al. The legacy of 4,500 years of polyculture agroforestry in the eastern Amazon. *Nature Plants* **4**, 540–547 (2018).
22. Watling, J. et al. Impact of pre-Columbian “geoglyph” builders on Amazonian forests. *Proc. Natl Acad. Sci. USA* **114**, 1868 (2017).
23. Rostain, S. et al. Two thousand years of garden urbanism in the Upper Amazon. *Science* **383**, 183–189 (2024).

24. Lombardo, U., May, J.-H. & Veit, H. Mid- to late-Holocene fluvial activity behind pre-Columbian social complexity in the southwestern Amazon basin. *The Holocene* **22**, 1035–1045 (2012).

25. Whitney, B. S., Dickau, R., Mayle, F. E., Soto, J. D. & Iriarte, J. Pre-Columbian landscape impact and agriculture in the Monumental Mound region of the Llanos de Moxos, lowland Bolivia. *Quat. Res.* **80**, 207–217 (2013).

26. Piperno, D. R., Holst, I., Wessel-Beaver, L. & Andres, T. C. Evidence for the control of phytolith formation in *Cucurbita* fruits by the hard rind (Hr) genetic locus: archaeological and ecological implications. *Proc. Natl Acad. Sci. USA* **99**, 10923 (2002).

27. Lombardo, U. Pre-Columbian legacy and modern land use in the Bolivian Amazon. *PAGES* **31**, 16–17 (2023).

28. Pearsall, D. M., Chandler-Ezell, K. & Chandler-Ezell, A. Identifying maize in neotropical sediments and soils using cob phytoliths. *J. Archaeol. Sci.* **30**, 611–627 (2003).

29. Rodrigues, L. et al. An insight into pre-Columbian raised fields: the case of San Borja, Bolivian lowlands. *SOIL* **2**, 367–389 (2016).

30. Knight, M. J. Structural analysis and mechanical origins of gilgai at Boorook, Victoria, Australia. *Geoderma* **23**, 245–283 (1980).

31. Florinsky, I. V. & Arlashina, H. A. Quantitative topographic analysis of gilgai soil morphology. *Geoderma* **82**, 359–380 (1998).

32. Kishné, A. S., Morgan, C. L. S. & Neely, H. L. How much surface water can gilgai microtopography capture? *J. Hydrol.* **513**, 256–261 (2014).

33. Verweij, M. Towards sustainable pond farming. *LEISA Mag.* **17**, 43–45 (2001).

34. Das, A. et al. Livelihood security of small holder farmers in eastern Himalayas, India: pond based integrated farming system a sustainable approach. *Curr. Res. Environ. Sustain.* **3**, 100076 (2021).

35. Alam, M. R., Ali, M. A., Hossain, M. A., Molla, M. & Islam, F. J. B. J. O. A. R. Integrated approach of pond based farming systems for sustainable production and income generation. *Banglad. J. Agric. Res.* **34**, 577–584 (2009).

36. Prestes-Carneiro, G., Takayuki, Y., Jean-Louis, D., Kélig, M. & Philippe, B. Reconstructing freshwater fishing seasonality in a neotropical savanna: first application of swamp eel (*Synbranchus marmoratus*) sclerochronology to a pre-Columbian Amazonian site (Loma Salvatierra, Bolivia). *J. Archaeol. Sci. Rep.* **37**, 102880 (2021).

37. Stahl, P. W. Adventive vertebrates and historical ecology in the pre-Columbian neotropics. *Diversity* **1**, 151–165 (2009).

38. Hutterer, R. Archaeozoological remains (Vertebrata, Gastropoda) from prehispanic sites at Pailón, Bolivia. *Beiträge zur Allgemeinen und Vergleichenden Archäologie* **17**, 325–342 (1997).

39. Denevan, W. M. *The Aboriginal Cultural Geography of the Llanos de Mojos of Bolivia* (Univ. of California Press, 1966).

40. Dickau, R. et al. Diversity of cultivars and other plant resources used at habitation sites in the Llanos de Mojos, Beni, Bolivia: evidence from macrobotanical remains, starch grains, and phytoliths. *J. Archaeol. Sci.* **39**, 357–370 (2012).

41. Bruno, M. Carbonized plant remains from Loma Salvatierra, Department of Beni, Bolivia. *Zeitschrift für Archäologie Außereuropäischer Kulturen* **3**, 151–206 (2010).

42. Staller, J. E. In *Andean Foodways. The Latin American Studies Book Series* (ed. Staller, J. E.) 283–310 (Springer, 2021).

43. Eva, H. D. et al. A proposal for defining the geographical boundaries of Amazonia. In *Synthesis of the Results from an Expert Consultation Workshop Organized by the European Commission in Collaboration with the Amazon Cooperation Treaty Organization – JRC Ispra* (eds Eva, H. D. & Huber, O.) <https://research.wur.nl/en/publications/a-proposal-for-defining-the-geographical-boundaries-of-amazonia-s> (European Commission 2005).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025

Article

Methods

Phytolith processing and identification

Phytoliths were extracted from sediments following previously published methods⁴⁴. Phytoliths were identified and counted using a Zeiss Axioscope 40 light microscope at $\times 500$ magnification. Phytolith identifications were made using published material for the Neotropics^{45–49}, and by direct comparison with the phytolith reference collection of the Archaeobotany and Palaeoecology Laboratory (Department of Archaeology, University of Exeter, UK) and at the phytolith laboratory of ICTA-UAB (Universitat Autònoma de Barcelona). A minimum of 200 diagnostic randomly placed phytoliths were counted per slide. A full scan of slides was performed to detect the presence of squash, manioc and maize. The average size of attributes measured on *Oryza* glume phytoliths identified in the pond/canal systems followed the model proposed by Hilbert et al.⁵⁰. Overall, all glume phytolith mean width and height measurements were compared with the Monte Castelo site to assess the likelihood of a domesticated origin. Glume phytoliths identified in upper layers from sites at which *O. sativa* is currently cultivated were analysed using prediction calculations proposed by Zhao et al.⁵¹. We confirmed the presence of Asian domesticated rice on all upper layers. Overall, our analysis indicates that the origin of rice phytoliths from our samples was statistically similar to both wild botanical specimens (*Oryza latifolia* and *Oryza alta*) and lower layers (I–J) from the Monte Castelo site⁵⁰.

Pollen processing and identification

Samples for pollen analysis were treated following a protocol designed to improve the recovery of large pollen grains—in particular, those of cultigen⁵². Two tablets of the exotic marker *Lycopodium clavatum* were added to each sample to facilitate the calculation of pollen concentration per cubic centimetre⁵³. Pollen and spores were analysed using a Leica DMLB microscope at $\times 400$ and $\times 1,000$ magnification, and identifications were made using the modern pollen reference collection at the University of Reading, as well as the Neotropical pollen database⁵⁴ and specialized atlases^{55–57}. In every sample, a total of 300 randomly placed terrestrial pollen grains were counted.

Drone light detection and ranging

A light detection and ranging survey was conducted using a Zenmuse L1 sensor mounted on a Matrice 300 real-time kinematic (RTK) drone and a D-RTK 2 base station. We used a postprocessing kinematic solution rather than RTK for data correction, because of malfunctioning of the latter device. Four flights at an altitude of 100 m and speed of 6 m s⁻¹ were needed to cover the entire area; point density was 477 m⁻². The missions were planned with DJI Pilot 2, v.9.0.5.5. We set the sensor to detect three returns, its maximum limit, to ensure the recording of laser bounce on the ground through the tree canopy, which covered around 50% of the surveyed area. Data were processed using D-RTK 2 data in the postprocessing kinematic workflow of DJITerra software according to the Zenmuse L1 v.1.1 operation guidebook⁵⁸. Terramatch software v.023.014 was used in the Spatix environment to align datasets, correct trajectories, delete overlapping points and smooth noise points, following the workflow steps explained in the user guide.

Radiocarbon dating

Accelerator mass spectrometry radiocarbon dating was performed on seven samples from profile 690 at the Oxford Radiocarbon Accelerator Unit and Beta Analytic; dates are reported in Extended Data Table 1. The samples dated at the Oxford Radiocarbon Accelerator Unit were chemically pretreated using an acid–base–acid protocol for the insoluble humin fraction of sediments, and subsequently dated following their protocols⁵⁹. The same acid–base–acid protocol was used by Beta Analytic. Radiocarbon dates were calibrated using SHCAL20 (ref. 60),

modelled using the P_Sequence command and outlier modelling in OxCal v.4.4.4 (refs. 61–64). The code used is available in Supplementary Information.

Inclusion and ethics

The study included several South American researchers (S.Q., J.I., L.H., E.N. and M.R.) who contributed to various aspects of the research project. The research is locally relevant, and several local institutions (Gobernación del Beni, Universidad Autónoma del Beni and Alcaldía de Trinidad) have repeatedly expressed public support. We have a collaboration Agreement with CIBIOMA at Universidad Autónoma del Beni José Balliván for training of local students in phytolith analysis (we are currently setting up a laboratory in Trinidad). We have provided training and materials to Museo Etnoarqueológico Kenneth Lee in Trinidad. The type of study we performed did not require the approval of a local ethics review committee. The local and regional research relevant to our study has been taken into account in citations.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

All phytolith and pollen data supporting the findings of this study are available in Supplementary Information. Phytoliths were identified using the sources referenced in Methods. Pollen was identified using the Neotropical pollen database (<https://research.fit.edu/paleolab/pollen-database/>) and the sources referenced in Methods.

44. Lombardo, U., Ruiz-Pérez, J. & Madella, M. Sonication improves the efficiency, efficacy and safety of phytolith extraction. *Rev. Palaeobot. Palynol.* **235**, 1–5 (2016).
45. Piperno, D. R. *Phytoliths* (AltaMira Press, 2006).
46. Piperno, D. R. Identifying crop plants with phytoliths (and starch grains) in Central and South America: a review and an update of the evidence. *Quat. Int.* **193**, 146–159 (2009).
47. Iriarte, J. Assessing the feasibility of identifying maize through the analysis of cross-shaped size and three-dimensional morphology of phytoliths in the grasslands of southeastern South America. *J. Archaeol. Sci.* **30**, 1085–1094 (2003).
48. Watling, J. et al. Differentiation of neotropical ecosystems by modern soil phytolith assemblages and its implications for palaeoenvironmental and archaeological reconstructions II: Southwestern Amazonian forests. *Rev. Palaeobot. Palynol.* **226**, 30–43 (2016).
49. Dickau, R. et al. Differentiation of neotropical ecosystems by modern soil phytolith assemblages and its implications for palaeoenvironmental and archaeological reconstructions. *Rev. Palaeobot. Palynol.* **193**, 15–37 (2013).
50. Hilbert, L. et al. Evidence for mid-Holocene rice domestication in the Americas. *Nat. Ecol. Evol.* **1**, 1693–1698 (2017).
51. Zhao, Z., Pearsall, D. M., Benfer, R. A. & Piperno, D. R. Distinguishing rice (*Oryza sativa* Poaceae) from wild *Oryza* species through phytolith analysis, II: finalized method. *Econ. Bot.* **52**, 134–145 (1998).
52. Whitney, B. S., Rushton, E. A., Carson, J. F., Iriarte, J. & Mayle, F. E. An improved methodology for the recovery of *Zea mays* and other large crop pollen, with implications for environmental archaeology in the Neotropics. *The Holocene* **22**, 1087–1096 (2012).
53. Stockmarr, J. Tablets with spores used in absolute pollen analysis. *Pollen et Spores* **13**, 615–621 (1971).
54. Bush, M. & Weng, C. Introducing a new (freeware) tool for palynology. *J. Biogeogr.* **34**, 377–380 (2007).
55. Colinvaux, P. A. & De Oliveira, P. E. *Amazon: Pollen Manual and Atlas* (Harwood, 1999).
56. Lima Lorenti, F., et al. *Atlas Palinológico. Laboratório 14C - CENA/USP*. Eds: Fundação de Estudos Agrários Luiz de Queiroz-FEALQ (2017).
57. Roubik, D. W. & Moreno, P. J. E. *Pollen and Spores of Barro Colorado Island* (Missouri Botanical Garden, 1991).
58. DJI L1 Operation Guidebook V1.1 (DJI, 2022).
59. Brock, F., Higham, T., Ditchfield, P. & Ramsey, C. B. Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (Orau). *Radiocarbon* **52**, 103–112 (2010).
60. Hogg, A. G. et al. SHCal20 Southern Hemisphere calibration, 0–55,000 years cal BP. *Radiocarbon* **62**, 759–778 (2020).
61. Bronk Ramsey, C. Bayesian analysis of radiocarbon dates. *Radiocarbon* **51**, 337–360 (2009).
62. Bronk Ramsey, C. Dealing with outliers and offsets in radiocarbon dating. *Radiocarbon* **51**, 1023–1045 (2009).
63. Ramsey, C. B. & Lee, S. Recent and planned developments of the program OxCal. *Radiocarbon* **55**, 720–730 (2013).
64. Ramsey, C. B. Deposition models for chronological records. *Quat. Sci. Rev.* **27**, 42–60 (2008).

Acknowledgements This research was funded by the AHRC-FAPESP MoU-funded HERCA project (Human-Environment Relationships in pre-Columbian Amazonia), Arts and Humanities Research Council project no. AH/S001662/1 (F.M. and C.B.R.), Fundação de Amparo à Pesquisa do Estado de São Paulo project no. 2019/07794-9 (E.N.); and ERC Consolidator project DEMODRIVERS funded by the European Research Council, project no. 101043738 (U.L.). This article contributes to ICTA-UAB 'Maria de Maeztu Unit of Excellence' (no. CEX2019-0000940-M), and also to EarlyFoods (Evolution and impact of early food production systems), which has received funding from Agència de Gestió d'Ajuts Universitaris i de Recerca de Catalunya (no. SGR-Cat-2021, 00527). We acknowledge the support of the Bolivian Ministerio de Culturas y Turismo, Gobierno Autónomo Departamental del Beni, Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma del Beni José Ballivian and the owners of the properties on which the study sites are located: the indigenous community of Poza Honda and Doña Vivian Galindo Hinojosa; D. J. Leige, D. M. Saavedra, D. P. Suarez, D. F. Boehme and the association of Beni's cattle ranchers (FEGABENI). We also thank S. Ten from CIBIOMA, M. Gonzales from WCS and E. Chavez for help with logistics. We thank G. P. Guevara for producing the artwork in Extended Data Fig. 4 and J. C. Briceño for identifying the bones shown in Extended Data Fig. 9.

Author contributions U.L. planned and conceptualized the study, U.L., L.H., A.G.-R. and S.Q. undertook the fieldwork. L.H. carried out phytolith analysis. M.R. performed pollen analysis. M.B. carried out radiocarbon analysis. A.M.R.R. and C.B.R. performed Bayesian analysis. U.L. and J.I. wrote the first draft. C.B.R., K.D., A.M.R.R., M.R., J.G.W., E.N. and F.M. contributed to later drafts. F.M., E.N., C.B.R. and U.L. secured funding for this research.

Competing interests The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41586-024-08473-y>.

Correspondence and requests for materials should be addressed to Umberto Lombardo.


Peer review information *Nature* thanks Roland Fletcher, Dolores Piperno and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at <http://www.nature.com/reprints>.


Article



Extended Data Fig. 1 | Field view of a circular farm pond. The diameter of the pond is ca. 70 metres.



Article




Extended Data Fig. 3 | Phytolith profiles of causeway (675), fields (676, 679, 680, 706), ponds (689, 695), forest (702, 703, 704, 705) and canal (708). Horizontal bars represent percentages; circles correspond to presence of plant taxa lower than 1% in abundance. See Fig. 2 of the main text for location.



Extended Data Fig. 4 | Artwork representing how farm ponds and drainage canals were probably used for maize agriculture. Maize was planted around the pond and along the edges of the canals. Canals and ponds are here depicted together for simplicity. Credit: J. P. Guevara.

Article





Extended Data Fig. 5 | Profile of pond 690 with associated age/depth model. The pond was in use for ca. 300 years, from ca. 1250 to ca. 1550 Cal yrs. CE. Samples for radiocarbon ages have been collected along the yellow measuring tape shown in the figure.



Extended Data Fig. 6 | Pond cluster with location of profile 695. Most of the ponds are aligned and are connected by depressions/canals. Image provided by ESRI under the ArcGIS Pro license.

Article





Extended Data Fig. 8 | Microphotographs showing examples of key phytolith morphologies identified and their taxonomic and anatomical associations. (A-D) *Zea mays* cob, WAVY-TOP RONDEL (A-690 80-85 cm; B-690 40-45 cm; C-690 65-70 cm; D-666 65-65 cm); (E) Poaceae non-diagnostic to maize leaf/stem/inflorescence, spiny WAVY-TOP RONDEL (666 40-45 cm); (F) Poaceae leaf, CROSS VARIANT-1 (695 30-35 cm); (G) *Manihot* sp. secretory cell, HEART-SHAPED (677 20-25 cm); (H) cf. *Calathea* sp. rhizome, FLAT DOMED CYLINDER (690 30-35 cm); (I) *Oryza* sp. husk, DOUBLE-PEAK GLUME

(695 20-25 cm); (J) *Cucurbita* sp. rind, SCALLOPED SPHEROID (680 25-30 cm); (K) *Lagenaria* sp. rind, irregularly-shaped SCALLOPED SPHEROID (667 45-50 cm); (L) cf. *Euterpe* sp. all plant parts, large dense SPHEROID ECHINATE (667 30-35 cm); (M) *Bactris/Astrocaryum* all plant parts, CONICAL TO HAT-SHAPED BODY (665 20-25); (N) Multiple plants species (e.g. *Commelinaceae* and *Phaseolus* sp.) trichome, HOOK-SHAPED HAIR (665 30-35 cm); (O) *Marantaceae* seed, CILINDRICAL BODY (671 35-40 cm). Scale = 20 μ m.

Article



Extended Data Fig. 9 | Bones of *Synbranchus marmoratus* found in pond 695 at a depth of 70 cm. Bones of *S. marmoratus* are often found in monumental mounds. It hibernates in the mud and it is a predictable and ready available source of proteins. Scale bar in cm.

Extended Data Table 1 | Radiocarbon dates conducted on pond 690 and used in the associated Bayesian age depth model

Lab code	Material/ Pretreatment	depth	±	14C age	±	Calibrated date range		Modelled date					
						(cm)	(range)	(BP)	(1σ)	from (95.4%)	to (95.4%)	median	from (95.4%)
Beta 690512	Seed/ABA	12.5	2.5	Modern: F14C = 1.0188 ± 0.0038		1950	-						
Beta 690513	Charred material/ABA	12.5	2.5	280	30	1511	1800	1534		1505	1668		
OxA-42634	Leaf/ABA	37.5	2.5	447	18	1447	1612	1463		1446	1494		
OxA-42637	Leaf/ABA	70	0.5	679	19	1295	1394	1374		1315	1396		
OxA-42636	Leaf/ABA	92.5	2.5	1061	19	990	1130	1311		1264	1344		
Beta 690511	Leaf/ABA	112.5	2.5	860	30	1163	1276	1258		1233	1279		
OxA-42635	Charred material/ABA	120	0.5	800	19	1226	1285	1239		1221	1268		

The date ranges are given as highest probability density ranges at the 95.4% level; the modelled dates are taken from Model 1 referred to in supplementary information which is an age-depth model and so the dates are estimates for the dates of the sediment at the given depths rather than estimates of age of the samples where they are outliers; for the modelled dates the median date for the posterior distributions is also given. ABA stands for acid-base-acid.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
 - Only common tests should be described solely by name; describe more complex techniques in the Methods section.*
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
 - Give P values as exact values whenever suitable.*
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection DJI proprietary software (Terramatch version 023.014 and DJITerra version 3.8.0) has been used for the generation of the LIDAR DEM. C2 Version 1.8 has been used to produce phytolith graphs. ArcGis Pro version version 3.3.2 has been used to produce all the mpas.

Data analysis Radiocarbon ages have been calibrated and modelled using OxCal 4.4.4. The code is available in Supplementary Information

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

All the phytolith data supporting the findings of this study are available as supplementary files. Phytoliths have been identified using the sources referenced in methods. Pollen has been identified using the Neotropical pollen database (<https://research.fit.edu/paleolab/pollen-database/>) and the sources referenced in Methods.

Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender	N/A
Reporting on race, ethnicity, or other socially relevant groupings	N/A
Population characteristics	N/A
Recruitment	N/A
Ethics oversight	N/A

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	Data are quantitative. The study includes mapping of landscape elements based on visual analysis of remote sensing data and LIDAR; soil and subsoil sampling of forest and savannah; 14C of the samples from a farm pond; phytolith extraction and visual (microscope) analysis (counting of individual phytoliths) of samples gathered from forest and savannah (canals, fields and farm ponds); pollen extraction and visual (microscope) analysis (counting of individual pollen grains) of samples gathered from a farm pond.
Research sample	Stratigraphic profiles have been investigated both using a motor corer and excavations. Sediments sampled for phytolith and pollen analysis have been taken from profiles exposed during excavations. Samples are representative of the local environments where they have been taken.
Sampling strategy	Samples have been taken from stratigraphic profiles at different depths. Some profiles were sampled with a 5 cm resolution (i.e. pond 690), others at 10 cm resolution. The depth of each sample is indicated in the Y axe of each phytolith or pollen graph.
Data collection	Samples have been collected in the field and air-dried in Bolivia before being shipped. Charcoal fragments and vegetal remains for 14C have been collected in situ, enveloped in aluminium foil and stored in plastic bags. Field observations have been wrote down on a notebook. The researchers were aware of the study hypothesis at the time of sampling.
Timing and spatial scale	Sampling has been done in 2021. Lidar has been done in 2023. Samples have been taken in the Beni department, Bolivia.
Data exclusions	No data was excluded
Reproducibility	The experiments consisted in counting a standard number (200) of diagnostic phytoliths and pollen grains. This number is considered sufficient to be representative of the sample, therefore it is not standard practice to repeat the counting.
Randomization	Sampling was not completely random. We identified several potential locations to sample on satellite imagery, we choose those to excavate based on their accessibility and ownership of the land. However, none of these criteria affect the representativeness of our sample.
Blinding	Sampling was not blind because we sampled soil and subsoil, so we knew the origin of each sample. In the lab samples where coded with numbers. Sample extraction and phytolith counting was blind because the origin of the sample was unknown during these steps. Pollen counting was not blind, because we analyzed pollen from only 1 profile, and this was known.

Did the study involve field work? Yes No

Field work, collection and transport

Field conditions	Sampling has always being performed during the dry season, between August and October
------------------	---

Location	Fieldwork took place in the Beni department, Bolivia. The area surveyed is enclosed in a square area: up right corner lat -14° 50" Lon -64° 10"; down left corner Lat -15°; Lon -64° 45". Average elevation 180 m a s l. All sampling was performed on land.
Access & import/export	Field sites have been accessed with the permission of the land owner. Authorizations for export of samples have been obtained by the Bolivian Ministry of Medioambiente y Agua (MMAYANMABCCGDF/DGBAP/MEGN°0342/2021) Autorization for flying the drone with the LIDAR has been obtained by DSO - OPERACIONES, Dirección General de Aeronáutica Civil de Bolivia (1ZNBL13)
Disturbance	All the savannah coring and excavation have been performed on land used for pasture with almost complete absence of wild fauna. The test pits in the forest were very small and we didn't produce any noise during the excavation or the sampling. The excavation pits were refilled with the excavated sediments in order to restore the aspect of the sites previous to the excavation.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems		Methods	
n/a	Involved in the study	n/a	Involved in the study
<input checked="" type="checkbox"/>	Antibodies	<input checked="" type="checkbox"/>	ChIP-seq
<input checked="" type="checkbox"/>	Eukaryotic cell lines	<input checked="" type="checkbox"/>	Flow cytometry
<input checked="" type="checkbox"/>	Palaeontology and archaeology	<input checked="" type="checkbox"/>	MRI-based neuroimaging
<input checked="" type="checkbox"/>	Animals and other organisms		
<input checked="" type="checkbox"/>	Clinical data		
<input checked="" type="checkbox"/>	Dual use research of concern		
<input checked="" type="checkbox"/>	Plants		

Plants

Seed stocks	Report on the source of all seed stocks or other plant material used. If applicable, state the seed stock centre and catalogue number. If plant specimens were collected from the field, describe the collection location, date and sampling procedures.
Novel plant genotypes	Describe the methods by which all novel plant genotypes were produced. This includes those generated by transgenic approaches, gene editing, chemical/radiation-based mutagenesis and hybridization. For transgenic lines, describe the transformation method, the number of independent lines analyzed and the generation upon which experiments were performed. For gene-edited lines, describe the editor used, the endogenous sequence targeted for editing, the targeting guide RNA sequence (if applicable) and how the editor was applied.
Authentication	Describe any authentication procedures for each seed stock used or novel genotype generated. Describe any experiments used to assess the effect of a mutation and, where applicable, how potential secondary effects (e.g. second site T-DNA insertions, mosaicism, off-target gene editing) were examined.