

# *Machine learning for improvement of upper-tropospheric relative humidity in ERA5 weather model data*

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Wang, Z., Bugliaro, L., Gierens, K., Hegglin, M. I. ORCID: <https://orcid.org/0000-0003-2820-9044>, Rohs, S., Petzold, A., Kaufmann, S. and Voigt, C. (2025) Machine learning for improvement of upper-tropospheric relative humidity in ERA5 weather model data. *Atmospheric Chemistry and Physics*, 25 (5). pp. 2845-2861. ISSN 1680-7324 doi: 10.5194/acp-25-2845-2025 Available at <https://centaur.reading.ac.uk/121823/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <https://doi.org/10.5194/acp-25-2845-2025>

To link to this article DOI: <http://dx.doi.org/10.5194/acp-25-2845-2025>

Publisher: Copernicus Publications

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online



# Machine learning for improvement of upper-tropospheric relative humidity in ERA5 weather model data

Ziming Wang<sup>1,2</sup>, Luca Bugliaro<sup>1</sup>, Klaus Gierens<sup>1</sup>, Michaela I. Hegglin<sup>3,4</sup>, Susanne Rohs<sup>5</sup>,  
Andreas Petzold<sup>5</sup>, Stefan Kaufmann<sup>1</sup>, and Christiane Voigt<sup>1,2</sup>

<sup>1</sup>Institute of Atmospheric Physics, Deutsches Zentrum für Luft- und Raumfahrt (DLR),  
Oberpfaffenhofen, 82234, Germany

<sup>2</sup>Institute for Atmospheric Physics, Johannes Gutenberg University Mainz, Mainz, 55128, Germany

<sup>3</sup>Institute of Climate and Energy Systems 4 – Stratosphere (ICE-4),  
Forschungszentrum Jülich, Jülich, 52428, Germany

<sup>4</sup>Department of Meteorology, University of Reading, Reading, RG6 6ET, UK

<sup>5</sup>Institute of Climate and Energy Systems 3 – Troposphere (ICE-3),  
Forschungszentrum Jülich, Jülich, 52428, Germany

**Correspondence:** Ziming Wang (ziming.wang@dlr.de)

Received: 1 July 2024 – Discussion started: 12 July 2024

Revised: 18 November 2024 – Accepted: 14 January 2025 – Published: 7 March 2025

**Abstract.** Knowledge of humidity in the upper troposphere and lower stratosphere (UTLS) is of special interest due to its importance for cirrus cloud formation and its climate impact. However, the UTLS water vapor distribution in current weather models is subject to large uncertainties. Here, we develop a dynamic-based humidity correction method using an artificial neural network (ANN) to improve the relative humidity over ice (RHi) in ECMWF numerical weather predictions. The model is trained with time-dependent thermodynamic and dynamical variables from ECMWF ERA5 and humidity measurements from the In-service Aircraft for a Global Observing System (IAGOS). Previous and current atmospheric variables within  $\pm 2$  ERA5 pressure layers around the IAGOS flight altitude are used for ANN training. RHi, temperature, and geopotential exhibit the highest impact on ANN results, while other dynamical variables are of low to moderate or high importance. The ANN shows excellent performance, and the predicted RHi in the UT has a mean absolute error (MAE) of 5.7 % and a coefficient of determination ( $R^2$ ) of 0.95, which is significantly improved compared to ERA5 RHi (MAE of 15.8 %;  $R^2$  of 0.66). The ANN model also improves the prediction skill for all-sky UT/LS and cloudy UTLS and removes the peak at RHi = 100 %. The contrail predictions are in better agreement with Meteosat Second Generation (MSG) observations of ice optical thickness than the results without humidity correction for a contrail cirrus scene over the Atlantic. The ANN method can be applied to other weather models to improve humidity predictions and to support aviation and climate research applications.

## 1 Introduction

The atmospheric region of the upper troposphere and lower stratosphere (UTLS) in the tropics (Dessler and Sherwood, 2009) and the extratropics (Gettelman et al., 2011) plays a crucial role in the climate system. Within the UTLS, atmospheric humidity significantly influences the radiation budget at the top of the atmosphere (TOA) (Riese et al., 2012). In fact, water vapor is the dominant atmospheric long-

wave absorber in the context of the global greenhouse effect (Schmidt et al., 2010). Further, observed increases in stratospheric water vapor (Hegglin et al., 2014) contribute to both stratospheric cooling and tropospheric warming (Forster and Shine, 2002) and act as positive feedback to surface temperature (Tao et al., 2023). Relative humidity (RH) over ice (RHi)  $> 100\%$  or ice supersaturation is of major importance for the formation and persistence of natural cirrus and aircraft-

induced contrail cirrus (Kärcher, 2018). Cirrus in this region can survive for hours if ambient conditions are ice supersaturated (Zhao et al., 2023), and they can have a positive cloud radiative effect on climate (Gasparini et al., 2020). Hence, accurate observations and representations of UTLS water vapor (Hegglin et al., 2014) are essential for climate and weather research.

During the past few decades, many in situ humidity measurements from aircraft (Krämer et al., 2009; Diao et al., 2015; Kaufmann et al., 2018), lidar (Groß et al., 2014; Krüger et al., 2022), balloon-borne instruments (Heymsfield et al., 1998; Dickson et al., 2010; Rollins et al., 2014), and polar-orbiting satellite instruments (Lamquin et al., 2012; Hegglin et al., 2013) show that high RHi may often occur in the UT. However, these observations are limited in space and time, and the uncertainties are relatively high (Gierens et al., 2020). Important in situ humidity data can also be provided by in-service passenger aircraft (Petzold et al., 2020; Reutter et al., 2020), but three-dimensional fields of RHi and dynamics for large geographic regions are currently only available from numerical weather prediction (NWP) models, for instance the Integrated Forecasting System (IFS) at the European Centre for Medium-Range Weather Forecasts (ECMWF, 2016) and ICOsahedral Non-hydrostatic (ICON; Zängl et al., 2015; Seifert and Siewert, 2024) at the German Weather Service. A wet bias in the RHi of the extratropical LS has been identified in the operational ECMWF IFS forecast and analysis data, as observed in comparison with in situ measurements from research aircraft (Kaufmann et al., 2018) and from Civil Aircraft for the Regular Investigation of the atmosphere Based on an Instrument Container (CARIBIC) passenger aircraft flights (Dyroff et al., 2015). In contrast, a dry bias of RHi is observed in the cloudy UT when compared with aircraft measurements in the In-Service Aircraft for a Global Observing System (IAGOS) (Teoh et al., 2022). The utilization of the saturation adjustment process (Tompkins et al., 2007), wherein supersaturation relaxes to saturation upon cloud formation, results in a systematic underestimation of both the frequency and the magnitude of ice supersaturation at cruise altitudes within NWPs and global climate models (Sperber and Gierens, 2023).

There is ongoing debate on the physical explanations of the NWP humidity bias in the UTLS, which is a crucial factor to consider for the improvement of atmospheric humidity prediction. According to Kunz et al. (2014), the influences of dynamical transport processes are challenging for the simulations of UTLS humidity distribution. Backward-trajectory analyses reveal a positive relationship between the moist bias at the aircraft flight level and air masses originating from high northern latitudes in the LS (Dyroff et al., 2015). The relationships between uncertainty in atmospheric mixing and the simulated composition of water vapor in the LS, as well as the radiative consequences in the UTLS, are highlighted by Krüger et al. (2022) and Riese et al. (2012), respectively. Small-scale stratospheric intrusions, which are

frequently observed in the UTLS but are unsolved by the NWP model, are another possible source of moisture bias (Dyroff et al., 2015). Numerical diffusion, which can easily smoothen the gradients of humidity across the hydopause, is also a possible reason for the moist bias in the LS (Stenke et al., 2008). Interestingly, Woiwode et al. (2020) show that there is little dependency of the moist bias on temporal or vertical model resolution in the ECMWF IFS analysis and forecast data.

The assimilation of observations into NWP models is the state-of-the-art way to improve weather forecast (Lawrence et al., 2019; van der Linden et al., 2020). However, as the primary data source for the assimilation system, calibrating RH instruments at temperatures below 0 °C is challenging, with the difficulty increasing as temperatures drop further. A great deal of effort has also been focused on post-processing of NWP data to improve the accuracy of atmospheric humidity and ice supersaturation prediction, as well as contrail cirrus prediction, utilizing long-term aircraft measurements from IAGOS (Teoh et al., 2020; Wolf et al., 2025). Teoh et al. (2022) employ in situ measurements from IAGOS to formulate a correction method for ERA5 RHi fields. With this method, the probability density function (PDF) of ERA5-corrected RHi inside ice supersaturation closely aligns with IAGOS measurements. Another humidity bias correction, also aiming to achieve consistency between IAGOS and ERA5 through a multivariate quantile approach, results in a notable reduction in the RHi bias (Wolf et al., 2025).

Gierens and Brinkop (2012) investigate the distributions of the dynamical quantities – divergence, relative vorticity, and vertical velocity – from ECMWF IFS within and outside ice-supersaturated regions and notice distinct patterns. Gierens et al. (2020) postulate that a more accurate prediction of ice supersaturation in NWP models may be achievable by further incorporating dynamical atmospheric fields with ERA5 RHi in a general regression method. Wilhelm et al. (2022) also suggest the possibility of basing an improved forecast of persistent contrails not only on the traditional quantities of temperature and RHi, but also on these dynamical proxies. In a recent study, Hofer et al. (2024) show that dynamical proxies taken only at the time and location of the forecast are insufficient for an improved prediction of ice supersaturation. However, they note the potential for improving RHi predictions by incorporating additional forecast data from earlier time points and upstream areas.

When improving the quality of meteorological data, machine learning techniques are widely used nowadays. Kadow et al. (2020) have demonstrated the skill of artificial intelligence in reconstructing surface temperatures when combined with climate model data. A machine-learning-based approach trained directly from historical NWP reanalysis data is introduced by Lam et al. (2023) to predict hundreds of weather variables at a remarkable speed. It outperforms the most accurate operational systems on 90 % of the verification tests, even without special consideration of vertical transport.


Teoh et al. (2020) also suggest in their outlook on RHi correction that further effort can be made to explore machine learning techniques to improve the accuracy of the ERA5 humidity fields.

This paper aims at improving predictions of atmospheric humidity, in particular RHi and ice supersaturation, in the UTLS region starting from ERA5 fields using machine learning. The previous humidity corrections of Teoh et al. (2022) and Wolf et al. (2025) for ERA5 model data were based on regression fitting methods using IAGOS observations but neglected the temporal evolution of dynamical quantities in the horizontal and vertical directions that led to the humidity bias. Targeting that gap, we develop an artificial neural network (ANN) model to correct relative humidity (and specific humidity in the Supplement) from ERA5, leveraging thermodynamic conditions and dynamical quantities from ERA5, along with measured water vapor data from IAGOS above the Atlantic Ocean, Europe, and Africa in 2020. The investigation is guided by three specific questions:

1. To what extent do atmospheric states impact the subsequent evolution of humidity fields?
2. Is it feasible to develop a dynamic-based machine learning method to correct the humidity bias in the UTLS?
3. How do the outcomes of the new method influence the ability to forecast ice supersaturation?

Finally, we apply the improved humidity fields for computing the optical properties of contrail cirrus in a particular situation using the contrail cirrus prediction (CoCiP) model and compare the simulation results to satellite observations.

This paper is outlined as follows: Sect. 2 provides an overview of the IAGOS humidity measurements (Sect. 2.1), ERA5 data, and IFS data as input to the ANN model and for the application (Sect. 2.2); the collocation procedure of the measurement data with ERA5 (Sect. 2.3) and the initial comparison (Sect. 2.4); the contrail cirrus prediction (CoCiP) model (Sect. 2.5); and satellite remote sensing techniques for retrieving the microphysical properties of cirrus clouds (Sect. 2.6). In Sect. 3, the concept of the temporal dependence of RHi on the evolution of meteorological parameters, the development of the RHi improvement model, and the importance of the selected synoptic variables for RHi prediction are explained in detail. The evaluation of the RHi improvement model using different metrics is presented in Sect. 4. The corresponding information for specific humidity is provided in the Supplement. Thereafter, Sect. 5 assesses the impact of the ANN humidity correction on the simulations of contrail cirrus in a case study. The conclusions are summarized in Sect. 6.



**Figure 1.** Number of IAGOS measurements per  $2^\circ \times 2^\circ$  latitude-longitude grid box between 200 and 400 hPa over the Atlantic Ocean, Europe, and Africa for the year 2020. The measurements are filtered based on data quality. See the text for details.

## 2 Data and application approaches

### 2.1 In-service Aircraft for a Global Observing System (IAGOS)

The In-service Aircraft for a Global Observing System (IAGOS; Petzold et al., 2015) is a European research infrastructure that implements instruments on long-range aircraft of internationally operating airlines to provide long-term in situ measurements of trace gases and meteorological conditions. These measurements are very valuable for the purpose of this study as most flight tracks are situated at heights between 9 and 13 km in the UTLS region. All aircraft within IAGOS have been equipped with a platinum sensor for temperature measurements with an accuracy of  $\pm 0.5$  K and a collocated capacitive sensor for monitoring RHi with an uncertainty of 5 % to 10 % (Petzold et al., 2020). The temperature detected at the sensor is transferred to the air temperature  $T_{\text{IAGOS}}$  by accounting for the (incomplete) adiabatic heating and the inlet heating. RHi is then derived using the measured water vapor mixing ratio, pressure, and  $T_{\text{IAGOS}}$  based on the saturation water vapor pressure equation from Sonntag (1994). The uncertainty in RHi increases with decreasing temperature due to a slower response time. In the dry conditions ( $\text{RHi} < 10\%$ ) of the LS, the sensor has only limited accuracy (Rolf et al., 2023), so these data have been excluded for further evaluation. The temporal resolution of IAGOS measurements amounts to 4 s.

The global distribution of IAGOS data is not uniform in every region since it is dependent on preferred flight routes and weather conditions. Western Europe and the eastern North Atlantic region (NAR) show a high density in IAGOS data; therefore we focus on this domain (Fig. 1, between

30° W and 50° E). We did not include the western regions of NAR, from where humid air is often advected, but instead focused on moisture originating and transported from lower-atmospheric levels up to cruise altitude. In addition, we aim to cover the latitude range between 80° N and the Equator because ice supersaturation occurs very frequently in the UT of the tropics. The geographic position of the aircraft, time, data quality flags, ambient pressure, temperature  $T_{\text{IAGOS}}$  (Berkes et al., 2017), and  $\text{RH}_{\text{IAGOS}}$  from IAGOS in the year 2020 are collected to produce the output humidity data set of the ANNs. We use only the IAGOS measurements that fulfill the following criteria: the IAGOS quality flag is not “limited” or “invalid”, and measurements are located between 0 and 80° N and 30° W and 50° E and between 400 and 200 hPa. The distribution of  $\text{RH}_{\text{IAGOS}}$  shows high-density values that gradually decrease starting from 110 % and drop significantly as they approach 150 % (see Fig. S2 in the Supplement), a trend consistent with findings in Wolf et al. (2025, Fig. 3) and Teoh et al. (2024, Fig. S4).

## 2.2 ECMWF reanalysis and forecast

Meteorological data for the year 2020 in the same region in Sect. 2.1 are sourced from the ERA5 reanalysis data, obtained from the ECMWF Copernicus Climate Data Store (Hersbach et al., 2020). The assimilation system takes new observations and combines them with IFS forecast data from 12 h before the given time to make the best estimate of the current state of the atmosphere. ERA5 data are on an equidistant latitude–longitude grid of 0.25° resolution with an hourly output on 37 pressure levels. Hourly atmospheric parameters on pressure levels between 200 and 250 hPa with a 25 hPa spacing and between 250 and 400 hPa with a 50 hPa spacing are used for model training. The IFS forecast data (137 model levels) are used for predicting contrail cirrus (see Sect. 5). Using pressure level data for ANN training reduces the size of the training data set and saves model training time.

We use the following thermodynamic parameters from ERA5 as the inputs for the ANN model: temperature  $T_{\text{ERA5}}$ ,  $\text{RH}_{\text{ERA5}}$  (in the main text), and specific humidity  $q_{\text{ERA5}}$  (in the Supplement). The saturation water vapor pressure equation from Alduchov and Eskridge (1996) is used here to calculate  $\text{RH}_{\text{ERA5}}$  from  $q_{\text{ERA5}}$ . In addition, specific cloud ice water content “ciwc” from ERA5 is utilized to differentiate between the cirrus and cirrus-free regions. Here, ciwc represents the mass of cloud ice particles per kilogram of moist air, averaged over a grid box. It is estimated using the prognostic equations of the cloud scheme (Tiedtke, 1993; Forbes and Tompkins, 2011; Forbes and Ahlgrimm, 2014), which account for cloud ice growth through deposition. As shown in Table 1, this study also considers dynamical parameters, including geopotential ( $z$ ) for adiabatic shifts, vertical velocity ( $w$ ) in  $\text{Pa s}^{-1}$  representing vertical air mass motion, divergence ( $d$ ) indicating air spread or convergence, horizontal wind speed ( $u$  and  $v$ ) in  $\text{m s}^{-1}$ , and relative and potential

vorticity (“vo” and “pv”) characterizing air rotation and stratosphere–troposphere exchanges. The use of  $pv$  specifically helps identify the dynamical tropopause and distinguish between UT and LS.

## 2.3 Data collocation


The ERA5 grid boxes that are closest to the IAGOS observations in both time and space are selected to align with the IAGOS-measured humidity and temperature data sets between 400 and 200 hPa. In contrast to other studies (Wolf et al., 2025; Hofer et al., 2024), this study also uses the temporal evolution of meteorological conditions before the IAGOS observation time. Specifically,  $\text{RH}$  in the UTLS is influenced by horizontal and vertical air motions like air mass uplift (Diao et al., 2015) in convection or frontal systems and stratospheric intrusions. To account for these, thermodynamic and dynamical data up to 6 h prior to the IAGOS data acquisition time, with 1 h intervals and within two pressure levels above and below the IAGOS acquisition location, are linked with  $\text{RH}_{\text{IAGOS}}$  at the IAGOS acquisition time and location. These ERA5 variables are vertically interpolated to match the IAGOS location based on pressure levels. While the ERA5 data retain their original temporal and spatial resolution, the water vapor data measured by IAGOS, which include several data points within a single ERA5 grid box, are averaged. This averaging reduces the autocorrelation in the measured data due to the response time of the sensor, accounts for internal ERA5 grid box variability, and maintains a proportion of ice supersaturation after averaging. This collocation of model meteorological variables and measured humidity values from the year 2020 comprises 3.99 million individual data points. To ensure that the training set consists of data from time periods that do not overlap with those used for either validation or testing, we now use 4 d of data to build the ANN model, followed by a 1 d gap, and then 1 d of data for validation or testing. This method accounts for considerable variability and sharp gradients in the humidity fields and can thus help to estimate realistic atmospheric humidity distributions for comparisons and model application.

## 2.4 Initial ERA5 $\text{RH}$ evaluation using IAGOS

We first compare and quantify the difference between ERA5 and in situ measurements provided by IAGOS with respect to temperature and specific and relative humidity.  $\text{RH}$  in cirrus clouds in NWP can have a low bias due to the application of saturation adjustment in cloud parameterizations (ECMWF, 2016); hence we differentiate between model clear (cloudy) conditions using ciwc equal to zero for all the current and  $\pm 2$  pressure layers from ERA5. We further distinguish between UT (LS) dependent on the threshold  $pv$  smaller than 2 PVU. This means we consider the dynamical tropopause as done for instance in Reutter et al. (2020). In the following,

**Table 1.** Overview of the variables used in this study. The spatial resolution of ERA5 is  $0.25^\circ$ . The vertical resolution of ERA5 on pressure levels is 25–50 hPa. The original temporal resolution of ERA5 and IAGOS is 1 h and 4 s. Study regions include the Atlantic, Europe, and Africa.

Source	Variable (*RHi ANN)	Description	Unit
ERA5 (seven conditions)	$*T_{\text{ERA5}}$	air temperature	K
	ciwc	specific cloud ice water content	$\text{Kg kg}^{-1}$
	$*\text{RHi}_{\text{ERA5}}$	relative humidity w.r.t. ice	%
	$q_{\text{ERA5}}$	specific humidity	$\text{g kg}^{-1}$
	$*z$	geopotential	$\text{m}^2 \text{s}^{-2}$
	$*w$	vertical velocity	$\text{Pa s}^{-1}$
	$*d$	divergency of wind	$\text{s}^{-1}$
	$*u$	eastward component of wind	$\text{m s}^{-1}$
	$*v$	northward component of wind	$\text{m s}^{-1}$
	$*\text{vo}$	relative vorticity	$\text{s}^{-1}$
	$\text{pv}$	potential vorticity	$\text{s}^{-1}$
	time	h	1
IAGOS at the current time	level	pressure	hPa
	$*\text{RHi}_{\text{IAGOS}}$	relative humidity w.r.t. ice	1
	$T_{\text{IAGOS}}$	air temperature	K
	pressure	air pressure	Pa



**Figure 2.** Comparisons of  $\text{RHi}_{\text{ERA5}}$  against  $\text{RHi}_{\text{IAGOS}}$  in (a) clear-sky and cloudy UTLS, (b) cloudy UTLS, (c) clear-sky UTLS, and (d) clear-sky and cloudy LS in the test data set between 200 and 400 hPa over the Atlantic, Europe, and Africa for the year 2020.

we focus on RHi (for comparison of other parameters, see also Sects. S1 and S4).

Figure 2 shows the comparison of  $\text{RHi}_{\text{ERA5}}$  and  $\text{RHi}_{\text{IAGOS}}$  separated between upper-stratospheric and lower-stratospheric conditions or between cloudy and clear-sky conditions, respectively. Here we use the test data set from the ERA5–IAGOS collection created in Sect. 2.3, which is

the same data set used for verifying the ANN model in Sect. 4.1. In the all-sky UT (Fig. 2a) and the cloudy UTLS (Fig. 2b), ERA5 RHi data show a considerably dry bias compared to IAGOS data, with mean absolute errors (MAEs) of 15.82 % and 16.28 %, respectively. RHi and the magnitude of ice supersaturation in ERA5 are underestimated. In addition, a partially artificial occurrence accumulation peak exists in the ERA5 data set at  $\text{RHi}_{\text{ERA5}} = 100\%$ . In  $\text{RHi}_{\text{IAGOS}}$ , a small peak is observed between 100 % and 110 % under cloudy conditions (Sanogo et al., 2024). However, much of the accumulation peak in the ERA5 data is attributed to the cloud saturation adjustment in NWP models. Nevertheless,  $\text{RHi}_{\text{ERA5}} > 100\%$  is also observed, either in partly cloudy model boxes or in clear-sky boxes, where only a fraction of the box is cloudy (with  $\text{RHi}_{\text{ERA5}} = 100\%$ ), and the clear-sky part is supersaturated due to the time required for the ice nucleation process. Consequently,  $\text{RHi}_{\text{ERA5}}$  values greater than 100 % can occur in cloudy conditions as well. In the clear-sky UTLS (Fig. 2c) and the all-sky LS (Fig. 2d) regions, MAEs are 11.21 % and 9.78 %, respectively. In the all-sky LS (Fig. 2d), few RHi data  $> 100\%$  have been measured by IAGOS, with most observations concentrated at low RHi values. In general, the extent and the degree of ice supersaturation underestimated by ERA5 are in line with the findings by Dyroff et al. (2015) for ECMWF analysis and forecast data. The comparison of ERA5 and IAGOS RHi serves as the motivation for our study, aiming to improve humidity predictions by NWPs.

## 2.5 Contrail cirrus prediction (CoCiP) model

The contrail cirrus prediction (CoCiP) model is used to predict the contrail cirrus cover and examine the contrail radiative forcing induced by individual flights (Schumann, 2012; Schumann et al., 2017, 2021b; Voigt et al., 2017, 2022; Teoh et al., 2024). The contrail model uses traffic data from the North Atlantic tracks for the Shanwick Oceanic Control area. When the ambient temperatures fall below the Schmidt–Appleman criterion threshold (Schumann, 1996) at two successive flight waypoints, a contrail segment forms. Contrail initial water content, width, and depth are determined by aircraft properties and emissions (non-volatile particulate matter). Plume dispersion is a function of turbulence, wind shear, and induced heating. RHi inside contrail plumes is set at saturation, and the ice water content of contrails grows or decreases in response to the ambient humidity. A Runge–Kutta integration simulates the contrail evolution until the end of its life by ambient drying or particle losses from aggregation and sedimentation. The contrail life ends when the maximum contrail lifetime of 24 h is reached, the ice number concentration is less than the background ice nuclei ( $< 10^3 \text{ m}^{-3}$ ), or the ice optical thickness (IOT) is less than  $10^{-6}$ . CoCiP simulations account for humidity exchange between contrails and the background air and the overlap of contrails above or below clouds present in the meteorological data from the NWP forecast (Schumann et al., 2021a). CoCiP's limitation in comparison to general circulation models is its absence of atmospheric interaction and feedback (Chen et al., 2012; Burkhardt et al., 2018; Bickel et al., 2020). In Sect. 5 of this study, we present an exemplary application of the model for RHi correction derived in this paper to contrail simulations and show its effect on contrail properties. This is done by performing two CoCiP runs: for the reference run, we use NWP data from ECMWF IFS for the contrail case on 14 April 2021 over the NAR within the ECLIF3 campaign (Märkl et al., 2024); for the second run, we correct the NWP humidity data with the ANN proposed in this work in the same situation.

## 2.6 Satellite remote sensing

CoCiP simulations are compared to spaceborne data from the SEVIRI radiometer (3 km sampling distance at nadir) aboard the geostationary Meteosat Second Generation (MSG) satellite in Sect. 5. To derive ice cloud properties, Cirrus Properties from SEVIRI (CiPS; Strandgren et al., 2017) is used. It consists of a set of ANNs trained on SEVIRI thermal observations, Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) cloud products, and ECMWF ERA-Interim surface temperature data and auxiliary data to retrieve IOT for identified cirrus clouds. Specifically developed for thin cirrus clouds, CiPS has been validated against CALIPSO, achieving detection rates of 20 %, 70 %, and 85 % for ice clouds with IOT values of 0.01, 0.1, and 0.2, respectively. For IOT between 0.35 and 1.8, CiPS demonstrates a

MAE smaller than 50 %, and MAE increases for IOT values between 0.07 and 0.35.

## 3 ANN model development

### 3.1 The temporal dependence of measured humidity on individual meteorological parameters

Which meteorological parameters should be chosen and at which time and pressure level for training the RHi improvement model? To answer this question, the dependence of measured RHi on meteorological variables at preceding times and surrounding pressure levels is considered by reviewing the sources of RHi bias in the UTLS within ECMWF data. As outlined in Dyroff et al. (2015), this bias is linked to air masses residing near the aircraft's flight level of approximately 230 hPa in high northern latitudes, likely influenced by air mass vertical intrusions and horizontal transport. This points out the connection between RHi and the temporal evolution of meteorological parameters.

Based on the physical definition of RHi, a negative correlation between  $\text{RH}_{\text{IAGOS}}$  and  $T_{\text{ERA5}}$  is expected because RHi is the ratio of the partial vapor pressure of water vapor to the saturation vapor pressure with respect to ice, the latter of which increases with temperature. For dynamical variables, the positive relationship between increased geopotential  $z$  values and  $\text{RH}_{\text{IAGOS}}$  is in accordance with the findings of Wilhelm et al. (2022). In addition, parameters such as vertical wind  $w$ , divergence  $d$ , horizontal wind speed components  $u$  and  $v$ , relative vorticity  $\text{vo}$ , and potential vorticity  $\text{pv}$  help represent the dynamical conditions at a given time and place that influence relative humidity in the model. For instance, an upward motion (negative values of  $w$  in ERA5) results in cooling and a decrease in RHi and promotes ice supersaturation. A relatively strong horizontal air mass movement with large divergence is typical for ice supersaturation. Large negative values of vorticity in anticyclonic systems are again also typical for supersaturation (Gierens et al., 2020). These connections suggest the potential to improve the RHi prediction by considering not only traditional thermodynamic variables like temperature but also dynamical proxies and their temporal evolution. Further computations of the Pearson correlation coefficient between  $\text{RH}_{\text{IAGOS}}$  and temporal meteorological variables from ERA5 and the impact of including data distributions from hours before the current time on improving the network prediction are explained in Sect. S2.

To balance information richness and modeling efficiency, only the current IAGOS humidity fields and ERA5 data at the current time of the IAGOS measurement – and 2 and 6 h time lags prior to IAGOS data acquisition – and  $\pm 1$  to  $\pm 2$  pressure layers from ERA5 are selected as input variables to account for the typical lifespans of water vapor transport mechanisms, including deep convection, warm conveyor belt uplift regimes, and slowly ascending flows (Wang et al., 2024). Notably,  $\text{pv}$  is not provided in ECMWF model-level

data and is subsequently excluded from the input data set during further training of the ANN model. In summary, the ANN model is trained with the variables shown in Table 1. The relevance of each input atmospheric variable to the RHi prediction model developed in Sect. 3.2 is discussed in Sect. 3.3.

### 3.2 Development and training of the ANN model for humidity improvement

An ANN is composed of a large number of units that exchange information with each other in a similar structure and function to neural networks in human brains. A basic ANN model contains three types of layers: an input layer, one or more hidden layer(s), and an output layer. Each layer is made up of neurons. Neurons receive the weighted sum of the results of the previous layer's neurons, use it as the argument of an activation function, and forward the results to the following layer. The feed-forward ANN used in this study employs a learning technique called back propagation, where the outputs are compared to the target values to calculate the differences in the form of the loss function. The error is then fed back to modify the weights and bias of each neuron based on the optimization method (see, e.g., Ma et al., 2020).

Here, the ANN model for RHi is trained using a large set of atmospheric variables obtained from ERA5 reanalysis and RHi measured from IAGOS, as explained in the previous section. The ANN learns to reproduce the nonlinear statistical relationships between the selected series of meteorological variables and humidity fields iteratively, adjusting its parameters until it can robustly and accurately predict RHi in the UTLS. This procedure considers not only  $RHi > 100\%$  to investigate ice supersaturation but also the full range of RHi to provide sufficient data for model building.

Based on the temporal dependence of measured humidity on individual meteorological parameters discussed in Sect. 3.1, the input variables for the ANN encompass  $RHi_{ERA5}$ ,  $T_{ERA5}$ ,  $z$ ,  $w$ ,  $d$ ,  $u$ ,  $v$ , and  $vo$ . They are extracted from the ERA5 fields (Sect. 2.2) at the time of the IAGOS observation, as well as 2 and 6 h prior, at the geographical and vertical location of the IAGOS measurement, along with  $\pm 1$  to  $\pm 2$  ERA5 pressure levels. The output (target) variable of the ANN is  $RHi_{IAGOS}$ . The ANN model consists of 56 inputs, derived from eight meteorological variables across seven conditions: the current time and level, two time lags ( $-2$  and  $-6$  h) for the current level, and four ERA5 pressure levels surrounding the IAGOS cruise altitude ( $-2$ ,  $-1$ ,  $+1$ ,  $+2$  levels) for the current time. They are summarized in Table 1. We apply min–max normalization to both input and output data, which prevents features with larger ranges from dominating and improves convergence speed during model training. We use three hidden layers, each with 100 neurons and a He weight initializer (He et al., 2015), along with batch normalization between layers to improve generalization. The humidity output is referred to as  $RHi_{ANN}$ . The rectified linear unit (Relu) serves as the activation function for the hidden

layers, while a linear function is used for the output layer. The mean squared error (MSE) is adopted as a loss function, and the ANN model is optimized using stochastic gradient descent with a learning rate of 0.001, decay of  $10^{-5}$ , and momentum of 0.99 after several tests. Training of the ANN model is executed with batch sizes of 1024 and 100 epochs. In Sect. 2.3, a period of 4 consecutive days of samples from the ERA5–IAGOS collection is allocated for model training, with the following day excluded to avoid overlap with the continuous weather system and another day reserved for validation, to evaluate the model's generalization to unseen data during training, or for testing. The trained model is validated against the test data set of ERA5 and IAGOS, which is used for comparative analysis in Sect. 2.4. To test the model's predictions, the results were transformed back to their original scale by applying the inverse of the normalization using the previously saved scaler.

Details on the preparation of training and validation data, particularly for specific humidity  $q$ , are provided in Sect. S3. Similarly, an ANN for  $q$  is implemented, with 300 neurons in each hidden layer and RHi replaced by  $q$  everywhere in both input and output layers (refer to the Supplement for more information). The ANN model can then be applied to ERA5 data for humidity correction in the UTLS region. The computational time required for each scene in Fig. 1 is approx. 5 s on a standard laptop (Intel i5-8250U CPU, 8G memory). This technique incorporates thermodynamic and dynamical meteorological values to account for the vertical and horizontal transport of water vapor and its temporal evolution and takes advantage of numerous humidity measurements.


### 3.3 Importance of the individual variables for the quality of ANN Rhi prediction

The ANN model is interpreted with an investigation of the relative contributions of input variables to the predicted  $RHi_{ANN}$ .  $K_x$  is the relative change in loss when one input, i.e., one feature of ERA5, is set to its mean value for the complete input data set, but the rest of the input features are kept unchanged:

$$K_x = \frac{L_x - L_0}{L_0}, \quad (1)$$

where  $L_x$  is the loss (MSE) for the test data set compared with IAGOS when setting one ERA5 feature input to its average value, and  $L_0$  is the loss for the full test data set. Low values of  $K_x$  indicate a small impact of the change in input quantity on the output accuracy or vice versa.


In Fig. 3, the importance ( $K_x$ ) analysis for all input variables (the current time and level,  $-2$  or  $-6$  h and  $-2$ ,  $-1$ ,  $+1$ ,  $+2$  levels above/below) in the ANN model reveals that  $RHi_{ERA5}$ ,  $T_{ERA5}$ , and  $z$  hold the highest level of significance and carry considerable weight among all parameters. The particular relevance of these three variables can be explained



**Figure 3.** Relative importance of the individual variables to the ANN model for predicting RHi.

by the inherent relationship wherein RHi typically rises in regions with decreasing temperature at higher geopotential  $z$  in the troposphere. Water vapor is carried to upper altitudes with the accompanying adiabatic cooling, which increases RHi. Thus, the ANN model already captures RHi<sub>AGOS</sub> effectively using RHi<sub>ERA5</sub>,  $T_{ERA5}$ , and  $z$ . However, although the other (dynamical) variables are less important for the prediction of RHi<sub>ANN</sub>, they provide a moderate and non-negligible contribution to the accuracy of the RHi prediction model. In fact,  $w$  and  $d$  show  $K_x$  values of 0.29 and 0.26, while  $K_x$  values for  $u$ ,  $v$ , and  $vo$  are even higher, at 0.98, 0.75, and 0.74. There is generally less importance in the contributions of the variables representing dynamical quantities, aligning with the findings in Hofer et al. (2024) based on meteorological variables from the given time.

The fact that dynamical variables, for instance  $u$ ,  $v$ , and  $vo$  in particular, are nearly as important as RHi<sub>ERA5</sub>,  $T_{ERA5}$ , and  $z$  for the description of the physical processes that lead to the decrease/increase in relative humidity in Sect. S2 but at the same time generally show only a moderate importance in the ANN model could be attributed to the correlation with other variables and the significant overlap between the conditional distributions of RHi<sub>ERA5</sub> on ice supersaturation determined by RHi<sub>AGOS</sub> or not (Hofer et al., 2024). Hofer et al. (2024) show that RHi<sub>ERA5</sub> is the most influential predictor for humidity predictions, while the explanatory power of dynamical proxies is insufficient when only using data from the current time and level. However, our updated analysis confirms that incorporating a broader vertical region and the historical time into the dynamical variables has a more significant impact on the ANN model and contributes to the understanding of humidity evolution.




**Figure 4.** Distribution of (a) RHi<sub>ERA5</sub> and (b) RHi<sub>ANN</sub> versus RHi<sub>AGOS</sub> in the UT in all-sky (clear and cloudy) conditions in the test data set. The number of data sets ( $N$ ), the mean absolute error (MAE), and the coefficient of determination ( $R^2$ ) are shown in the panels.

## 4 Model evaluation and results

### 4.1 Validation of ANN RHi in clear and cloudy conditions in the UTLS

This study aims to use the ANN model to resolve biases inherent in NWP model output evaluated in Sect. 2.4. To quantify the accuracy of the ANN model, RHi<sub>ANN</sub> (and  $q_{ANN}$  in the Supplement) is evaluated based on the test data set under four conditions: all-sky UT, cloudy UTLS, clear-sky UTLS, and all-sky LS. Validation results of RHi<sub>ANN</sub> are shown in Figs. 4, 5, and 6 and  $q_{ANN}$  in Sect. S4.

In the UT all-sky condition, important for cirrus clouds and contrails, a high number of measurements, comprising 21 031 data points, are used for the inter-comparison between ERA5 and the outputs of the ANN model. RHi<sub>ANN</sub> (Fig. 4b) demonstrates better agreement with RHi<sub>AGOS</sub> compared to RHi<sub>ERA5</sub> (Fig. 4a). In particular, RHi<sub>ANN</sub> shows values consistent with RHi<sub>AGOS</sub> at RHi  $> 100\%$ , which is a major improvement in comparison to the ERA5 data set. RHi<sub>ANN</sub> exhibits a significant higher correlation with RHi<sub>AGOS</sub> for all uncertainty parameters (mean absolute error, MAE; coefficient of determination,  $R^2$ ) compared to ERA5. The MAE decreases significantly from 15.82 % (ERA5) to 5.71 % (ANN), the  $R^2$  values increase from 0.66 (ERA5) to 0.95 (ANN), and the root mean square error (RMSE) decreases from 20.52 % (ERA5) to 7.88 % (ANN). Notably, the ANN model also effectively corrects the existing peak at RHi<sub>ERA5</sub> = 100 % in Fig. 4a and does not show a peak at RHi  $\sim 100\%$ , similar to the IAGOS measurements. Hence, the ANN exhibits a significant improvement in RHi that would be beneficial for cirrus and cloud predictions. For other scenarios, such as the cloudy UTLS, the clear-sky UTLS, and the all-sky LS between 400 and 200 hPa, the comparison of RHi is shown in Fig. 5. Notably, also in the cloudy UTLS, RHi<sub>ANN</sub> results (Fig. 5b) exhibit a closer correlation with RHi<sub>AGOS</sub> than those in the cloudy region of RHi<sub>ERA5</sub> (Fig. 5a). In the cloudy (Fig. 5a–b) and clear-sky (Fig. 5c–d) conditions in the UTLS, the MAE of the RHi decreases from 16.28 % and 11.21 % to 5.95 % and 4.28 %,




**Figure 5.** Comparison of RHieRA5 (left column) and RHieANN (right column) against RHieAGOS in the (a, b) cloudy UTLS, (c, d) clear-sky UTLS, and (e, f) clear-sky and cloudy (or all-sky) LS regions in the test data set. The number of data sets ( $N$ ), the mean absolute error (MAE), and the coefficient of determination ( $R^2$ ) are indicated in the individual panels.

respectively. Moreover,  $R^2$  increases by 0.30 (0.23) to 0.95 (0.95) for the two scenarios. Again, the peak at 100 % in the RHieRA5 distribution in the cloudy UTLS disappears in RHieANN, in line with the IAGOS observations. In particular, the ANN model shows very good performance in the cloudy UTLS, and RHieANN and RHieAGOS align close to the 1 : 1 line. While the majority of data in the cloudy UTLS is allocated at  $RH_i > 80\%$ , some clouds were observed in ice-subsaturated conditions, or ice particles had sedimented into drier air.

The ANN model also has strong skills of RHi correction in the LS; see Fig. 5e and f.  $R^2$  values increase from 0.59 (ERA5) to 0.95 (ANN), similar to the UT region in Fig. 4. The improvement in RHi prediction by the ANN model is also documented by the decrease in MAE by 6.07 %. The ANN model successfully learns interconnections within the data, as evidenced by its more accurate RHieANN.

Figure 6 presents a detailed relative comparison (mean bias error, MBE) of either RHieRA5 or RHieANN as a function of RHieAGOS. In Fig. 6a, the occurrences of RHieRA5  $> 105\%$  are underestimated compared to the distribution of RHieAGOS. In Fig. 6b, the distribution of RHieANN closely re-



**Figure 6.** Frequency distribution (a and c) and overall mean biased error (MBE) (%) (b and d) of RHieRA5 and RHieANN against RHieAGOS in the clear and cloudy UT (gray) in the test data set.

sembles RHieAGOS, showing a smoother distribution around RHi of 100 %. In Fig. 6c, RHieRA5 shows an increasing dry bias in the UT, reaching 37 % at RHi  $> 120\%$ . The few data points at RHi  $> 120\%$  even exhibit larger deviations by more than 60 % within ERA5. As opposed to this, Fig. 6d shows that RHieANN and RHieAGOS have closer agreement, with an MBE of about  $\pm 11\%$  for all UT measurements up to 140 %. The RHi between 80 % and 130 % in the important range for cirrus clouds is represented well by the ANN with an MBE better than  $\pm 7\%$ . This suggests that the saturated region of RHi, which presents a requisite environmental condition for new ice crystal nucleation and subsequent growth, can be more accurately parameterized.

Wolf et al. (2025) developed a humidity correction technique for RHieRA5 using IAGOS measurements through a multivariate quantile method. The differences between the corrected ERA5 data and RHieRA5 in cloudy regions are documented in their Table 3, with the mean absolute difference ranging from  $-2.2\%$  to  $12.08\%$  depending on cloud fraction. Although not directly comparable within the same time frame, RHieANN also shows good performance, with a MAE of approximately 5.8 % in the same region of all-sky UT and cloudy UTLS.

RHieANN also shows better agreement with independent airborne measurements compared to ERA5 data. For detailed information on the humidity data on 21 July 2021 during the CIRRUS-HL campaign, refer to Sect. S5, which includes measured data from an atmospheric ionization mass spectrometer (AIMS; Kaufmann et al., 2016) instrument and RHieRA5 and RHieANN.

The investigations related to validating ANN $q$  in clear and cloudy conditions in the UTLS are shown in Sect. S4.

#### 4.2 Skill of ANN and ERA5 in predicting RH<sub>i</sub> > 100 % versus IAGOS data

For cirrus clouds and contrails, accurately representing RH<sub>i</sub> > 100 %, and thus ice supersaturation, is of great importance. Hence, here we focus on the data sets with RH<sub>i</sub> > 100 %. The skill of ice supersaturation prediction from RH<sub>iERA5</sub> and RH<sub>iANN</sub> is evaluated based on the equitable threat score (ETS), as described in Gierens et al. (2020). The ETS measures forecasting performance by assessing the proportion of correctly forecasted events and is often used in weather forecast verification (Wang, 2014). First, events are labeled according to the contingency table, with a ( $Y_{\text{IAGOS}}/Y_{\text{ERA5}}$  or  $Y_{\text{IAGOS}}/Y_{\text{ANN}}$ , ice supersaturation predicted and observed), b ( $Y_{\text{IAGOS}}/N_{\text{ERA5}}$  or  $Y_{\text{IAGOS}}/N_{\text{ANN}}$ , no ice supersaturation predicted but observed), c ( $N_{\text{IAGOS}}/Y_{\text{ERA5}}$  or  $N_{\text{IAGOS}}/Y_{\text{ANN}}$ , ice supersaturation predicted but not observed), and d ( $N_{\text{IAGOS}}/N_{\text{ERA5}}$  or  $N_{\text{IAGOS}}/N_{\text{ANN}}$  ice supersaturation neither predicted nor observed).  $Y_{\text{IAGOS}}$  indicates that the waypoint is in ice supersaturation based on the IAGOS measurements, while  $N_{\text{IAGOS}}$  indicates the absence of ice supersaturation. The same notations are applied when analyzing the statistics for ERA5 and ANN. The ETS is then calculated using the following equations:

$$\text{ETS} = \frac{a - r}{a + b + c - r} \quad (2)$$

with

$$r = \frac{(a + b)(a + c)}{a + b + c + d}. \quad (3)$$

The ETS value gets larger when the ice supersaturation prediction is closer to the measured values (here IAGOS data). ETS = 1 indicates that all RH<sub>iIAGOS</sub> values perfectly agree with RH<sub>iERA5</sub> or RH<sub>iANN</sub>. ETS = 0 means a completely random distribution, while negative ETS implies a negative correlation.

Table 2 shows ETS values for both the ERA5- and the ANN-predicted ice supersaturation across all test data sets in Sect. 2.4. The scores for ERA5 in all-sky UT, cloudy UTLS, and all-sky LS classes are 0.23, 0.21, and 0.14, respectively, indicating limited predictive skill, particularly in the all-sky LS region. In contrast, the ANN model significantly enhances the ice supersaturation prediction, yielding scores of 0.71, 0.70, and 0.52 for the respective regions. This represents an approximate 0.44 increase in ETS across all classes, thereby facilitating related studies on the formation and persistence of cirrus clouds. The clear-sky UTLS region is not discussed here, as we are focusing on RH<sub>i</sub> > 100 %. According to Fig. 5c, few ERA5 data points fall within the ice supersaturation region in the clear-sky data sets.


Teoh et al. (2022) developed a statistical approach to correct the ERA5 humidity fields, in particular for ice supersaturation, with the aim of adjusting the PDF of RH<sub>iERA5</sub> in

order to achieve a similar PDF to RH<sub>iIAGOS</sub>. After applying this humidity correction method, the ETS for ice supersaturation in the all-sky UTLS reached a value of 0.424 when compared to IAGOS measurements in 2019, as shown in their Table S4. This statistical method outperforms RH<sub>iERA5</sub> in predicting ice supersaturation, as the latter achieves an ETS of approximately 0.2. In a recent study, Hofer et al. (2024) use RH<sub>iERA5</sub>,  $T_{\text{ERA5}}$ , and dynamical proxies only from the given time in several regression models to predict RH<sub>i</sub> > 100 % and find the best regression with an ETS of 0.378 for 16 588 flights of the Measurement of Ozone and Water Vapour on Airbus In-service Aircraft (MOZAIC) between 2000 and 2009. While not directly comparable within the same time frame, RH<sub>iANN</sub> excels in forecasting ice supersaturation relative to ERA5 and methods from Teoh et al. (2022) and Hofer et al. (2024), demonstrating a higher accuracy with an ETS as high as 0.71.

#### 5 CoCiP predictions and MSG contrail cirrus observations

As an application, this section investigates the impact of improved humidity prediction in the UT on the estimation of contrail cirrus optical thickness using CoCiP simulations (Sect. 2.5) and compares the results with retrieved IOT from MSG observations using the CiPS algorithm (Sect. 2.6). The selected case is from 10:00 UTC on 14 April 2021 (during the ECLIF3 campaign) over the North Atlantic region (NAR; Sect. 2.1), representing a typical contrail cirrus situation just off the coast of Ireland. The MSG observation scene is from 09:45 UTC, as SEVIRI scans from the south, taking about 12 min per scene and reaching the upper edge (near the NAR) around 09:56 UTC, which is more consistent with the CoCiP simulation time. For the CoCiP simulations, specific humidity  $q_{\text{IFS}}$  and other atmospheric trace gas profiles from the ECMWF IFS model-level data between level 73 (about 190 hPa) and level 90 (about 400 or 410 hPa) are used as input to the ANN model to produce  $q_{\text{ANN}}$  profiles. We perform two CoCiP experiments: the  $q_{\text{IFS}}$  and the resulting  $q_{\text{ANN}}$  profiles serve as input for CoCiP, while other parameters, particularly cloud liquid water content and ciwc, are kept constant using IFS values to include the same natural cloud effects in both CoCiP simulations.

Figure 7 provides a comparison of the spatial distributions and frequency of occurrence (histogram) of CoCiP-simulated and MSG-observed IOT. In the MSG scene (Fig. 7a), contrails and contrail cirrus, represented by linear structures with IOT  $\sim 0.3\text{--}0.5$  (Wang et al., 2023), are situated above the Atlantic Ocean, extending from west to east and surrounded by thicker cirrus clouds with higher IOT (even exceeding 1.0). The simulated IOTs (Fig. 7b and c) show patterns of higher IOT surrounded by lower IOT cirrus, as in the MSG observations, although the single structures are not directly comparable. In addition, due to satellite detection limitations, CiPS



**Figure 7.** Distributions of IOT for contrails and cirrus retrieved from (a) MSG observations using the CiPS algorithm and simulated using the CoCiP model with (b)  $q_{\text{ANN}}$  or (c)  $q_{\text{IFS}}$  at 10:00 UTC on 14 April 2021. The IOT distribution for contrails from CoCiP simulations is shown in (e) and (f). The linear structures of contrails correspond to the associated flight tracks. The IOT frequencies (histograms) for contrail cirrus and contrails are shown in (d) and (g), respectively.

**Table 2.** ETS values for the prediction of  $\text{RHi} > 100\%$  from  $\text{RHi}_{\text{ERA5}}$  and  $\text{RHi}_{\text{ANN}}$  in the test data set between 200 and 400 hPa over the Atlantic, Europe, and Africa in 2020.

Scenarios	$Y_{\text{IAGOS}}/Y_{\text{ERA5}}$	$Y_{\text{IAGOS}}/N_{\text{ERA5}}$	$N_{\text{IAGOS}}/Y_{\text{ERA5}}$	$N_{\text{IAGOS}}/N_{\text{ERA5}}$	ETS
Clear and cloudy UT	66.34 %	3.31 %	19.42 %	10.94 %	0.23
Cloudy UTLS	54.61 %	4.99 %	24.06 %	16.34 %	0.21
Clear and cloudy LS	97.48 %	0.21 %	1.96 %	0.36 %	0.14
Clear-sky UTLS	95.43 %	0.05 %	4.23 %	0.28 %	0.06
Scenarios	$Y_{\text{IAGOS}}/Y_{\text{ANN}}$	$Y_{\text{IAGOS}}/N_{\text{ANN}}$	$N_{\text{IAGOS}}/Y_{\text{ANN}}$	$N_{\text{IAGOS}}/N_{\text{ANN}}$	ETS
Clear and cloudy UT	67.07 %	2.57 %	4.57 %	25.79 %	0.71
Cloudy UTLS	56.26 %	3.34 %	4.95 %	35.44 %	0.70
Clear and cloudy LS	97.29 %	0.40 %	0.89 %	1.42 %	0.52
Clear-sky UTLS	94.99 %	0.49 %	2.07 %	2.44 %	0.47

cannot capture the thinnest ice clouds. For these reasons, a quantitative pixel-to-pixel comparison between the CoCiP simulations and the MSG observation is not meaningful in this context, and we thus rather consider frequency distributions of IOT in the following. In general, the simulated IOT using  $q_{\text{ANN}}$  (Fig. 7b) is closer to CiPS-retrieved IOT compared to the CoCiP simulation with  $q_{\text{IFS}}$  (Fig. 7c). The histogram (Fig. 7d) shows decreasing frequencies of occurrence with increasing IOT. Better agreement is also exhibited between IOT from CiPS and the CoCiP simulation using  $q_{\text{ANN}}$  than that using  $q_{\text{IFS}}$ . For natural cirrus, larger IOT ( $\sim 0.75$ ) is observed, while the smaller IOT ( $< 0.5$ ) for contrails is of particular interest. In the lower panels showing only con-

trails, the simulation with increased humidity exhibits larger IOT (Fig. 7e) compared to that without humidity correction (Fig. 7f). Higher IOT up to 0.85 from CoCiP with  $q_{\text{ANN}}$ , compared to IOT values below 0.5 with  $q_{\text{IFS}}$ , is due to the growth of contrail ice crystals from the increased amount of available water vapor in  $q_{\text{ANN}}$  and is also evident in the frequency analysis (Fig. 7g).

In general, the model results demonstrate increased consistency with MSG observations when  $q_{\text{ANN}}$  is incorporated in CoCiP. Given that ice supersaturation typically exhibits large horizontal but shallow vertical extensions (Spichtinger et al., 2003), a minor adjustment in cruising altitude, avoiding regions of high humidity, can potentially reduce contrail

radiative forcing. An improved representation of humidity thanks to an ANN approach is thus crucial for more accurate predictions of the contrail cirrus cover and radiative effect (Kaufmann et al., 2024).

## 6 Summary and conclusions

The distribution of relative humidity in the UTLS from NWP models, which plays a vital role in the parameterizations of natural cirrus and contrail cirrus properties, is subject to large uncertainties. In this study we propose a humidity bias correction method for relative humidity from ERA5,  $RH_{ERA5}$ , particularly for ice supersaturation in the UT, using an ANN technique. The novelty of this study lies in the incorporation of thermodynamic and dynamical atmospheric quantities for the given time and height together with atmospheric properties from previous times and nearby altitudes. The atmospheric humidity improvement method consists of an ANN developed using atmospheric variables from ERA5, along with collocated measurements of water vapor from IAGOS. The ERA5 data include temporal and vertical dependencies of humidity on meteorological conditions, combining not only the historic data ( $-6, -2$  h) and current time but also  $\pm 2$  ERA5 pressure layers around the flight latitude of IAGOS. The target region covers the Atlantic, Europe, and Africa, spanning  $0$  to  $80^\circ$  N and  $30^\circ$  W to  $50^\circ$  E and pressure levels from  $400$  to  $200$  hPa.

The analysis of biases between collocated  $RH_{ERA5}$  and  $RH_{IAGOS}$  reveals an underestimation of  $RH_{ERA5}$  within the UT and an  $RH_i$  occurrence peak near 100 % due to the cloud saturation adjustment by ECMWF NWP. The ERA5–IAGOS collocated data are processed and the variables for training the ANN model for humidity correction are selected based on the discussion of the temporal evolution of meteorological variables. Humidity, temperature, and geopotential (as a variable for the altitude) have the biggest impact on the  $RH_{ANN}$  results, while other meteorological variables, including horizontal wind speed, relative vorticity, vertical velocity, and divergence, have a high or moderate to minor but measurable influence.

Using this ANN humidity correction, the MAE of  $RH_{ANN}$  compared to  $RH_{IAGOS}$ , relative to  $RH_{ERA5}$  and  $RH_{IAGOS}$ , is reduced from 15.82 % to 5.71 % in all-sky UT, 16.28 % to 5.95 % in cloudy UTLS, 11.21 % to 4.28 % in clear UTLS, and 9.78 % to 3.71 % in all-sky LS regions, respectively, presenting remarkable improvements, particularly in the all-sky UT and cloudy UTLS regions. A previously existing occurrence peak at  $RH_i = 100$  % in  $RH_{ERA5}$ , which is caused by the cloud saturation adjustment in NWP, has been removed completely by the ANN.

The representation of ice supersaturation in  $RH_{ERA5}$  and  $RH_{ANN}$  with respect to  $RH_{IAGOS}$  was assessed with the calculation of the ETS value. The dynamic-based humidity correction leads to an increase in ETS from 0.23, 0.21, and 0.14

(ERA5) to 0.71, 0.70, and 0.52 by the ANN, respectively, in the all-sky UT, cloudy UTLS, and all-sky LS regions. The skill of ice supersaturation prediction improves considerably.

The forecast of optical and radiative properties of cirrus and contrail cirrus, based on the ANN humidity correction, is exemplarily assessed with CoCiP simulations using IFS weather data and the ANN-corrected data and MSG satellite observations for one case between  $35$  and  $60^\circ$  N (over the NAR) at 10:00 UTC on 14 April 2021. The result shows better agreement in ice optical thickness between model simulations with humidity correction and satellite observations in this contrail situation.

Teoh et al. (2022) and Wolf et al. (2025) utilize IA-GOS measurements to correct  $RH_{ERA5}$  with statistical methods. Our study shows the potential of the emerging field of machine-learning-based weather prediction post-processing, in which forecast outputs are improved using historical observations and analysis data. How the current atmospheric states influence the future development of humidity patterns has been highlighted. One issue in the existing model data, where the frequency and degree of ice supersaturation in the UT are consistently underestimated due to the practice of cloud saturation adjustment, has been successfully addressed by the ANN model. The method demonstrates competitive performance, as seen by the decreased MAE and larger ETS compared to the accuracy of the aforementioned statistical methods.

Incorporating more water vapor data from the fleet-wide observations of humidity within the UTLS can further improve this method. Our findings suggest potential applications for aircraft diversion strategies to avoid ice supersaturation regions and reduce contrail cirrus climate impact. Further research on applying humidity correction methods to weather forecasts is vital for improving our understanding of the global cloud radiation budget. Our improved humidity predictions could serve as benchmarks for the further measurements of aircraft campaigns, as the assimilation or reference data set for NWP or climate models for a better parameterization of ice supersaturation. The method could also be applied to other weather forecast models, including those from ECMWF and national weather services. Additionally, increased resolution of NWP models at the tropopause is required for better cirrus and contrail forecasting. Combining modeled meteorological conditions and their temporal changes, along with measured humidity from long-term data sets, is essential for a more realistic representation of  $RH_i$ , subsequent processes like cloud and contrail formation in the UTLS, and their climate impact.

**Data availability.** IAGOS measurements are available at <https://doi.org/10.25326/20> (Boulanger et al., 2020). The ERA5 and IFS atmospheric profiles can be obtained from the Climate Data Store (<https://doi.org/10.24381/cds.bd0915c6>, Hersbach et al., 2023) or ECMWF directly. The SEVIRI data are provided

by the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT). The CoCiP model code can be accessed from <https://doi.org/10.5281/zenodo.7776686> (Shapiro et al., 2023). The machine learning technique implementation is based on the open-source platform TensorFlow (<https://arxiv.org/abs/1603.04467>, Abadi et al., 2016). The required software packages are Python (van Rossum and Drake, 2009), Keras (Gulli and Pal, 2017), and Scikit-learn (Pedregosa et al., 2011).

**Supplement.** The supplement related to this article is available online at <https://doi.org/10.5194/acp-25-2845-2025-supplement>.

**Author contributions.** ZW conceived the study concept, developed the methods, and wrote the paper. LB and CV advised on the study and provided feedback on the paper. KG contributed to explanations of the dependence of humidity on meteorological conditions. MIH contributed to the initial method and ERA5 data evaluation. SR and AP coordinated the IAGOS research infrastructure and provided the data. SK helped with the interpretation of humidity bias in ERA5. All authors contributed to and commented on the paper.

**Competing interests.** At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

**Acknowledgements.** We thank the IAGOS European research infrastructure for excellent global aircraft measurements and ECMWF and EUMETSAT for providing the modeled atmospheric data and MSG/SEVIRI observations. ZW was supported by the DLR/DAAD (Deutsches Zentrum für Luft- und Raumfahrt/Deutscher Akademischer Austauschdienst) Research Fellowships – Doctoral Studies in Germany, 2020, under grant no. 57540125 and now by the LuFo project MEFKON. CV, LB, and SK are supported by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) within HALO SPP 1294 under project nos. VO1504/9-1 and 522359172 and TRR 301 – project ID 428312742, the European Union CONCERTO, and SESAR JU CICONIA. This research is also supported by the European Union A4Climate. IAGOS data were created with support from the European Commission; national agencies in Germany (BMBF), France (MESR), and the UK (NERC); and the IAGOS member institutions (<https://www.iagos.org/iagos-partner/>, last access: 2 March 2025). The participating airlines (in the year 2020: Deutsche Lufthansa, Air France, China Airlines, Hawaiian Airlines) supported IAGOS by

carrying the measurement equipment free of charge. The data are available at <http://www.iagos.fr> (last access: 2 March 2025) thanks to additional support from AERIS. We thank Andreas Schäfer from DLR for valuable discussions.

**Financial support.** This research has been supported by the Deutscher Akademischer Austauschdienst (DLR/DAAD) Research Fellowships – Doctoral Studies in Germany, 2020, grant no. 57540125), the Bundesministerium für Wirtschaft und Klimaschutz (LuFo, MEFKON grant no. 20F2202B), the Deutsche Forschungsgemeinschaft (grant nos. SPP-1294 HALO, VO1504/9-1, 522359172, and TRR 301, 428312742), the European Commission, EU Horizon (A4Climate grant no. 101192301, CONCERTO grant no. 101101999), and the SESAR Joint Undertaking (CICONIA grant no. 101114613).

The article processing charges for this open-access publication were covered by the German Aerospace Center (DLR).

**Review statement.** This paper was edited by Annika Oertel and reviewed by two anonymous referees.

## References

Abadi, M., Agarwal, A., Barham, P., Brevdo, E., Chen, Z., Citro, C., Corrado, G. S., Davis, A., Dean, J., Devin, M., and Ghemawat, S.: Tensorflow: Large-scale machine learning on heterogeneous distributed systems, arXiv preprint, Cornell University [data set], <https://arxiv.org/abs/1603.04467> (last access: 1 March 2025), 2016.

Alduchov, O. A. and Eskridge, R. E.: Improved Magnus form approximation of saturation vapor pressure, *J. Climatol. Appl. Meteorol.*, 35, 601–609, [https://doi.org/10.1175/1520-0450\(1996\)035<0601:imfaos>2.0.co;2](https://doi.org/10.1175/1520-0450(1996)035<0601:imfaos>2.0.co;2), 1996.

Berkes, F., Neis, P., Schultz, M. G., Bundke, U., Rohs, S., Smit, H. G. J., Wahner, A., Konopka, P., Boulanger, D., Nédélec, P., Thouret, V., and Petzold, A.: In situ temperature measurements in the upper troposphere and lowermost stratosphere from 2 decades of IAGOS long-term routine observation, *Atmos. Chem. Phys.*, 17, 12495–12508, <https://doi.org/10.5194/acp-17-12495-2017>, 2017.

Bickel, M., Ponater, M., Bock, L., Burkhardt, U., and Reineke, S.: Estimating the effective radiative forcing of contrail cirrus, *J. Climate*, 33, 1991–2005, <https://doi.org/10.1175/JCLI-D-19-0467.1>, 2020.

Boulanger, D., Thouret, V., and Petzold, A.: IAGOS Data Portal, AERIS [data set], <https://doi.org/10.25326/20>, 2020.

Burkhardt, U., Bock, L., and Bier, A.: Mitigating the contrail cirrus climate impact by reducing aircraft soot number emissions, *npj Clim. Atmos. Sci.*, 1, 37, <https://doi.org/10.1038/s41612-018-0046-4>, 2018.

Chen, C.-C., Gettelman, A., Craig, C., Minnis, P., and Duda, D. P.: Global contrail coverage simulated by CAM5 with the inventory of 2006 global aircraft emissions, *J. Adv. Model. Earth Syst.*, 4, M04003, <https://doi.org/10.1029/2011MS000105>, 2012.

Dessler, A. E. and Sherwood, S. C.: ATMOSPHERIC SCIENCE: A Matter of Humidity, *Science*, 323, 1020–1021, <https://doi.org/10.1126/science.1171264>, 2009.

Diao, M., Jensen, J. B., Pan, L. L., Homeyer, C. R., Honomichl, S., Bresch, J. F., and Bansemer, A.: Distributions of ice supersaturation and ice crystals from airborne observations in relation to upper tropospheric dynamical boundaries, *J. Geophys. Res.-Atmos.*, 120, 5101–5121, <https://doi.org/10.1002/2015JD023139>, 2015.

Dickson, N. C., Gierens, K. M., Rogers, H. L., and Jones, R. L.: Probabilistic description of ice-supersaturated layers in low resolution profiles of relative humidity, *Atmos. Chem. Phys.*, 10, 6749–6763, <https://doi.org/10.5194/acp-10-6749-2010>, 2010.

Dyroff, C., Zahn, A., Christner, E., Forbes, R., Tompkins, A. M., and van Velthoven, P. F. J.: Comparison of ECMWF analysis and forecast humidity data with CARIBIC upper troposphere and lower stratosphere observations, *Q. J. Roy. Meteor. Soc.*, 141, 833–844, <https://doi.org/10.1002/qj.2400>, 2015.

ECMWF: IFS Documentation CY41R2 – Part IV: Physical Processes, IFS Documentation CY41R2, 4, <https://doi.org/10.21957/tr5rv27xu>, 2016.

Forbes, R. M. and Ahlgrimm, M.: On the representation of high-latitude boundary layer mixed-phase cloud in the ECMWF global model, *Mon. Weather Rev.*, 142, 3425–3445, <https://doi.org/10.1175/MWR-D-13-00325.1>, 2014.

Forbes, R. and Tompkins, A.: An improved representation of cloud and precipitation, Tech. Rep., European Center for Medium-Range Weather Forecasting, <https://doi.org/10.21957/nfgulzhe>, 2011.

Forster, P. M. de F. and Shine, K. P.: Assessing the climate impact of trends in stratospheric water vapor, *Geophys. Res. Lett.*, 29, 10–1–10–4, <https://doi.org/10.1029/2001GL013909>, 2002.

Gasparini, B., McGraw, Z., Storelvmo, T., and Lohmann, U.: To what extent can cirrus cloud seeding counteract global warming?, *Environ. Res. Lett.*, 15, 054002, <https://doi.org/10.1088/1748-9326/ab71a3>, 2020.

Gettelman, A., Hoor, P., Pan, L. L., Randel, W. J., Hegglin, M. I., and Birner, T.: THE EXTRATROPICAL UPPER TROPOSPHERE AND LOWER STRATOSPHERE, *Rev. Geophys.*, 49, 3, <https://doi.org/10.1029/2011RG000355>, 2011.

Gierens, K. and Brinkop, S.: Dynamical characteristics of ice supersaturated regions, *Atmos. Chem. Phys.*, 12, 11933–11942, <https://doi.org/10.5194/acp-12-11933-2012>, 2012.

Gierens, K., Matthes, S., and Rohs, S.: How Well Can Persistent Contrails Be Predicted?, *Aerospace*, 7, 169, <https://doi.org/10.3390/aerospace7120169>, 2020.

Groß, S., Wirth, M., Schäfler, A., Fix, A., Kaufmann, S., and Voigt, C.: Potential of airborne lidar measurements for cirrus cloud studies, *Atmos. Meas. Tech.*, 7, 2745–2755, <https://doi.org/10.5194/amt-7-2745-2014>, 2014.

Gulli, A. and Pal, S.: Deep learning with Keras, Packt Publishing, ISBN 9781787128422, 2017.

He, K., Zhang, X., Ren, S., and Sun, J.: Delving Deep into Rectifiers: Surpassing Human-Level Performance on ImageNet Classification, 2015 IEEE International Conference on Computer Vision (ICCV), Santiago, Chile, 1026–1034, <https://doi.org/10.1109/ICCV.2015.123>, 2015.

Hegglin, M. I., Tegtmeier, S., Anderson, J., Froidevaux, L., Fuller, R., Funke, B., Jones, A., Lingenfelter, G., Lumpe, J., Pendlebury, D., Remsberg, E., Rozanov, A., Toohey, M., Urban, J., von Clar mann, T., Walker, K. A., Wang, R., and Weigel, K.: SPARC data initiative: Comparison of water vapor climatologies from international satellite limb sounders: Sparc Data Initiative Water Vapor Comparisons, *J. Geophys. Res.-Atmos.*, 118, 11824–11846, <https://doi.org/10.1002/jgrd.50752>, 2013.

Hegglin, M. I., Plummer, D. A., Shepherd, T. G., Scinocca, J. F., Anderson, J., Froidevaux, L., Funke, B., Hurst, D., Rozanov, A., Urban, J., von Clar mann, T., Walker, K. A., Wang, H. J., Tegtmeier, S., and Weigel, K.: Vertical structure of stratospheric water vapour trends derived from merged satellite data, *Nat. Geosci.*, 7, 768–776, <https://doi.org/10.1038/NGEO2236>, 2014.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Q. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: Copernicus Climate Change Service (C3S) Climate Data Store (CDS), ERA5 hourly data on pressure levels from 1940 to present, ECMWF [data set], <https://doi.org/10.24381/cds.bd0915c6>, 2023.

Heymsfield, A. J., Miloshevich, L. M., Twohy, C., Sachse, G., and Oltmans, S.: Upper-tropospheric relative humidity observations and implications for cirrus ice nucleation, *Geophys. Res. Lett.*, 25, 1343–1346, <https://doi.org/10.1029/98GL01089>, 1998.

Hofer, S., Gierens, K., and Rohs, S.: How well can persistent contrails be predicted? An update, *Atmos. Chem. Phys.*, 24, 7911–7925, <https://doi.org/10.5194/acp-24-7911-2024>, 2024.

Kadow, C., Hall, D. M., and Ulbrich, U.: Artificial intelligence reconstructs missing climate information, *Nat. Geosci.*, 13, 408–413, <https://doi.org/10.1038/s41561-020-0582-5>, 2020.

Kärcher, B.: Formation and radiative forcing of contrail cirrus, *Nat. Commun.*, 9, 1824, <https://doi.org/10.1038/s41467-018-04068-0>, 2018.

Kaufmann, S., Voigt, C., Jurkat, T., Thornberry, T., Fahey, D. W., Gao, R.-S., Schlage, R., Schäuble, D., and Zöger, M.: The airborne mass spectrometer AIMS – Part 1: AIMS-H<sub>2</sub>O for UTLS water vapor measurements, *Atmos. Meas. Tech.*, 9, 939–953, <https://doi.org/10.5194/amt-9-939-2016>, 2016.

Kaufmann, S., Voigt, C., Heller, R., Jurkat-Witschas, T., Krämer, M., Rolf, C., Zöger, M., Giez, A., Buchholz, B., Ebert, V., Thornberry, T., and Schumann, U.: Intercomparison of midlatitude tropospheric and lower-stratospheric water vapor measurements and comparison to ECMWF humidity data, *Atmos. Chem. Phys.*, 18, 16729–16745, <https://doi.org/10.5194/acp-18-16729-2018>, 2018.

Kaufmann, S., Dischl, R., and Voigt, C.: Regional and seasonal impact of hydrogen propulsion systems on potential contrail cirrus cover, *Atmos. Environ.*, 24, 100298, <https://doi.org/10.1016/j.aeaoa.2024.100298>, 2024.

Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrman, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, *Atmos. Chem. Phys.*, 9, 3505–3522, <https://doi.org/10.5194/acp-9-3505-2009>, 2009.

Krüger, K., Schäfler, A., Wirth, M., Weissmann, M., and Craig, G. C.: Vertical structure of the lower-stratospheric moist bias in the ERA5 reanalysis and its connection to mixing processes, *Atmos. Chem. Phys.*, 22, 15559–15577, <https://doi.org/10.5194/acp-22-15559-2022>, 2022.

Kunz, A., Spelten, N., Konopka, P., Müller, R., Forbes, R. M., and Wernli, H.: Comparison of Fast In situ Stratospheric Hygrometer (FISH) measurements of water vapor in the upper troposphere and lower stratosphere (UTLS) with ECMWF (re)analysis data, *Atmos. Chem. Phys.*, 14, 10803–10822, <https://doi.org/10.5194/acp-14-10803-2014>, 2014.

Lam, R., Sanchez-Gonzalez, A., Willson, M., Wirnsberger, P., Fortunato, M., Alet, F., Ravuri, S., Ewalds, T., Eaton-Rosen, Z., Hu, W., Merose, A., Hoyer, S., Holland, G., Vinyals, O., Stott, J., Pritzel, A., Mohamed, S., and Battaglia, P.: Learning skillful medium-range global weather forecasting, *Science*, 382, 1416–1421, <https://doi.org/10.1126/science.adi2336>, 2023.

Lamquin, N., Stubenrauch, C. J., Gierens, K., Burkhardt, U., and Smit, H.: A global climatology of upper-tropospheric ice supersaturation occurrence inferred from the Atmospheric Infrared Sounder calibrated by MOZAIC, *Atmos. Chem. Phys.*, 12, 381–405, <https://doi.org/10.5194/acp-12-381-2012>, 2012.

Lawrence, H., Bormann, N., Sandu, I., Day, J., Farnan, J., and Bauer, P.: Use and impact of Arctic observations in the ECMWF Numerical Weather Prediction system, *Q. J. Roy. Meteor. Soc.*, 145, 3432–3454, <https://doi.org/10.1002/qj.3628>, 2019.

Ma, R., Letu, H., Yang, K., Wang, T., Shi, C., Xu, J., Shi, J., Shi, C., and Chen, L.: Estimation of surface shortwave radiation from Himawari-8 satellite data based on a combination of radiative transfer and deep neural network, *IEEE T. Geosci. Remote*, 58, 5304–5316, <https://doi.org/10.1109/tgrs.2019.2963262>, 2020.

Märkl, R. S., Voigt, C., Sauer, D., Dischl, R. K., Kaufmann, S., Harlaß, T., Hahn, V., Roiger, A., Weiß-Rehm, C., Burkhardt, U., Schumann, U., Marsing, A., Scheibe, M., Dörnbrack, A., Renard, C., Gauthier, M., Swann, P., Madden, P., Luff, D., Sallinen, R., Schripp, T., and Le Clercq, P.: Powering aircraft with 100 % sustainable aviation fuel reduces ice crystals in contrails, *Atmos. Chem. Phys.*, 24, 3813–3837, <https://doi.org/10.5194/acp-24-3813-2024>, 2024.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, É.: Scikit-learn: Machine Learning in Python, *J. Mach. Learn. Res.*, 12, 2825–2830, ISBN 1532-4435, 2011.

Petzold, A., Thouret, V., Gerbig, C., Zahn, A., Brenninkmeijer, C. A. M., Gallagher, M., Hermann, M., Pontaud, M., Ziereis, H., Boulanger, D., Marshall, J., Nédélec, P., Smit, H. G. J., Frieß, U., Flaud, J.-M., Wahner, A., Cammas, J.-P., Volz-Thomas, A., and IAGOS-Team: Global-Scale Atmosphere Monitoring by In-Service Aircraft – Current Achievements and Future Prospects of the European Research Infrastructure IAGOS, *Tellus B*, 67, 28452, <https://doi.org/10.3402/tellusb.v67.28452>, 2015.

Petzold, A., Neis, P., Rütimann, M., Rohs, S., Berkes, F., Smit, H. G. J., Krämer, M., Spelten, N., Spichtinger, P., Nédélec, P., and Wahner, A.: Ice-supersaturated air masses in the northern mid-latitudes from regular in situ observations by passenger aircraft: vertical distribution, seasonality and tropospheric fingerprint, *Atmos. Chem. Phys.*, 20, 8157–8179, <https://doi.org/10.5194/acp-20-8157-2020>, 2020.

Reutter, P., Neis, P., Rohs, S., and Sauvage, B.: Ice supersaturated regions: properties and validation of ERA-Interim reanalysis with IAGOS in situ water vapour measurements, *Atmos. Chem. Phys.*, 20, 787–804, <https://doi.org/10.5194/acp-20-787-2020>, 2020.

Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, *J. Geophys. Res.*, 117, D16305, <https://doi.org/10.1029/2012JD017751>, 2012.

Rolf, C., Rohs, S., Smit, H. G. J., Krämer, M., Bozóki, Z., Hofmann, S., Franke, H., Maser, R., Hoor, P., and Petzold, A.: Evaluation of compact hygrometers for continuous airborne measurements, *Meteorol. Z.*, 33, 15–34, <https://doi.org/10.1127/metz/2023/1187>, 2023.

Rollins, A. W., Thornberry, T. D., Gao, R. S., Smith, J. B., Sayres, D. S., Sargent, M. R., Schiller, C., Krämer, M., Spelten, N., Hurst, D. F., Jordan, A. F., Hall, E. G., Vömel, H., Diskin, G. S., Podolske, J. R., Christensen, L. E., Rosenlof, K. H., Jensen, E. J., and Fahey, D. W.: Evaluation of UT/LS hygrometer accuracy by intercomparison during the NASA MACPEX mission, *J. Geophys. Res.-Atmos.*, 119, 1915–1935, <https://doi.org/10.1002/2013JD020817>, 2014.

Sanogo, S., Boucher, O., Bellouin, N., Borella, A., Wolf, K., and Rohs, S.: Variability in the properties of the distribution of the relative humidity with respect to ice: implications for contrail formation, *Atmos. Chem. Phys.*, 24, 5495–5511, <https://doi.org/10.5194/acp-24-5495-2024>, 2024.

Schmidt, G. A., Ruedy, R. A., Miller, R. L., and Lacis, A. A.: Attribution of the present-day total greenhouse effect, *J. Geophys. Res.-Atmos.*, 115, D20106, <https://doi.org/10.1029/2010JD014287>, 2010.

Schumann, U.: On conditions for contrail formation from aircraft exhausts, *Meteorol. Z.*, 5, 4–23, <https://doi.org/10.1127/metz/5/1996/4>, 1996.

Schumann, U.: A contrail cirrus prediction model, *Geosci. Model Dev.*, 5, 543–580, <https://doi.org/10.5194/gmd-5-543-2012>, 2012.

Schumann, U., Baumann, R., Baumgardner, D., Bedka, S. T., Duda, D. P., Freudenthaler, V., Gayet, J.-F., Heymsfield, A. J., Minnis, P., Quante, M., Raschke, E., Schlager, H., Vázquez-Navarro, M., Voigt, C., and Wang, Z.: Properties of individual contrails: a compilation of observations and some comparisons, *Atmos. Chem. Phys.*, 17, 403–438, <https://doi.org/10.5194/acp-17-403-2017>, 2017.

Schumann, U., Bugliaro, L., Dörnbrack, A., Baumann, R., and Voigt, C.: Aviation Contrail Cirrus and Radiative Forcing Over Europe During 6 Months of COVID-19, *Geophys. Res. Lett.*, 48, e2021GL092771, <https://doi.org/10.1029/2021GL092771>, 2021a.

Schumann, U., Poll, I., Teoh, R., Koelle, R., Spinielli, E., Molloy, J., Koudis, G. S., Baumann, R., Bugliaro, L., Stettler, M., and

Voigt, C.: Air traffic and contrail changes over Europe during COVID-19: a model study, *Atmos. Chem. Phys.*, 21, 7429–7450, <https://doi.org/10.5194/acp-21-7429-2021>, 2021b.

Seifert, A. and Siewert, C.: An ML-based P3-like multimodal two-moment ice microphysics in the ICON model, *J. Adv. Model. Earth Syst.*, 16, e2023MS004206, <https://doi.org/10.1029/2023MS004206>, 2024.

Shapiro, M., Engberg, Z., Teoh, R., and Dean, T.: pycontrails: Python library for modeling aviation climate impacts, *Zenodo [code]*, <https://doi.org/10.5281/zenodo.7776686>, 2023.

Sonntag, D.: Advancements in the field of hygrometry, *Meteorol. Z.*, 3, 51–66, <https://doi.org/10.1127/metz/3/1994/51>, 1994.

Sperber, D. and Gierens, K.: Towards a more reliable forecast of ice supersaturation: concept of a one-moment ice-cloud scheme that avoids saturation adjustment, *Atmos. Chem. Phys.*, 23, 15609–15627, <https://doi.org/10.5194/acp-23-15609-2023>, 2023.

Spichtinger, P., Gierens, K., and Read, W.: The global distribution of ice-supersaturated regions as seen by the Microwave Limb Sounder, *Quart. J. Roy. Met. Soc.*, 129, 3391–3410, <https://doi.org/10.1256/qj.02.141>, 2003.

Stenke, A., Grewe, V., and Ponater, M.: Lagrangian transport of water vapor and cloud water in the ECHAM4 GCM and its impact on the cold bias, *Clim. Dynam.*, 31, 491–506, <https://doi.org/10.1007/s00382-007-0347-5>, 2008.

Strandgren, J., Bugliaro, L., Sehnke, F., and Schröder, L.: Cirrus cloud retrieval with MSG/SEVIRI using artificial neural networks, *Atmos. Meas. Tech.*, 10, 3547–3573, <https://doi.org/10.5194/amt-10-3547-2017>, 2017.

Tao, M., Konopka, P., Wright, J. S., Liu, Y., Bian, J., Davis, S., Jia, Y., and Ploeger, F.: Multi-decadal variability controls short-term stratospheric water vapor trends, *Commun. Earth Environ.*, 4, 2662–4435, <https://doi.org/10.1038/s43247-023-01094-9>, 2023.

Teoh, R., Schumann, U., Majumdar, A., and Stettler, M. E. J.: Mitigating the Climate Forcing of Aircraft Contrails by Small-Scale Diversions and Technology Adoption, *Environ. Sci. Technol.*, 54, 2941–2950, <https://doi.org/10.1021/acs.est.9b05608>, 2020.

Teoh, R., Schumann, U., Gryspeerdt, E., Shapiro, M., Molloy, J., Koudis, G., Voigt, C., and Stettler, M. E. J.: Aviation contrail climate effects in the North Atlantic from 2016 to 2021, *Atmos. Chem. Phys.*, 22, 10919–10935, <https://doi.org/10.5194/acp-22-10919-2022>, 2022.

Teoh, R., Engberg, Z., Schumann, U., Voigt, C., Shapiro, M., Rohs, S., and Stettler, M. E. J.: Global aviation contrail climate effects from 2019 to 2021, *Atmos. Chem. Phys.*, 24, 6071–6093, <https://doi.org/10.5194/acp-24-6071-2024>, 2024.

Tiedtke, M.: Representation of clouds in large-scale models, *Mon. Weather Rev.*, 121, 3040–3061, [https://doi.org/10.1175/1520-0493\(1993\)121<3040:ROCILS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<3040:ROCILS>2.0.CO;2), 1993.

Tompkins, A., Gierens, K., and Rädel, G.: Ice supersaturation in the ECMWF Integrated Forecast System, *Q. J. Roy. Meteor. Soc.*, 133, 53–63, <https://doi.org/10.1002/qj.14>, 2007.

van der Linden, R., Knippertz, P., Fink, A. H., Ingleby, B., Maranan, M., and Benedetti, A.: The influence of DACCIA radiosonde data on the quality of ECMWF analyses and forecasts over southern West Africa, *Q. J. Roy. Meteor. Soc.*, 146, 1719–1739, <https://doi.org/10.1002/qj.3763>, 2020.

van Rossum, G. and Drake, F. L.: Python Reference Manual, CreateSpace, Scotts Valley, CA, USA, ISBN 1441412697, 2009.

Voigt, C., Schumann, U., Minikin, A., Abdelmonem, A., Afchine, A., Borrman, S., Boettcher, M., Buchholz, B., Bugliaro, L., Costa, A., Curtius, J., Dollner, M., Dörnbrack, A., Dreiling, V., Ebert, V., Ehrlich, A., Fix, A., Forster, L., Frank, F., Fütterer, D., Giez, A., Graf, K., Grooß, J.-U., Groß, S., Heimerl, K., Heinold, B., Hünemeier, T., Järvinen, E., Jurkat, T., Kaufmann, S., Kenntner, M., Klingebiel, M., Klimach, T., Kohl, R., Krämer, M., Krisna, T. C., Luebke, A., Mayer, B., Mertes, S., Molleker, S., Petzold, A., Pfeilsticker, K., Port, M., Rapp, M., Reutter, P., Rolf, C., Rose, D., Sauer, D., Schäfler, A., Schlage, R., Schnaiter, M., Schneider, J., Spelten, N., Spichtinger, P., Stock, P., Walser, A., Weigel, R., Weinzierl, B., Wendisch, M., Werner, F., Wernli, H., Wirth, M., Zahn, A., Ziereis, H., and Zöger, M.: ML-CIRRUS: The Airborne Experiment on Natural Cirrus and Contrail Cirrus with the High-Altitude Long-Range Research Aircraft HALO, *B. Am. Meteorol. Soc.*, 98, 271–288, <https://doi.org/10.1175/BAMS-D-15-00213.1>, 2017.

Voigt, C., Lelieveld, J., Schlager, H., Schneider, J., Curtius, J., Meerkötter, R., Sauer, D., Bugliaro, L., Bohn, B., Crowley, J. N., Erbertseder, T., Groß, S., Hahn, V., Li, Q., Mertens, M., Pöhlker, M. L., Pozzer, A., Schumann, U., Tomsche, L., Williams, J., Zahn, A., Andreae, M., Borrman, S., Bräuer, T., Dörich, R., Dörnbrack, A., Edtbauer, A., Ernle, L., Fischer, H., Giez, A., Granzin, M., Grewe, V., Harder, H., Heinritzi, M., Holanda, B. A., Jöckel, P., Kaiser, K., Krüger, O. O., Lucke, J., Marsig, A., Martin, A., Matthes, S., Pöhlker, C., Pöschl, U., Reifenberg, S., Ringsdorf, A., Scheibe, M., Tadic, I., Zauner-Wieczorek, M., Henke, R., and Rapp, M.: Cleaner skies during the COVID-19 lockdown, *B. Am. Meteorol. Soc.*, 103, E1796–E1827, <https://doi.org/10.1175/BAMS-D-21-0012.1>, 2022.

Wang, C.-C.: On the Calculation and Correction of Equitable Threat Score for Model Quantitative Precipitation Forecasts for Small Verification Areas: The Example of Taiwan, *Weather Forecast.*, 29, 788–798, <https://doi.org/10.1175/WAF-D-13-00087.1>, 2014.

Wang, Z., Bugliaro, L., Jurkat-Witschas, T., Heller, R., Burkhardt, U., Ziereis, H., Dekoutsidis, G., Wirth, M., Groß, S., Kirschler, S., Kaufmann, S., and Voigt, C.: Observations of microphysical properties and radiative effects of a contrail cirrus outbreak over the North Atlantic, *Atmos. Chem. Phys.*, 23, 1941–1961, <https://doi.org/10.5194/acp-23-1941-2023>, 2023.

Wang, Z., Letu, H., Shang, H., and Bugliaro, L.: Technical note: Retrieval of the supercooled liquid fraction in mixed-phase clouds from Himawari-8 observations, *Atmos. Chem. Phys.*, 24, 7559–7574, <https://doi.org/10.5194/acp-24-7559-2024>, 2024.

Wilhelm, L., Gierens, K., and Rohs, S.: Meteorological Conditions That Promote Persistent Contrails, *Appl. Sci.-Basel*, 12, 4450, <https://doi.org/10.3390/app12094450>, 2022.

Woiwode, W., Dörnbrack, A., Polichtchouk, I., Johansson, S., Harvey, B., Höpfner, M., Ungermaier, J., and Friedl-Vallon, F.: Technical note: Lowermost-stratosphere moist bias in ECMWF IFS model diagnosed from airborne GLORIA observations during winter–spring 2016, *Atmos. Chem. Phys.*, 20, 15379–15387, <https://doi.org/10.5194/acp-20-15379-2020>, 2020.

Wolf, K., Bellouin, N., Boucher, O., Rohs, S., and Li, Y.: Correction of ERA5 temperature and relative humidity biases by bivariate quantile mapping for contrail formation analysis, *Atmos. Chem. Phys.*, 25, 157–181, <https://doi.org/10.5194/acp-25-157-2025>, 2025.

Zängl, G., Reinert, D., Rípodas, P., and Baldauf, M.: The ICON (ICOahedral Non-hydrostatic) modelling framework of DWD and MPI-M: Description of the non-hydrostatic dynamical core, *Q. J. Roy. Meteor. Soc.*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.

Zhao, Y., Li, J., Zhang, L., Deng, C., Li, Y., Jian, B., and Huang, J.: Diurnal cycles of cloud cover and its vertical distribution over the Tibetan Plateau revealed by satellite observations, reanalysis datasets, and CMIP6 outputs, *Atmos. Chem. Phys.*, 23, 743–769, <https://doi.org/10.5194/acp-23-743-2023>, 2023.