

Platelet endothelial cell adhesion molecule-1 inhibits platelet response to thrombin and von Willebrand factor by regulating the internalization of glycoprotein Ib via AKT/glycogen synthase kinase-3/dynamin and integrin α IIb β 3

Article

Accepted Version

Jones, C. I. ORCID: <https://orcid.org/0000-0001-7537-1509>, Sage, T., Moraes, L. A., Vaiyapuri, S. ORCID: <https://orcid.org/0000-0002-6006-6517>, Hussain, U., Tucker, K. L., Barrett, N. E. ORCID: <https://orcid.org/0000-0001-9123-1100> and Gibbins, J. M. ORCID: <https://orcid.org/0000-0002-0372-5352> (2014) Platelet endothelial cell adhesion molecule-1 inhibits platelet response to thrombin and von Willebrand factor by regulating the internalization of glycoprotein Ib via AKT/glycogen synthase kinase-3/dynamin and integrin α IIb β 3. *Arteriosclerosis Thrombosis and Vascular Biology*, 34 (9). pp. 1968-1976. ISSN 1079-5642 doi: 10.1161/ATVBAHA.114.304097 Available at <https://centaur.reading.ac.uk/37995/>

work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1161/ATVBAHA.114.304097>

Publisher: Lippincott, Williams & Wilkins

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Platelet Endothelial Cell Adhesion Molecule-1 inhibits platelet response to thrombin and von Willebrand Factor by regulating the internalisation of Glycoprotein Ib via AKT/Glycogen Synthase Kinase-3/Dynamin and Integrin α IIb β 3

Chris I. Jones, Tanya Sage, Leonardo A. Moraes, Sakthivel Vaiyapuri, Umara Hussain, Katherine L. Tucker, Natasha E. Barrett, Jonathan M. Gibbins.

Institute for Cardiovascular and Metabolic Research, School of Biological Sciences, University of Reading, Reading, Berkshire, RG6 6AS, UK

Running title: PECAM-1 mediated internalisation of GPIb

Corresponding Author: Dr Chris Jones, Institute for Cardiovascular and Metabolic Research, School of Biological Sciences, University of Reading, Harbourne Building, Whiteknights, Reading, Berkshire, RG6 6AS, UK. Tel: +44 (0) 118 378 4653, Fax: +44(0) 118 378 7096, E-mail: c.i.jones@reading.ac.uk

Arterioscler Thromb Vasc Biol. 2014;34:1968-1976.

Abstract.

Objectives – Platelet endothelial cell adhesion molecule-1 (PECAM-1) regulates platelet response to multiple agonists. How this immunoreceptor tyrosine-based inhibitory motif-containing receptor inhibits G protein-coupled receptor-mediated thrombin-induced activation of platelets is unknown.

Approach and Results – Here, we show that the activation of PECAM-1 inhibits fibrinogen binding to integrin $\alpha\text{IIb}\beta 3$ and P-selectin surface expression in response to thrombin (0.1–3U/ml) but not thrombin receptor-activating peptides SFLLRN (3×10^{-7} – 1×10^{-5} M) and GYPGQV (3×10^{-6} – 1×10^{-4} M). We hypothesised a role for PECAM-1 in reducing the tethering of thrombin to GPIIb α on the platelet surface. We show that PECAM-1 signalling regulates the binding of FITC-labelled thrombin to the platelet surface and reduces the levels of cell surface GPIIb α by promoting its internalisation, while concomitantly reducing the binding of platelets to von Willebrand factor under flow *in vitro*. PECAM-1 mediated internalisation of GPIIb α was reduced in the presence of both EGTA, and/or Cytochalasin D and Latrunculin, but not either individually, and was reduced in mice in which tyrosines 747 and 759 of the cytoplasmic tail of $\beta 3$ integrin were mutated to phenylalanine. Furthermore PECAM-1 cross-linking led to a significant reduction in the phosphorylation of GSK-3 β Ser⁹, but interestingly an increase in GSK-3 α pSer²¹. PECAM-1 mediated internalisation of GPIIb α was reduced by inhibitors of Dynamin (Dynasore) and GSK-3 (CHIR99021), an effect that was enhanced in the presence of EGTA.

Conclusions – PECAM-1 mediates internalisation of GPIIb α in platelets through dual AKT/AKT/protein kinase B/glycogen synthase kinase-3/dynamin-dependent and $\alpha\text{IIb}\beta 3$ -dependent mechanisms. These findings expand our understanding of how PECAM-1 regulates non-immunoreceptor signalling pathways and helps to explain how PECAM-1 regulates thrombosis.

Nonstandard abbreviations and acronyms	
--	--

PMI	PECAM-1 mediated internalisation
-----	----------------------------------

Introduction

Platelet endothelial cell adhesion molecule-1 (PECAM-1) is a 130-kDa transmembrane glycoprotein expressed on the platelet surface at between 5000 and 8800 copies per cell.¹⁻³ It is composed of a 574 amino acid residue extracellular portion organised into six immunoglobulin (Ig)-like homology domains, a 19 amino acid transmembrane domain and a cytoplasmic domain that contains an Immunoreceptor Tyrosine-based Inhibitory Motif (ITIM) ((L/V/I/S/T)XYXX(L/V)) and an Immunoreceptor Tyrosine-based Switch Motif (ITSM) (TxYxx(V/I)).^{4, 5} The tyrosine residues (residues 663 and 686) within the ITIM and ITSM become tyrosine phosphorylated by Src or Csk family kinases,⁶ following homophilic ligation or stimulation of platelets through protease-activated receptor (PAR)-1 and 4, glycoprotein VI (GPVI), or GPIb in a process that is largely, but not wholly, dependent on integrin $\alpha\text{IIb}\beta\text{3}$ -dependent platelet aggregation.⁷⁻¹⁰

The inhibition by PECAM-1 of immunoreceptor signalling downstream of GPVI is well established. Once phosphorylated these twin ITIM and ITSM tyrosine residues support the recruitment of Src-homology 2 (SH2) domain containing proteins, including the protein-tyrosine phosphatases SHP-2, and to a lesser extent SHP-1,¹¹ to which the p85 regulatory subunit of phosphatidylinositol 3-kinase (PI3-K) becomes associated destabilising its association with the linker for activation of T cells (LAT) and the scaffolding molecule Grb-2-associated binding protein-1 (Gab1).¹² By disturbing the LAT-GAB1-PI3K signalling complex PECAM-1, therefore, disrupts the recruitment and activation of phospholipase C (PLC γ 2),¹² thereby inhibiting immune-like signalling downstream of GPVI and reducing the activation of platelets by collagen, the GPVI specific ligand cross-linked collagen-related peptide (CRP-XL) and convulxin (CVX).^{10, 13, 14}

More surprising is the ability of PECAM-1 to reduce platelet activation through GPCR mediated signalling pathways following stimulation by thrombin or ADP. Whilst hyper-reactivity to thrombin and ADP has not been detected in PECAM-1 deficient mice,¹⁵ pre-activation of PECAM-1 has been shown to reduce fibrinogen binding to platelets, secretion and calcium mobilisation in response to ADP and thrombin,¹⁴ and in a laser injury model of thrombosis in cremaster muscle arterioles, that is dependent on thrombin generation rather than collagen exposure,¹⁶ thrombi formed more rapidly, were more stable and approximately 35% larger in PECAM-1 deficient mice.¹⁷ The mechanism by which PECAM-1 inhibits GPCR-mediated activation of platelets is as yet unclear.

While the inhibitory effect of PECAM-1 on individual pathways is modest, corresponding to 5-15% inhibition in platelet function, the physiological importance of PECAM-1 comes from its ability to inhibit multiple activation pathway. The combination of these effects account for a substantial inhibition in thrombus formation.^{14, 17}

We have previously reported that although PECAM-1 inhibits the activation of platelets by thrombin, it does not inhibit activation stimulated by TRAP. Here we investigate this disparity and show that the activation of PECAM-1 triggers the internalisation of GPIb by dual AKT/GSK-3/dynamin and $\alpha\text{IIb}\beta\text{3}$ -dependent mechanisms, which result in the reduction in platelet binding of, and hence response to, thrombin, and reduced platelet binding to VWF.

Materials and methods

Materials and Methods are available in the online-only Data Supplement

Results

Activation of PECAM-1 inhibits the binding of thrombin to human platelets and their subsequent activation.

Consistent with our previous work, activating PECAM-1 by cross-linking antibodies that recognise the sixth Ig domain of PECAM-1 (AB468), resulting in PECAM-1 phosphorylation,¹²⁻¹⁴ led to a small (5-15%) but significant inhibition in platelet fibrinogen binding and P-selectin exposure in response to thrombin across a range of concentration ($p < 0.0001$ and $p = 0.001$, respectively, two-way ANOVA) (Figure 1A). Again confirming our previous findings PECAM-1 cross-linking did not inhibit platelet response to either PAR-1 or PAR-4 activating peptides (Figure 1A). The inhibition of thrombin activation of platelets without inhibiting PAR signalling suggests a role for PECAM-1 in modulating the binding of thrombin to platelets. Consistent with this cross-linking PECAM-1 led to a significant reduction in the binding to FITC-labelled thrombin and a concomitant reduction of platelet activation measured by the surface exposure of P-selectin (Figure 1B and Supplementary Figure I).

This reduction of thrombin binding was specific to the activation of PECAM-1. To control for the potential steric inhibition caused by the binding and cross-linking of the PECAM-1 antibody we repeated these experiment using a different PECAM-1 antibody (WM59) that binds PECAM-1 but prevents its activation, thereby providing a control for any steric effects. No reduction in thrombin binding was seen when platelets were incubated with antibodies against the 1st or 2nd Ig domain of PECAM-1 (WM59) (which prevent homophilic ligation and the activation of PECAM-1¹⁸) and the cross-linking antibody (Figure 1C). Similarly the binding and cross-linking of antibodies against GPIb α , GPVI, PAR-1 and PAR-4 had no effect on the binding of thrombin to the platelet surface (Figure 1C).

PECAM-1 crosslinking reduces the levels of GPIb α at the cell surface.

GPIb α as part of the GPIb/V/IX complex plays an integral role in thrombin activation. It is a high affinity receptor for thrombin¹⁹ and acts as a cofactor for PAR cleavage.²⁰ Furthermore PECAM-1 plays a role in GPIb signalling, becoming tyrosine phosphorylated upon VWF binding to GPIb α , and PECAM-1 deficient mice show enhanced aggregation in response to VWF.⁹ We hypothesised that the reduction in thrombin binding to platelets may be due to reduced surface expression of GPIb α following PECAM-1 activation. PECAM-1 cross-linking resulted in a significant reduction in the binding of antibodies against GPIb α , and a reduction in antibodies against GPIb β (Figure 2A and Supplementary Figure II). Although the level of GPIb α expressed on the surface of resting platelets varied between subjects the reduction in GPIb α following PECAM-1 cross-linking occurred in all subjects. The action of PECAM-1 appeared to be specific for GPIb and did not cause a reduction in the binding of antibodies against PAR-1 and PAR-4, nor GPVI (Figure 2A).

To confirm that this reduction was due to PECAM-1, PECAM-1 on the surface of WT mouse platelets was cross-linked using antibodies that recognise the sixth Ig domain of mouse PECAM-1 (M-185), and, as with human blood resulted in a significant decrease in GPIb α on the platelet surface. No such decrease was seen in PECAM-1 deficient mouse platelets (Figure 2B). Again to control for the effects of steric inhibitions caused by the binding and cross-linking of the PECAM-1 antibody we repeated these experiments using a different PECAM-1 antibody (WM59) that binds PECAM-1 but prevents its activation. No reduction in GPIb α was seen when platelets were incubated with antibodies against the 1st or 2nd Ig domain of PECAM-1 (WM59) and the cross-linking antibody (Figure 2C).

PECAM-1 activation inhibits platelet binding to VWF

Given these findings we hypothesised that PECAM-1 cross-linking should also reduce platelet adhesion to VWF. The number of platelets in whole blood binding to VWF (100 μ g/ml) under arterial flow conditions was indeed reduced following PECAM-1 cross-

linking (Figure 2D-G). This supports the role of PECAM-1 in regulating surface GPIb α and is in line with previous reports showing enhanced aggregation in response to VWF in PECAM-1 deficient mice.⁹

PECAM-1 mediated reduction of platelet surface GPIb α is due to internalisation and not cleavage.

We sought to determine if the PECAM-1 mediated reduction in the surface expression of GPIb α was due to internalisation or cleavage of the receptor. Previous reports have demonstrated the cleavage of GPIb α by ADAM17, generating a ~95kDa fragment and a ~45kDa N-terminal fragment.²¹ Using a purified cobra metalloproteinase, Nk (10 μ g/mL)²² as a positive control we identified both the ~130kDa intact GPIb α and the ~95kDa and ~45kDa fragments by Western blotting, using antibodies against the C-terminal and N-terminal of GPIb α respectively (Figure 3A). Neither of these cleaved forms of GPIb α were observed in lysates of resting, CRP-XL stimulated or PECAM-1 stimulated platelets, indicating that no detectable cleavage of GPIb α occurred (Figure 3A). Furthermore, there was no evidence of any GPIb α cleavage products in these samples. Although there is a smear in the lane of the PECAM-1 stimulated platelets, this appears at the same height when probing with both the N-terminal and C-terminal antibodies and is present in the antibody control lanes. We therefore conclude that this smear is not due to cleavage of GPIb α .

To further test if GPIb α was cleaved or internalised we measured the binding of GPIb α antibodies to non-permeabilised and permeabilised platelets. As in all previous experiments the binding of GPIb α antibodies to non-permeabilised platelets (i.e. binding solely to GPIb α expressed on the surface of platelets) decreased following PECAM-1 stimulation. There was, however, no reduction in binding of GPIb α antibodies to permeabilised platelets (i.e. when the GPIb antibody could access both external and internal pools of GPIb) indicating that there was no loss of GPIb α from the platelets (Figure 3B). Taken together these two pieces of evidence suggest that PECAM-1 mediated reduction of platelet surface GPIb α is due to internalisation and not cleavage.

PECAM-1 internalisation of GPIb α is dependent on cytoskeletal rearrangement and exogenous calcium ions.

The internalisation of GPIb α is well known following platelet activation,²³ and PECAM-1 has, in other cells types, been shown to be involved in non-clathrin mediated endocytosis.²⁴ To elucidate the mechanism by which PECAM-1 mediated internalisation (PMI) of GPIb α occurs in platelets the reduction in surface expression of GPIb α was measured following PECAM-1 activation in the presence of EGTA, to chelate extracellular calcium ions, and/or Cytochalasin D and Latrunculin, to prevent actin polymerisation. Individually none of the compounds significantly inhibited PMI of GPIb α . A combination of EGTA and either Cytochalasin D or Latrunculin did, however, completely abrogate the effect of PECAM-1 activation on the surface expression of GPIb α (Figure 4A). By contrast GPIb α internalisation following stimulation of GPVI by CRP-XL was not inhibited by EGTA alone but was completely inhibited by either Cytochalasin D or Latrunculin (Figure 4B). This indicates that while GPVI mediated internalisation of GPIb α is entirely dependent on cytoskeletal rearrangement, the activation of PECAM-1 initiates two separate mechanisms, one dependent on cytoskeleton and the other dependent on exogenous divalent cations, both of which are capable of modulating the surface exposure of GPIb α independently.

Chelating extracellular calcium inhibits the activation of integrin α IIb β 3, the activation of which is known to be enhanced by PECAM-1.^{15, 25} To establish whether integrin α IIb β 3 is involved in PMI of GPIb α , we used knock-in (KI) mice (DiYF mice) in which tyrosines 747 and 759 of the cytoplasmic tail of the β 3 integrin were mutated to phenylalanine, preventing tyrosine phosphorylation, and selectively impairing outside-in α IIb β 3 signalling.²⁶ Internalisation of GPIb α following PECAM-1 cross-linking was reduced in heterozygous mice

and abolished in KI mice (Figure 4C). In corroboration of our findings in humans, internalisation of GPIIb α following stimulation with CRP-XL was unaltered in DiYF mice (Figure 4D). There was no significant difference between WT, heterozygous or homozygous KI mice in the surface expression of either α IIb β 3 or GPVI (Figure 4E, F).

PMI of GPIIb α occurs in an AKT/GSK-3/Dynamin dependent manner

PECAM-1 clustering has been reported to induce Dynamin-2 dependent endocytosis in endothelial cells,^{24, 27} while in platelets dynamin has been shown to be critical to the internalisation of both P2Y₁ and P2Y₁₂.²⁸ To investigate the role of dynamin in PMI of GPIIb α , platelets were incubated with Dynasore (100 μ M) prior to antibody mediated PECAM-1 cross-linking. PMI of GPIIb α was reduced in the presence of Dynasore and almost abolished in the presence of Dynasore and EGTA (Figure 5A). In part the action of dynamin in endocytosis is regulated by the de-phosphorylation (by calcineurin^{29, 30}) and re-phosphorylation (by cdk5 and GSK-3^{31, 32}) of two serine residues in the C-terminal proline-rich domain (PRD) of dynamin. While de-phosphorylation of dynamin is required for clathrin-mediated endocytosis in neurons, non-clathrin-mediated bulk endocytosis (of the type previously shown to result from PECAM-1 activation in endothelial cells²⁴) requires both de-phosphorylation and re-phosphorylation.^{32, 33} We have previously shown that PECAM-1 crosslinking resulted in reduced phosphorylation of AKT at Ser⁴⁷³.¹² This should lead to a reduction in the phosphorylation of GSK-3, which would in turn regulate its kinase activity. Following stimulation of platelets with CRP-XL the phosphorylation of GSK-3 β Ser⁹ increased compared to the level in unstimulated cells, however, following PECAM-1 crosslinking the phosphorylation of GSK-3 β Ser⁹ decreased (Figure 5B-C). Interestingly the same pattern was not repeated for GSK-3 α . Ser²¹ phosphorylation of GSK-3 α did not increase significantly in response to CRP-XL but did increase significantly following PECAM-1 crosslinking (Figure 5B-C). To confirm the role of GSK-3 in PMI we used the highly selective inhibitor of both GSK-3 isoforms, CHIR99021, which inhibited the internalisation of GPIIb α following the cross-linking of PECAM-1 (Figure 5D). Together this data suggests that the internalisation of GPIIb α following PECAM-1 stimulation occurs in a non-clathrin, AKT/GSK-3/Dynamin dependant manner which may be similar to bulk endocytosis reported in endothelial or neuronal cells.

Discussion

Two of the enduring questions surrounding the role of PECAM-1 in regulating platelet response are how this ITIM containing molecule inhibits non-immunoreceptor signalling, and how a molecule with what appears to be a moderate impact on individual signalling pathways can have such a profound effect on thrombus formation *in vitro* and *in vivo*.^{14, 17} By investigating the disparity between the action of PECAM-1 on platelet response to thrombin and TRAP we have identified a PECAM-1 mediated receptor internalisation pathway mediated via dual AKT/GSK-3/Dynamin and α IIb β 3 dependent mechanisms that result in the internalisation of GPIIb α , a reduction in thrombin binding to, and activation of, platelets and reduced platelet binding to VWF. The physiological impact of PMI of GPIIb α on the binding of platelets to vessel-bound VWF is uncertain as it is unlikely that PECAM-1 is activated before platelets bind to VWF and adhere to the site of vessel damage. This novel role for PECAM-1 in platelets does, however, explain how PECAM-1 inhibits GPCR-mediated activation of platelets by thrombin, and why it has such a potent impact on thrombosis. By simultaneously reducing thrombin stimulation through PMI of GPIIb α , and GPVI signalling through its action as an ITIM containing receptor, PECAM-1 directly regulates two of the major pathways by which platelets become activated and propagate thrombus formation.

PMI of GPIIb α appears to occur via a distinct mechanism that is separate from the internalisation of GPIIb α that follows GPVI stimulation. This was seen in the internalisation of

GPIb α in response to CRP-XL in PECAM-1 KO mice, and the ability of Cytochalasin D and Latrunculin alone to inhibit CRP-XL mediated internalisation of GPIb α , but not PMI. It is likely, however, that *in vivo* these two mechanisms work in concert, particularly as PECAM-1 becomes phosphorylated following GPVI stimulation, and as PECAM-1 and GPVI have opposing effects on AKT and hence GSK-3 phosphorylation. Why such opposing signalling events result in such similar outcomes is not yet known. The importance of PMI during thrombus formation may in part come from the activation of PECAM-1 by homophilic ligation, which only occurs during close platelet-platelet contact. Under these circumstances stimulation of receptor internalisation or endocytosis by PECAM-1 at the site of contact may be a mechanism for recycling receptors that are not occupied or bound into focal adhesion complexes. Trafficking these receptors away from the site of contact to areas of the platelet that are not in contact with other cells provides an attractive mechanism by which to maximise the efficiency of platelet response to multiple simultaneous stimuli and orchestrate thrombus growth.

The mechanism by which integrin β 3 regulates PMI is as yet unknown. Its effect, however, appears to be critical. One explanation for this, perhaps lies with filamin. Filamin binds to GPIb α in resting platelets anchoring it to the cytoskeleton,³⁴ helping to maintain the structure and integrity of the plasma membrane following platelet tethering to VWF under high shear.³⁵ Upon stimulation filamin becomes detached from the cytoplasmic tail GPIb α allowing its translocation.³⁶ By contrast filamin binds to α IIb β 3 following platelet activation. One tempting hypothesis is that PECAM-1 plays a role, either directly or indirectly, in the decoupling of filamin from GPIb α and its subsequent binding to integrin β 3. We have as yet been unable to confirm or refute this hypothesis but it will be the focus of future work.

Of the two isoforms of GSK-3 it is the β form that is thought to be predominant in platelets.³⁷ Proteomic analysis has, however, identified that the copy number per platelet for each isoform is similar, 1300 for GSK-3 β and 1000 for GSK-3 α .³⁸ Both isoforms are constitutively active in resting platelets but are inhibited by phosphorylation of Ser²¹ (GSK-3 α) or Ser⁹ (GSK-3 β) by PKC or AKT upon platelet stimulation.³⁹ It is thought that GSK-3 acts as a negative regulator of platelets; GSK-3 inhibitors increase platelet responses, GSK-3 β +/- mice display increased aggregation and thrombus formation, and platelets from mice in which GSK-3 α Ser²¹ and GSK-3 β Ser⁹ have been mutated to Ala show reduced response.^{37, 39} The reduction on GSK-3 β pSer⁹ following PECAM-1 cross-linking fits with this understanding of platelet GSK-3. PECAM-1, which has a well-established inhibitory effect on platelet signalling, reduces AKT phosphorylation and therefore reduces the phosphorylation and inhibition GSK-3 β , thereby enhancing its inhibitory effect, some of which may be mediated through Dynamin and translocation of surface receptors, as we have shown to be the case with GPIb. The increase in GSK-3 α phosphorylation following PECAM-1 cross-linking does not, however, fit with our current understanding. Part of the reason for this is that all previous work on GSK-3 in platelets has focused either on GSK-3 β alone, as in the case of the GSK-3 β +/- mice, or has assumed that both isoforms work in a similar way and have therefore inhibited both isoforms. In the case of dual inhibition the effect of GSK-3 α may be masked by GSK-3 β making any interpretation of the physiological relevance of GSK-3 α impossible. Although there is 85% similarity in the amino acid sequence between the isoforms and 98% homology in the kinase domain⁴⁰ there is growing evidence that the two isoforms of GSK-3 have distinct roles.⁴¹⁻⁴³ Quite why the activation of PECAM-1 has such contrasting effects on the two GSK-3 isoforms and what significance GSK-3 α phosphorylation has in PECAM-1 signalling are as yet unknown.

The inhibition of platelet activation by PECAM-1 is well known but the mechanisms by which this ITIM-containing receptor inhibits platelet response to thrombin were not. Here we have shown that the activation of PECAM-1 results in the internalisation of GPIb by dual GSK-3/Dynamin and α IIb β 3 dependent mechanisms, which result in the reduction in platelet binding of, and hence response to, thrombin, and reduced platelet binding to VWF.

Furthermore, we have for the first time shown the differential regulation of GSK-3 α/β by PECAM-1 in platelets. This novel role for PECAM-1 expands our understanding of how this promiscuous molecule regulates multiple diverse signalling pathways and why it has such a potent impact on thrombosis. It may also suggest a role for PECAM-1 in regulating conditions where the generation of thrombin, and the response of platelets to thrombin is central to pathology.

Sources of Funding

This study was supported by British Heart Foundation grants to J.M.G. (RG/05/007, RG/09/011 and FS/07/018).

Disclosures

None.

References

1. Newman PJ. The role of pecam-1 in vascular cell biology. *Ann N Y Acad Sci.* 1994;714:165-174
2. Wu XW, Lian EC. Binding properties and inhibition of platelet aggregation by a monoclonal antibody to cd31 (pecam-1). *Arterioscler Thromb Vasc Biol.* 1997;17:3154-3158
3. Metzelaar MJ, Korteweg J, Sixma JJ, Nieuwenhuis HK. Biochemical characterization of pecam-1 (cd31 antigen) on human platelets. *Thromb Haemost.* 1991;66:700-707
4. Jackson DE, Kupcho KR, Newman PJ. Characterization of phosphotyrosine binding motifs in the cytoplasmic domain of platelet/endothelial cell adhesion molecule-1 (pecam-1) that are required for the cellular association and activation of the protein-tyrosine phosphatase, shp-2. *J. Biol. Chem.* 1997;272:24868-24875
5. Vivier E, Daeron M. Immunoreceptor tyrosine-based inhibitory motifs. *Immunol. Today.* 1997;18:286-291
6. Cao MY, Huber M, Beauchemin N, Famiglietti J, Albelda SM, Veillette A. Regulation of mouse pecam-1 tyrosine phosphorylation by the src and csk families of protein-tyrosine kinases. *The Journal of biological chemistry.* 1998;273:15765-15772
7. Cicmil M, Thomas JM, Sage T, Barry FA, Leduc M, Bon C, Gibbins JM. Collagen, convulxin, and thrombin stimulate aggregation-independent tyrosine phosphorylation of cd31 in platelets. Evidence for the involvement of src family kinases. *J. Biol. Chem.* 2000;275:27339-27347
8. Jackson DE, Ward CM, Wang R, Newman PJ. The protein-tyrosine phosphatase shp-2 binds platelet/endothelial cell adhesion molecule-1 (pecam-1) and forms a distinct signaling complex during platelet aggregation. *J. Biol. Chem.* 1997;272:6986-6993
9. Rathore V, Stapleton MA, Hillery CA, Montgomery RR, Nichols TC, Merricks EP, Newman DK, Newman PJ. Pecam-1 negatively regulates gpib/v/ix signaling in murine platelets. *Blood.* 2003;102:3658-3664
10. Jones KL, Hughan SC, Dopheide SM, Farndale RW, Jackson SP, Jackson DE. Platelet endothelial cell adhesion molecule-1 is a negative regulator of platelet-collagen interactions. *Blood.* 2001;98:1456-1463
11. Hua CT, Gamble JR, Vadas MA, Jackson DE. Recruitment and activation of shp-1 protein-tyrosine phosphatase by human platelet endothelial cell adhesion molecule-1 (pecam-1). Identification of immunoreceptor tyrosine-based inhibitory motif-like binding motifs and substrates. *J. Biol. Chem.* 1998;273:28332-28340
12. Moraes LA, Barrett NE, Jones CI, Holbrook LM, Spyridon M, Sage T, Newman DK, Gibbins JM. Pecam-1 regulates collagen-stimulated platelet function by modulating the association of pi3 kinase with gab1 and lat. *J Thromb Haemost.* 2010;8:2530-2541
13. Cicmil M, Thomas JM, Leduc M, Bon C, Gibbins JM. Pecam-1 signalling inhibits the activation of human platelets. *Blood.* 2002;99:137-144
14. Jones CI, Garner SF, Moraes LA, Kaiser WJ, Rankin A, Ouwehand WH, Goodall AH, Gibbins JM. Pecam-1 expression and activity negatively regulate multiple platelet signaling pathways. *FEBS Lett.* 2009;583:3618-3624
15. Wee JL, Jackson DE. The ig-itim superfamily member pecam-1 regulates the "outside-in" signaling properties of integrin alpha(iiib)beta(3) in platelets. *Blood.* 2005;106:3816-3823
16. Falati S, Gross P, Merrill-Skoloff G, Furie BC, Furie B. Real-time in vivo imaging of platelets, tissue factor and fibrin during arterial thrombus formation in the mouse. *Nature medicine.* 2002;8:1175-1181
17. Falati S, Patil S., Gross PL, Stapleton M, Merrill-Skoloff G., Barrett NE, Pixton KL, Weiler H., Cooley B., Newman D.K., Newman PJ, Furie BC, Furie B., Gibbins JM. Platelet pecam-1 inhibits thrombus formation in vivo. *Blood.* 2006;107:535-541
18. Sun QH, DeLisser HM, Zukowski MM, Paddock C, Albelda SM, Newman PJ. Individually distinct ig homology domains in pecam-1 regulate homophilic binding and modulate receptor affinity. *The Journal of biological chemistry.* 1996;271:11090-11098

19. Buchanan SG, Gay NJ. Structural and functional diversity in the leucine-rich repeat family of proteins. *Progress in Biophysics and Molecular Biology*. 1996;65:1-44
20. De Candia E, Hall SW, Rutella S, Landolfi R, Andrews RK, De Cristofaro R. Binding of thrombin to glycoprotein Ib accelerates the hydrolysis of PAR-1 on intact platelets. *The Journal of biological chemistry*. 2001;276:4692-4698
21. Gardiner EE, Karunakaran D, Shen Y, Arthur JF, Andrews RK, Berndt MC. Controlled shedding of platelet glycoprotein (gp)VI and GPIIb-IIIa by ADAM family metalloproteinases. *J Thromb Haemost*. 2007;5:1530-1537
22. Wijeyewickrema LC, Gardiner EE, Shen Y, Berndt MC, Andrews RK. Fractionation of snake venom metalloproteinases by metal ion affinity: A purified cobra metalloproteinase, nk, from *Naja kaouthia* binds Ni²⁺-agarose. *Toxicon : official journal of the International Society on Toxinology*. 2007;50:1064-1072
23. Han Y, Nurden A, Combrie R, Pasquet JM. Redistribution of glycoprotein Ib within platelets in response to protease-activated receptors 1 and 4: Roles of cytoskeleton and calcium. *J Thromb Haemost*. 2003;1:2206-2215
24. Muro S, Wiewrodt R, Thomas A, Koniaris L, Albelda SM, Muzykantov VR, Koval M. A novel endocytic pathway induced by clustering endothelial ICAM-1 or PECAM-1. *Journal of cell science*. 2003;116:1599-1609
25. Varon D, Jackson DE, Shenkman B, Dardik R, Tamarin I, Savion N, Newman PJ. Platelet/endothelial cell adhesion molecule-1 serves as a costimulatory agonist receptor that modulates integrin-dependent adhesion and aggregation of human platelets. *Blood*. 1998;91:500-507
26. Law DA, DeGuzman FR, Heiser P, Ministri-Madrid K, Killeen N, Phillips DR. Integrin cytoplasmic tyrosine motif is required for outside-in α IIb β 3 signalling and platelet function. *Nature*. 1999;401:808-811
27. Garnacho C, Shuvaev V, Thomas A, McKenna L, Sun J, Koval M, Albelda S, Muzykantov V, Muro S. RhoA activation and actin reorganization involved in endothelial ICAM-mediated endocytosis of anti-PECAM carriers: Critical role for tyrosine 686 in the cytoplasmic tail of PECAM-1. *Blood*. 2008;111:3024-3033
28. Mundell SJ, Barton JF, Mayo-Martin MB, Hardy AR, Poole AW. Rapid resensitization of purinergic receptor function in human platelets. *J Thromb Haemost*. 2008;6:1393-1404
29. Liu JP, Sim AT, Robinson PJ. Calcineurin inhibition of dynamin I GTPase activity coupled to nerve terminal depolarization. *Science*. 1994;265:970-973
30. Marks B, McMahon HT. Calcium triggers calcineurin-dependent synaptic vesicle recycling in mammalian nerve terminals. *Current biology : CB*. 1998;8:740-749
31. Clayton EL, Sue N, Smillie KJ, O'Leary T, Bache N, Cheung G, Cole AR, Wyllie DJ, Sutherland C, Robinson PJ, Cousin MA. Dynamin I phosphorylation by GSK3 controls activity-dependent bulk endocytosis of synaptic vesicles. *Nature neuroscience*. 2010;13:845-851
32. Tan TC, Valova VA, Malladi CS, Graham ME, Berven LA, Jupp OJ, Hansra G, McClure SJ, Sarcevic B, Boadle RA, Larsen MR, Cousin MA, Robinson PJ. Cdk5 is essential for synaptic vesicle endocytosis. *Nature cell biology*. 2003;5:701-710
33. Clayton EL, Anggono V, Smillie KJ, Chau N, Robinson PJ, Cousin MA. The phospho-dependent dynamin-syndapin interaction triggers activity-dependent bulk endocytosis of synaptic vesicles. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2009;29:7706-7717
34. Andrews RK, Fox JE. Identification of a region in the cytoplasmic domain of the platelet membrane glycoprotein GPIIb-IIIa complex that binds to purified actin-binding protein. *The Journal of biological chemistry*. 1992;267:18605-18611
35. Cranmer SL, Ashworth KJ, Yao Y, Berndt MC, Ruggeri ZM, Andrews RK, Jackson SP. High shear-dependent loss of membrane integrity and defective platelet adhesion following disruption of the GPIIb-IIIa-filamin interaction. *Blood*. 2011;117:2718-2727

36. Dyson JM, Munday AD, Kong AM, Huysmans RD, Matzaris M, Layton MJ, Nandurkar HH, Berndt MC, Mitchell CA. Ship-2 forms a tetrameric complex with filamin, actin, and gpib-ix-v: Localization of ship-2 to the activated platelet actin cytoskeleton. *Blood*. 2003;102:940-948
37. Li D, August S, Woulfe DS. Gsk3beta is a negative regulator of platelet function and thrombosis. *Blood*. 2008;111:3522-3530
38. Burkhardt JM, Vaudel M, Gambaryan S, Radau S, Walter U, Martens L, Geiger J, Sickmann A, Zahedi RP. The first comprehensive and quantitative analysis of human platelet protein composition allows the comparative analysis of structural and functional pathways. *Blood*. 2012;120:e73-82
39. Moore SF, van den Bosch MT, Hunter RW, Sakamoto K, Poole AW, Hers I. Dual regulation of glycogen synthase kinase 3 (gsk3)alpha/beta by protein kinase c (pkc)alpha and akt promotes thrombin-mediated integrin alphaIIb beta3 activation and granule secretion in platelets. *The Journal of biological chemistry*. 2013;288:3918-3928
40. Woodgett JR. Molecular cloning and expression of glycogen synthase kinase-3/factor a. *The EMBO journal*. 1990;9:2431-2438
41. Azoulay-Alfaguter I, Yaffe Y, Licht-Murava A, Urbanska M, Jaworski J, Pietrokovski S, Hirschberg K, Eldar-Finkelman H. Distinct molecular regulation of glycogen synthase kinase-3alpha isozyme controlled by its n-terminal region: Functional role in calcium/calpain signaling. *The Journal of biological chemistry*. 2011;286:13470-13480
42. Ciaraldi TP, Nikoulina SE, Bandukwala RA, Carter L, Henry RR. Role of glycogen synthase kinase-3 alpha in insulin action in cultured human skeletal muscle cells. *Endocrinology*. 2007;148:4393-4399
43. Hoeflich KP, Luo J, Rubie EA, Tsao MS, Jin O, Woodgett JR. Requirement for glycogen synthase kinase-3beta in cell survival and nf-kappab activation. *Nature*. 2000;406:86-90

Significance.

We show two new facets of PECAM-1 biology in platelets. 1) That PECAM-1 mediates the internalisation of GPIIb/IIIa leading to a reduction in the activation of platelets by thrombin and a reduction in their binding to VWF. 2) That PECAM-1 differentially regulates the phosphorylation of the two isoforms of GSK-3. In so doing this article answers a number of the outstanding questions concerning the regulation of platelets by PECAM-1; how this ITIM containing molecule inhibits non-immunoreceptor signalling, and importantly, how PECAM-1 has such a profound effect on thrombus formation *in vitro* and *in vivo*.

Figure 1 – Activation of PECAM-1 inhibits the binding of thrombin to human platelets and their subsequent activation. (A) Fibrinogen binding (top row) and P-selectin expression (bottom row) to human platelets in response to thrombin, PAR-1 or PAR-4 activating peptides in the presence (black) or absence (grey) of PECAM-1 (AB468) crosslinking and activation was measured by flow cytometry (n=5). P-values compare the difference between the two curves over the entire concentration range using Two-way ANOVA with Bonferroni post-test analysis of the difference at individual concentrations. (B) To ascertain if reduced platelet response to thrombin was due to reduced thrombin binding, the reduction in the binding of FITC-labelled thrombin to human platelets (left) and the resultant surface exposure of P-selectin (right) was measured following PECAM-1 crosslinking. The binding of FITC-thrombin to platelets was measured by flow cytometry and is presented as the median fluorescent intensity (MFI) (left hand axis - presence (black line) or absence (grey line) of PECAM-1 (AB468) crosslinking) and the percentage reduction in median fluorescent intensity (MFI) in PECAM-1 cross-linked samples compared to the isotype control (right hand axis) (n=5). (C) To check that this reduction in thrombin binding was specific to the activation of PECAM-1 and not the binding of antibodies to the platelet surface, the binding of FITC labelled thrombin to human platelets in the presence of cross-linked isotype control, PECAM-1 (WM59 - inhibitory), GPIIb/IIIa, GPVI, PAR-1 or PAR-4 antibodies was measured (n=3). In all cases *p<0.05, **p<0.01, ***p<0.001.




Figure 2 – Activation of PECAM-1 reduces the surface expression of GPIIb α and reduces platelet binding to vWF. (A) Human platelet surface expression of GPIIb α , GPIIb β , GPVI, PAR-1, PAR-4 in the presence (black) or absence (grey) of PECAM-1 (AB468) crosslinking was measured by flow cytometry (n=12 for anti-GPIIb α , and n=3 for all other experiments). (B) To confirm that this was PECAM-1 specific the expression of GPIIb α on the surface of mouse platelets from WT (n=4) or PECAM-1 KO (n=6) mice following the crosslinking of isotype control (grey) or PECAM-1 stimulatory antibodies (black) was measured. (C) To check that this reduction in the surface of GPIIb α was specific to the activation of PECAM-1 and not the binding of antibodies to the platelet surface, human platelet surface expression of GPIIb α was measured in the presence of cross-linked isotype control (grey), PECAM-1 (WM59 - inhibitory)(black)(n=4). The difference in MFI between (A) and (C) are because they were performed on different flow cytometers, a FACSCalibur and an Accuri C6, respectively. (D-G) Given the importance of GPIIb α in platelet tethering to VWF we assessed the impact of PECAM-1 crosslinking on the binding of platelets to VWF under flow. (D) Representative images showing human platelets binding to VWF (100 μ g/ml) under flow conditions following the crosslinking of isotype control or PECAM-1 stimulatory antibodies, or, as a positive control, antibodies that block the binding of GPIIb to VWF (Haematologic Technologies Inc). (E) The number of human platelets binding to VWF (100 μ g/ml), (F) the area covered by, and (G) the sum fluorescence intensity of these platelets following the crosslinking of isotype control (grey) or PECAM-1 (black) stimulatory antibodies (n=3). *p<0.05, **p<0.01, ***p<0.001




Figure 3 – GPIb α is internalised and not cleaved following PECAM-1 activation. (A) To determine whether GPIb α is cleaved following PECAM-1 activation, human platelets that had been stimulated for 90 or 600 seconds with CRP-XL ($\mu\text{g/mL}$), or undergone PECAM-1 activation via antibody crosslinking, or been exposed to purified cobra metalloproteinase, Nk (10 $\mu\text{g/mL}$), were probed by western blotting using antibodies that recognise either the C or the N-terminal domain of GPIb α . Cleavage products were clearly visible in samples treated with Nk but not CRP-XL or PECAM-1 stimulated platelets. The smear present in the PECAM-1 stimulated samples appears at the same height when probing with both the N-terminal and C-terminal antibodies and appears when the antibodies are run alone in the absence of platelets. We therefore conclude that it is the result of protein in the stimulating antibodies rather than a result of cleavage. (B) To identify the internalisation of GPIb α following PECAM-1 activation the binding of GPIb α antibodies to non-permeabilised human platelets (i.e. binding to GPIb α expressed on the surface of platelets) or to permeabilised platelets (i.e. binding to all GPIb α , internal and surface expressed) following PECAM-1 cross-linking was measured by flow cytometry (n=3). *p<0.05.




Figure 4 – Actin polymerisation and α IIb β 3 are required for PECAM-1 mediated internalisation of GPIIb α . Surface expression of GPIIb α on human platelets following (A) PECAM-1 crosslinking or (B) stimulation with CRP-XL in the presence of EGTA (2mM), Cytochalasin D (50 μ M) or Latrunculin (8.5 μ M) was measured by flow cytometry (n=5). To test the involvement of α IIb β 3 in PMI of GPIIb α , platelets from DiYF mice, in which tyrosines 747 and 759 of the cytoplasmic tail of the β 3 integrin were mutated to phenylalanine, underwent (C) PECAM-1 crosslinking or (D) CRP-XL stimulation followed by measurement of surface GPIIb α . The surface expression of (E) α IIb β 3 and (F) GPVI were also measured in these mice and showed no significant difference between the genotypes in resting or CRP-XL stimulated conditions (WT: n=5, Het: n=9, KI: n=5). Asterisks indicate significant difference from isotype control or resting platelets in all cases. *p<0.05, **p<0.01, ***p<0.001




Figure 5 – PECAM-1 mediated internalisation of GPIb α is dependent on AKT/GSK-3/Dynamin signalling. (A) The reduction in the expression of GPIb α on the platelet surface resulting from PECAM-1 crosslinking was measured in the absence or presence of Dynasore (100 μ M) or Dynasore and EGTA together (n=8). (B) A representative image showing phosphorylation of GSK-3 α Ser21 and GSK-3 β Ser9, and total GSK-3 α and β following platelet stimulation by CRP-XL (1 μ g/mL) or PECAM-1 cross-linking. (C) Quantification of Western blots showing phosphorylation of GSK-3 α Ser21 and GSK-3 β Ser9, and total GSK-3 α and β following platelet stimulation by CRP-XL (1 μ g/mL) or PECAM-1 cross-linking (n=3). (D) The reduction in the expression of GPIb α on the platelet surface resulting from PECAM-1 cross-linking was measured in the absence or presence of the GSK-3 inhibitor CHIR99021 (3 μ M), or CHIR99021 (3 μ M) and EGTA (2mM) together (n=3). *p<0.05, **p<0.01, ***p<0.001

