

*Walking in a winter wonderland?
Strategies for Early and Middle
Pleistocene survival in mid-latitude Europe*

Article

Accepted Version

Hosfield, R. ORCID: <https://orcid.org/0000-0001-6357-2805>
(2016) Walking in a winter wonderland? Strategies for Early
and Middle Pleistocene survival in mid-latitude Europe.
Current Anthropology, 57 (5). pp. 653-682. ISSN 0011-3204
doi: 10.1086/688579 Available at
<https://centaur.reading.ac.uk/43354/>

It is advisable to refer to the publisher's version if you intend to cite from the
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1086/688579>

Publisher: University of Chicago Press

All outputs in CentAUR are protected by Intellectual Property Rights law,
including copyright law. Copyright and IPR is retained by the creators or other
copyright holders. Terms and conditions for use of this material are defined in
the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Site	T _{min} (°C)	T _{max} (°C)	Evidence ¹	Age (MIS)	Source
Happisburgh III (Bed E)	-3 – 0	+16 – 18	Coleoptera	21 or 25	2,10
Pakefield (Bed Cii–Ciii)	-6 – +4	+17 – 23	Coleoptera	17 or 19	2,5
Boxgrove (Unit 4c & Freshwater Silt Bed ≈ Units 4b & 4c)	-4 – +4	+15 – 20	Ostracods (MOTR) & Herpetofauna (MCR)	13	2,7,8
Happisburgh I (Organic Mud)	-11 – -3	+12 – 15	Coleoptera	13/15 or 17	2,5
High Lodge (Bed C1)	-4 – +1	+15 – 16	Coleoptera	13	5
Waverley Wood (Channel 2, Organic Mud)	-	+10 – 15	Coleoptera	13 or 15	5,11
Brooksby (Redland's Brooksby Channel)	-10 – +2	+15 – 16	Coleoptera	13 or 15	5
Barnham (Unit 5c; Holl)	-	+17 – 18	Herpetofauna	11	6
Hoxne (Stratum D ⁵ ; Holla ⁶)	-10 – +6	+15 – 19	Coleoptera	11	3,4
Bilzingsleben	-0.5 – +3	+20 – 25	Mollusca & ostracods ⁹	11	9

Table 1: Winter and summer temperature estimates for Early and Middle Pleistocene north-western and north-central European sites. ¹Sensitivity tests on coleoptera-based MCR procedures suggest that winter temperature estimates are usually too warm (Pettitt and White 2012:35); ²Ashton and Lewis 2012 (Pakefield listed as -4 – +6°C); ³Ashton *et al.* 2008a; ⁴Coope 1993; ⁵Coope 2006; ⁶Holman 1998; ⁷Holman 1999; ⁸Holmes *et al.* 2009; ⁹Mania 1995 (the specific source of the palaeo-temperature estimates is not stated, but the fauna includes molluscs and ostracods); ¹⁰Parfitt *et al.* 2010; ¹¹Shotton *et al.* 1993.

Species	Home range ¹	Density ¹	Mobility ¹	Site examples
<i>M. martes</i>	3–82km ²	1/0.8–10km ²	Solitary; not highly territorial; hunting trips upto 28km	Swanscombe (LL) ⁴
<i>F. sylvestris</i>	0.6–3.5km ²	1/0.7–10km ²	Sedentary; nomadic	Boxgrove ³ Swanscombe (LG) ⁴
<i>C. fiber</i>	500m–5.5km (along river)	1.0–1.8/km ²	Family movements within territory	Boxgrove ³ Bilzingsleben ² Hoxne (Beds C & E) ⁵ Swanscombe (LL) ⁴
<i>C. lupus</i>	100–10,000km ² (food-dependent)	1/50–80km ²	Territorial (and correlating with prey migrations)	Bilzingsleben ² Swanscombe (LL/LG) ⁴

Table 2: Fur-bearing animals, with modern distribution data for comparison, documented on northern European Middle Pleistocene sites. Other documented species include: *V. vulpus*; *M. putorius*; *M. erminea*; *M. lutreola*; and *L. lutra*.

¹Macdonald and Barrett 1993 (modern European data; it is fully acknowledged that Early and Middle Pleistocene species' ecology would not have been identical to their modern equivalents); ²Mania and Mania 2005; ³Parfitt 1999; ⁴Schreve 1996; ⁵Stuart *et al.* 1993. Site units: Swanscombe (LL): Lower Loam; Swanscombe (LG): Lower Gravels.

Species	Home range ³	Density ³	Mobility ³	Site examples
<i>C. capreolus</i>	0.05–1km ²	15–25/km ² Solitary/small groups in closed woodland ⁷	Reduced territoriality in winter & congregation (herds up to 30)	Boxgrove ¹ Bilzingsleben ⁴ Hoxne ¹⁰ Swanscombe (LL) ⁸
<i>D. dama</i> ⁶	0.5–2.5km ²	12(?) / km ² Small groups (<7/8) in closed/open woodland ¹²	Habitat use shifts seasonally (e.g. summer in open habitats & autumn → spring in woodlands ⁷)	Barnham ⁵ Bilzingsleben ⁴ High Lodge? ⁹ Hoxne ¹⁰ Swanscombe (LG) ⁸
<i>C. elaphus</i>	0.5–8km ² Smaller upper limits also suggested ²	5–45/km ² Small groups (1–3) in closed woodland ⁷	Summer → winter range migrations up to 6km (e.g. lowland woodlands → open uplands [UK])	Barnham ⁵ Bilzingsleben ⁴ Boxgrove ⁶ High Lodge? ⁹ Hoxne ¹⁰ Schöningen 13-I & 13 II-4 ^{11,12} Swanscombe (LL) ⁸

Table 3: Modern home range, density and mobility data for selected ungulate species, documented on Middle Pleistocene sites. ¹Bello, Parfitt, and Stringer 2009; ²Clutton-Brock, Guinness, and Albon 1982; ³Macdonald and Barrett 1993 (modern European data; it is fully acknowledged that Early and Middle Pleistocene species' ecology would not have been identical to their modern equivalents); ⁴Mania and Mania 2005; ⁵Parfitt 1998; ⁶Parfitt 1999 (notes that the fallow deer's late rut results in males' poor condition during winter); ⁷Putman 1988; ⁸Schreve 1996; ⁹Stuart 1992; ¹⁰Stuart *et al.* 1993; ¹¹Thieme 2005; ¹²Voormolen 2008. Site units: Swanscombe (LL): Lower Loam; Swanscombe (LG): Lower Gravels.

Species	Home range ²	Density ²	Mobility ²	Site examples
<i>C. fiber</i>	500m–5.5km (along river)	1.0–1.8/km ²	Family movements within territory	Bilzingsleben ³ Boxgrove ⁵ Hoxne ⁸ Swanscombe (LL) ⁶
<i>S. scrofa</i>	2–20km ²	ND	Sedentary (if stable env.); ♀ Small herds; ♂ Solitary	Barnham ⁴ Bilzingsleben ³
<i>U. arctos</i>	150–4000km ²	1–190/ 10,000km ²	Solitary; Travel 2– 3.5km/day; Hibernation (with accumulated fat) ¹	Swanscombe (LL/LG) ⁶ Barnham ⁴ Hoxne ⁸
<i>D. bicornis</i>	Few ha–75 sq. km	ND	♀ + young; ♂ Solitary; Resident & local (if resources sufficient)	Barnham ⁴ Bilzingsleben ³ Boxgrove ⁵ Hoxne ⁸ High Lodge ⁷ Swanscombe (LG) ⁶

Table 4: Fat-bearing and/or residential winter animals, with modern distribution data for comparison, documented on Middle Pleistocene sites. ¹Jochim 1981; ²Macdonald and Barrett 1993 (modern European data); ³Mania and Mania 2005; ⁴Parfitt 1998; ⁵Parfitt 1999; ⁶Schreve 1996; ⁷Stuart 1992; ⁸Stuart *et al.* 1993.

Species	Butchery evidence	Sites
<i>Bos or Bison sp.</i>	Marrow extraction & cut-marks (filleting?); Filleting; Cut-marks, defleshing and marrow bone breakage; Dismembering, filleting, defleshing & marrow bone breakage	Barnham ⁴ Boxgrove ⁵ Happisburgh I ² Schöningen 13 II-4 ⁷
<i>C. capreolus</i>	Cut-marks; Defleshing	Boxgrove ³ Happisburgh I ²
<i>C. elaphus</i>	Skinning, dismemberment, filleting & marrow bone breakage; Marrow bone breakage & cut-marks (<u>seasonality data: late Summer → Spring</u>); Skinning, dismemberment & filleting; Cut-mark	Boxgrove ⁵ Hoxne ⁶ Schöningen 13 II-4 ⁷ Westbury ¹
<i>E. ferus</i>	Disarticulation, filleting & marrow bone breakage; Marrow bone breakage & cut-marks; Dismemberment, filleting, boning, defleshing & marrow bone breakage	Boxgrove ⁵ Hoxne ⁶ Schöningen 13 II-4 ⁷
<i>S. hundsheimensis</i>	Disarticulation & filleting; Disarticulation	Boxgrove ⁵ Happisburgh I ²
<i>U. deningeri</i>	Skinning	Boxgrove ⁵

Table 5: Butchery by species and technique, from selected Lower Palaeolithic sites.

¹Andrews and Ghaleb 1999; ²Ashton *et al.* 2008b; ³Bello, Parfitt, and Stringer 2009;

⁴Parfitt 1998; ⁵Parfitt and Roberts 1999; ⁶Stopp 1993; ⁷Voormolen 2008.

Family/Species identified at Hoxne ³	Modern winter foraging species ^{1,2}		
	Species	Habitat	Key Nutrients
<i>Caryophyllaceae</i>	Common chickweed (<i>Stellaria media</i>)	Woodland fringe	Vitamins A, D, B complex, C, and Rutin
	Common mouse-ear chickweed (<i>Cerastium holosteoides</i>)	Grassland	-
<i>Brassicaceae</i> (previously <i>Cruciferae</i>)	Garlic mustard or Jack-by-the-hedge (<i>Alliaria petiolata</i>)	Woodland fringe	Vitamins A, C & E
<i>Ericaceae</i>	Cowberry (<i>Vaccinium vitis-idaea</i>)	Pine forest	Vitamins A, B & C
<i>Apiaceae</i> (or <i>Umbelliferae</i>)	Wild parsnip (<i>Pastinaca sativa</i>)	Grassland	Potassium
<i>T. latifolia</i>	Reed mace/Bulrush (<i>Typha latifolia</i>)	Wetland	Protein & carbohydrate
<i>Urticaceae</i>	Stinging nettle (<i>Urtica dioica</i>)	Woodland & river valley	Protein and vitamin C

Table 6: Plant families identified at Hoxne, with comparison to modern plant species available to winter foragers. ¹Mabey 2012; ²Mears and Hillman 2007; ³Mullenders 1993, table 6.3 & figs. 6.1–6.3.




Figure 1: Comparison of winter temperature ranges for Spanish (Early Pleistocene; EP) and British (Early Pleistocene and Middle Pleistocene; EP & MP) sites. Number of sites calculated according to the temperature ranges for each site (e.g. 7 sites have a T_{min} range which spans -3°C). Spanish site data (Almenara-Casablanca 3; Cal Guardiola; Cúllar Baza 1; Barranca León 5; Fuente Nueva 3; Trinchera Dolina (TD6); Trinchera Elefante (TERc)) from Agusti *et al.* (2009); British site data (Boxgrove, Brooksby, Happisburgh I, Happisburgh III, High Lodge, Hoxne, Pakefield) from Ashton *et al.* 2008a; Ashton and Lewis 2012; Coope 1993, 2006; Holman 1998, 1999; Holmes *et al.* 2009; Mania 1995; Parfitt *et al.* 2010.




Figure 2: Mean winter air temperature data (°C) from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, fig. 5.7 [Stage 3 Warm Phase DJF]). Dashed white line: Modern European coastline.




Figure 3: Summer/winter contrasts in mean air temperature data (°C) from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, appendix 5.1). Dashed white line: Modern European coastline.




Figure 4: Snow depth (cm) data from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, fig. 5.9).
Dashed white line: Modern European coastline.




Figure 5: Number of days with snow cover data from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, fig. 5.9). Dashed white line: Modern European coastline.




Figure 6: Wind chill ($^{\circ}\text{F}$) data from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, appendix 5.1). Dashed white line: Modern European coastline.




Figure 7: Precipitation (mm/day) data from the Stage 3 Project's MIS-3 'warm' simulation (Barron, van Andel, and Pollard 2003, appendix 5.1). Dashed white line: Modern European coastline.




Figure 8: Relationship between effective temperature and average distance/residential move (after Kelly 1995, fig. 4-7). Note the examples (circled) for groups making relatively short mean residential moves in low effective temperature environments (see Kelly 1995:128–130 for details). Effective Temperature (ET) is derived from the mean temperatures ($^{\circ}C$) of the warmest and coldest months (W and C); where $ET = \frac{18W - 10C}{(W - C) + 8}$, and its value varies from 26 (equator) to 8 (poles). High ET values are associated with tropical, non-seasonal environments (in terms of temperature, not precipitation) with long growing seasons. Low ET values are associated with cold, seasonal environments with short growing seasons (Kelly 1995:66–69).




Figure 9: Selected sources of vitamins in Arctic hunter-gatherer diets (data from Hidiroglou *et al.* 2008; Kuhnlein *et al.* 2006). Values per 100g of fresh raw caribou liver (e.g. 1.58mg for Riboflavin) compared against alternative food sources (e.g. raw moose liver [6.51mg] and raw beluga muktuk [0.02mg] for Riboflavin). G&D: growth and development.



Figure 10: A winter residency model.

Supplementary Materials

	<i>H. erectus</i> ¹		<i>H. sapiens</i> ²	
	Kleiber BMR ³	Elevated BMR ⁴	Kleiber BMR ³	Elevated BMR ⁴
Body Mass (kg)	68	68	70	70
Stature (cm)	185	185	177	177
BMR	80.512	92.589	82.282	94.624
Body surface area ⁵	1.900	1.900	1.862	1.862
Human Conductance A ⁶	5.0	5.0	5.0	5.0
Total Conductance A ⁷	9.498	9.498	9.312	9.312
Lower Critical Temperature A (°C) ⁸	28.5	27.3	28.2	26.8
Minimum Sustainable Temperature A (°C) ⁹	11.6	7.8	10.5	6.5
Human Conductance B ¹⁰	4.750	4.750	4.750	4.750
Total Conductance B ⁷	9.023	9.023	8.846	8.846
Lower Critical Temperature B (°C) ⁸	28.1	26.7	27.7	26.3
Minimum Sustainable Temperature B (°C) ⁹	10.2	6.2	9.1	4.9
Human Conductance C ¹¹	2.817	2.817	2.817	2.817
Total Conductance C ⁷	5.351	5.351	5.246	5.246
Lower Critical Temperature C (°C) ⁸	22.0	19.7	21.3	19.0
Minimum Sustainable Temperature C (°C) ⁹	-8.1	-14.9	-10.1	-17.1

Table 1: Lower critical and minimum sustainable ambient temperatures for *H. erectus* and *H. sapiens* (after Aiello and Wheeler 2003, tables 9.1–9.3). ¹*H. erectus* data from KNM-WT 15000 (Ruff 1994); ²*H. sapiens* data from Předmost 3 & 9, Skhul 4 and Grotte des Enfants 4 (Ruff 1994); ³BMR = $3.4 \times \text{mass (kg)}^{0.75}$ (Kleiber 1961); ⁴Elevated BMR = BMR raised by 15% to account for climatic and dietary-induced increases (after Aiello and Wheeler 2003:150); ⁵Body surface area (m²) = $0.00718 \times \text{mass (kg)}^{0.425} \times \text{stature (cm)}^{0.725}$; ⁶Typical human conductance = $5 \text{ W} \cdot \text{m}^{-2} \cdot ^\circ\text{C}^{-1}$; ⁷Total conductance = typical human conductance $\times$ surface area (m²); ⁸Critical temperature (°C) = $37^\circ\text{C} - (\text{BMR}/\text{Total conductance})$; ⁹Minimum sustainable ambient temperature (°C) = $37^\circ\text{C} - ((3 \times \text{BMR})/\text{Total conductance})$; ¹⁰Typical human conductance reduced by 5% to account for hominin muscularity (after Aiello and Wheeler 2003:150); ¹¹Typical human conductance reduced by c. 44% to account for 1 clo of insulation (after Aiello and Wheeler 2003:150). 1 clo is roughly equivalent to the insulation provided by a western business suit.

Palaeoclimate Measure ¹	MIS 3 'warm' interval			Modern data		
	52°N 0°E	45°N 0°E	50°N 10°E	52°N 0°E	45°N 0°E	50°N 10°E
Min. monthly lowest-level air temperature (°C)	-4 – 0	0 – +4	-4 – -8	4 – 8	4 – 8	0 – 4
T _{max} – T _{min} (°C)	12 – 16	12 – 16	16 – 20	9 – 12	9 – 12	12 – 16
Diurnal range of lowest level air temperature (°C)	1 – 2	2 – 3	1 – 2	1 – 2	3 – 4	1 – 2
No. of days/year with snow cover	90 – 120	10 – 30	150 – 180	< 10	< 10	30 – 60
Snow depth, actual (cm)	5 – 10	0 – 5	20 – 50	0 – 5	0	0 – 5
Wind chill (°F)	0 – 10	10 – 20	0 – 10	20 – 50	20 – 50	10 – 20
Precipitation (mm/day)	2 – 3	3 – 5	2 – 3	2 – 3	3 – 5	1.5 – 2
Net primary productivity (gC/m ² /year)	200 – 300	300 – 400	200 – 300	600 – 700	900 – 1000	600 – 700
Annual growing days above 5°C (°C.day)	750 – 1000	1000 – 1500	1000 – 1500	1500 – 2000	2000 – 3000	1500 – 2000
Annual growing days above 0°C (°C.day)	1500 – 2000	2000 – 3000	2000 – 3000	3000 – 4000	4000 – 5000	3000 – 4000

Table 2: Selected palaeoclimate simulation data for three point-specific locations, for an MIS 3 'warm' interval and the present day. Data from Barron, van Andel, and Pollard (2003). ¹Descriptions of palaeoclimate measures from Barron, van Andel, and Pollard (2003:78).