

# *Rhodosporidium toruloides cultivated in NaCl-enriched glucose-based media: adaptation dynamics and lipid production*

Article

Accepted Version

Tchakouteu, S. S., Kopsahelis, N., Chatzifragkou, A. ORCID: <https://orcid.org/0000-0002-9255-7871>, Kalantzi, O., Stoforos, N. G., Koutinas, A. A., Aggelis, G. and Papanikolaou, S. (2017) *Rhodosporidium toruloides cultivated in NaCl-enriched glucose-based media: adaptation dynamics and lipid production*. *Engineering in Life Sciences*, 17 (3). pp. 237-248. ISSN 1618-2863 doi: 10.1002/elsc.201500125 Available at <https://centaur.reading.ac.uk/56639/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/elsc.201500125>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# *Rhodosporidium toruloides* cultivated in NaCl-enriched glucose-based media: adaptation dynamics and lipid production

Sidoine Sadjeu Tchakouteu<sup>a</sup>, Nikolaos Kopsahelis<sup>a</sup>, Afroditi Chatzifragkou,<sup>a,1</sup>, Ourania Kalantzi<sup>a</sup>,  
Nikolaos G. Stoforos<sup>a</sup>, Apostolis A. Koutinas<sup>a</sup>, George Aggelis<sup>b,c</sup>, Seraphim Papanikolaou<sup>a</sup>

<sup>a</sup> Department of Food Science and Technology, Agricultural University of Athens, 75 Iera Odos, 11855 – Athens, Greece

<sup>b</sup> Unit of Microbiology, Department of Biology, Division of Genetics, Cell and Developmental Biology, University of Patras, 26500 – Patras, Greece

<sup>c</sup> Department of Biological Sciences, King Abdulaziz University, 21589 – Jeddah, Saudi Arabia

\*Corresponding author: Dr. S. Papanikolaou, Associate Professor, Laboratory of Food Microbiology and  
Technology, Department of Food Science and Human Nutrition, Agricultural University of Athens, Iera Odos 75 –  
Athens, Greece, e-mail: [spapanik@hua.gr](mailto:spapanik@hua.gr), tel., fax: +30-210-5294700

<sup>1</sup> Present address: Department of Food and Nutritional Sciences, University of Reading, Whiteknights Campus, RG6 6AP – Reading, UK

19 Key-words: NaCl, microbial lipids, pasteurized conditions, fed-batch culture, *Rhodosporidium*  
20 *toruloides*

22 Abbreviations and units: X – total biomass (dry cell weight, DCW), (g/L); L – total cellular lipids, (g/L); Glc – glucose, (g/L);  $Y_{L/X}$  – lipid in DCW (%), w/w;  $Y_{X/Glc}$  – total biomass yield, g of total biomass produced per g of glucose consumed;  $Y_{L/Glc}$  – lipid yield, g of cellular lipids produced per g of glucose consumed;  $q_{Glc}$  – specific glucose uptake rate, g of glucose/g of biomass/h;  $r_{Glc}$  – glucose uptake rate, g/L/h;  $\mu$  – specific glucose uptake rate, g of glucose/g of biomass/h.

28    **Practical application**

29        The yeast strain *Rhodosporidium toruloides* DSM 4444 was found capable of producing sufficient lipid  
30    amounts when grown in media supplemented with NaCl. This particular feature of the strain, combined with its  
31    tolerance against relatively high amounts of glucose, could denote the feasibility of microbial oil production without the  
32    need of stringent sterile conditions. The utilization of agro-industrial residues such as salty or brackish waste-water  
33    streams deriving from fisheries or olives production establishments as fermentation media could reduce the cost of  
34    microbial oil production by the particular yeast, whereas waste bio-remediation could offer an additional environmental  
35    benefit in the process. Likewise, the accomplishment of microbial lipid fermentation by this strain in substrates  
36    supplemented with NaCl and without previous sterilization of the culture medium can further reduce the cost of the  
37    bioprocess.

38

39 **Abstract**

40 In the present report and for the first time in the international literature, the impact of the addition of NaCl  
41 upon growth and lipid production on the oleaginous yeast *Rhodosporidium toruloides* was studied. Moreover, equally  
42 for first time, lipid production by *R. toruloides* was performed under non-aseptic conditions. Therefore, the potentiality  
43 of *R. toruloides* DSM 4444 to produce lipid in media containing several initial concentrations of NaCl with glucose  
44 employed as carbon source was studied. Preliminary batch-flask trials with increasing amounts of NaCl revealed the  
45 tolerance of the strain against NaCl content up to 6.0% (w/v). However, 4.0% (w/v) of NaCl stimulated lipid  
46 accumulation for this strain, by enhancing lipid production up to 71.3% (w/w) per dry cell weight. The same amount of  
47 NaCl was employed in pasteurized batch-flask cultures in order to investigate the role of the salt as bacterial inhibiting  
48 agent. The combination of NaCl and high glucose concentrations was found to satisfactorily suppress bacterial  
49 contamination of *R. toruloides* cultures under these conditions. Batch-bioreactor trials of the yeast in the same media  
50 with high glucose content (up to 150 g/L) resulted in satisfactory substrate assimilation, with almost linear kinetic  
51 profile for lipid production, regardless of the initial glucose concentration imposed. Finally, fed-batch bioreactor  
52 cultures led to the production of 37.2 g/L of biomass, accompanied by 64.5% (w/w) of lipid yield. Lipid yield per unit  
53 of glucose consumed received the very satisfactory value of 0.21 g/g, a value amongst the highest ones in the literature.  
54 The yeast lipid produced contained mainly oleic acid and to lesser extent palmitic and stearic acids, thus constituting a  
55 perfect starting material for “second generation” biodiesel.

56

57 **Introduction**

58 Environmental concerns have driven scientific research towards alternative energy  
59 resources, as means of disengagement from fossil oil [1, 2]. Biodiesel consists one of the major  
60 renewable transportation fuels, deriving by trans-esterification process of long chain fatty acids of  
61 plant or animal origin. However, constant rising demand of biodiesel production competes with the  
62 availability of existing raw materials and as a result, other non-conventional oil resources are  
63 explored, mainly of non-edible nature [2]. In this light, scientific interest on microbial lipids as  
64 alternative source of oil has gain momentum the last decades.

65 Microbial oil production can be carried out by a number of heterotrophic (mostly yeast and  
66 fungi) or phototrophic (algae) organisms which are found to accumulate oil up to 80% of their dry  
67 weight [1, 3]. This lipid, namely single cell oil (SCO), is mainly composed of neutral fractions  
68 (principally triacylglycerols-TAGs and to lesser extent steryl-esters) [4], while these lipid-  
69 accumulating microorganisms are called “oleaginous” [1-4]. It has been well established that when  
70 culture is performed on sugars or similarly metabolized compounds (e.g. polysaccharides, glycerol,  
71 etc), the conditions required to trigger lipid production are met in a culture environment with carbon  
72 excess and (at least) one essential nutrient depletion (usually nitrogen) [1, 3-5]. As indicated, SCOs  
73 could constitute the starting material for the synthesis of the “2<sup>nd</sup>” or the “3<sup>rd</sup> generation” biodiesel  
74 [2, 6]. Nevertheless, and despite the huge upsurge of interest concerning the production of microbial  
75 oils amenable to be converted into biodiesel [2, 4], the first industrial applications related with the  
76 utilization of oleaginous microorganisms referred to the production of specialty (and expensive)  
77 lipids, rarely found in the plant or animal kingdom, like the cocoa-butter substitutes [1-4, 7, 8]. In  
78 any case though, the feasibility of sustainable bioprocess development for SCO production is  
79 determined by the cost of both raw materials and (mainly) the fermentation process [1-3, 9-12].  
80 Yeast strains belonging to the genera *Cryptococcus* sp., *Lipomyces* sp., *Rhodotorula* sp.,  
81 *Rhodosporidium* sp, and *Trichosporon* sp. are among those reported as possible biodiesel producers  
82 [13-19]. Among those, *Rhodosporidium toruloides* Y4 has been reported capable of producing

83 106.5 g/L of biomass containing 67.5% (w/w) of oil, during cultivation in a 15-L bioreactor under  
84 fed-batch mode [20], designated as the highest oil production from the particular strain so far.

85 One important aspect developed in several industrial fermentations, refers to the potential of  
86 the accomplishment of the microbial conversion under non-aseptic conditions, due to obvious  
87 process cost reduction [6]. While this is feasible in fermentations in which inhibiting (for any  
88 contaminant microorganisms) metabolites are accumulated into the production media (e.g. ethanol  
89 during the alcoholic fermentation processes; see Sarris and Papanikolaou [6]), generation of such  
90 extra-cellular metabolites inhibiting contaminant cells is not obvious during the fermentation of  
91 SCO production. Thus, addition of such a “hurdle” compound into the medium should be  
92 considered. Preliminary works have identified the potential of the microorganism *R. toruloides*  
93 DSM 4444 to grow on media containing NaCl (an important inhibiting agent towards contaminant  
94 microorganisms) quantities, while in some previous studies it has been demonstrated that amongst  
95 other yeasts, *Rhodosporidium* sp. strains have been isolated from hyper-saline habitats [13] and,  
96 therefore, could potentially grow on media presenting relatively high salinity. The objective of the  
97 study, thus, was double: to evaluate the performance of this yeast strain on glucose-based media  
98 supplemented with different initial quantities of NaCl and to perform SCO production by this yeast  
99 under non-aseptic conditions, due to the addition of salt into the medium.

100

## 101 **Materials and Methods**

102 i) Microorganism and media. *Rhodosporidium toruloides* DSM 4444, provided by the  
103 DSMZ culture collection (Leibniz, Germany), was maintained on yeast peptone dextrose agar  
104 (YPD), supplemented with malt extract, at T=4 °C and sub-cultured every month in order to  
105 maintain its viability. Pre-cultures of the strain contained glucose, yeast extract and peptone at 10  
106 g/L each. The synthetic medium used had the following salt composition (g/L): KH<sub>2</sub>PO<sub>4</sub>, 7.0;  
107 Na<sub>2</sub>HPO<sub>4</sub>, 2.5; MgSO<sub>4</sub>·7H<sub>2</sub>O, 1.5; CaCl<sub>2</sub>, 0.15; FeCl<sub>3</sub>·6H<sub>2</sub>O, 0.15; ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.02;  
108 MnSO<sub>4</sub>·H<sub>2</sub>O, 0.06. Peptone and yeast extract were used as nitrogen sources in concentrations of

109 0.75 g/L and 0.5 g/L respectively. Unless otherwise stated, cultures were supplemented with NaCl  
110 at concentrations of 0.5, 1.0, 1.5, 2.5, 4.0 and 6.0 % (w/v). Commercial glucose provided by the  
111 “Hellenic Industry of Sugar SA” (Thessaloniki, Greece) was used as carbon source in the  
112 fermentations performed [purity *c.* 95%, w/w, impurities composed of maltose (2%, w/w), malto-  
113 dextrines (0.5%, w/w), water (1.5%, w/w) and salts (1.0%, w/w)]. The initial pH for all media  
114 before and after sterilization (121 °C/ 20 min) was 6.0±0.1. Glucose was added at different  
115 concentrations into the medium before heat sterilization. In all trials, initial glucose (Glc<sub>i</sub>)  
116 concentration was measured after the sterilization. Assay of glucose before and after the  
117 sterilization demonstrated very small glucose destruction (*c.* 2%) due to the heat sterilization.

118 ii) Culture conditions. Batch-flask cultures were conducted in 250-mL conical flasks,  
119 containing 50±1 mL of growth medium, previously sterilized (121 °C/ 20 min) and inoculated with  
120 1 mL of a 24-h exponential pre-culture (*c.* 3×10<sup>7</sup> cells, initial biomass concentration at the flasks at  
121 *c.* 0.12±0.02 g/L). Cultures were performed in an orbital shaker (Lab-Line, Illinois-USA) at an  
122 agitation rate of 185±5 rpm and incubation temperature T=26±1 °C. It was desirable to maintain a  
123 medium pH in a value greater than 5.2, therefore an appropriate volume of KOH (5 M) was  
124 periodically and aseptically added into the flasks when needed, in order to maintain the pH value at  
125 6.0±0.2 [19].

126 Batch-flask cultures were also realized under previously pasteurized media by subjecting the  
127 flasks filled with the fermentation media at 100 °C for 7 min. Then 3 mL of 24-h pre-culture (*c.*  
128 9×10<sup>7</sup> cells, initial biomass concentration at the flasks at ~0.36±0.06 g/L) were used as inoculum. A  
129 Jenway 3020 pH-meter was used for pH-measurements of cultures. Dissolved oxygen (DO)  
130 concentration was determined using a selective electrode (OXI 96, B-SET, Germany). Before  
131 harvesting, the shaker was stopped and the probe was placed into the flask. Then, the shaker was  
132 switched on and the measurement was taken after DO equilibration (usually within 10 min).  
133 Oxygen saturation was kept above 20% (v/v) during all growth phases.

134       Batch-bioreactor experiments were carried out in a 3.5-L bioreactor (Infors HT, Labfors 5),  
135    with a working volume of 2.0 L. A 7.5% (v/v) inoculum was employed using 24-h exponential  
136    yeast pre-culture. The stirrer speed was on cascade mode, automatically varying from 200 to 500  
137    rpm to maintain a dissolved oxygen (DO) concentration above 20% (v/v) of saturation. Aeration  
138    and temperature were maintained at 1.0 vvm and T=26 °C, respectively. The pH was maintained at  
139    6.0±0.1 by automatic addition of 5 M KOH. Fed-batch fermentations were initiated in batch mode  
140    and when glucose concentration was reduced to 10 g/L, a volume of concentrated glucose solution  
141    (60%, w/v) was added in the bioreactor. Samples were taken periodically from the bioreactor  
142    throughout fermentation for subsequent analysis as described in the next section.

143       iii) Analytical methods. The whole content of flasks (c. 50 mL) or bioreactor samples (c. 20  
144    mL) were periodically collected and cells were harvested by centrifugation (9000 × g/15 min at 10  
145    °C) in a Hettich Universal 320R (Germany) centrifuge and washed twice with distilled water.  
146    Biomass (X, g/L) was determined by means of dry cell weight (DCW) (95±2 °C/ 24 h). Consumed  
147    glucose was determined by 3,5-dinitrosalicylic acid (DNS) assay [21]. Total cellular lipid (L, g/L)  
148    was extracted from the dry biomass with a mixture of chloroform/methanol 2/1 (v/v) and was  
149    determined gravimetrically. Cellular lipids were converted to their methyl-esters in a two-step  
150    reaction [19]. FAMEs were analyzed according to Fakas et al. [21] and were identified by reference  
151    to standards. In some of the performed trials, in order to investigate whether glycerol (or other  
152    polyols) was secreted into the culture medium, HPLC analysis [21] of the supernatant obtained after  
153    centrifugation was performed. All experiments were performed in duplicate by using different  
154    inocula. All of the experimental points presented in the tables and the figures are the mean value of  
155    two independent determinations, with standard error ≤10%.

156

## 157    **Results**

158       This study was focused on the evaluation of the *Rhodosporidium toruloides* yeast ability to  
159    grow on media containing glucose as carbon source, supplemented with various amounts of NaCl.

160 Special attention was paid to the evolution of lipid production and the potential effect of NaCl on  
161 yeast metabolism, in batch-flask trials as well as batch- and fed-batch bioreactor experiments.

162 i) Effect of NaCl concentration on *Rhodosporidium toruloides* cultures. The yeast strain *R.*  
163 *toruloides* DSM 4444 was cultivated in batch-flask trials, in media containing 50 g/L glucose  
164 supplemented with increasing NaCl concentrations. Cultures were done under nitrogen-limited  
165 conditions (carbon-to-nitrogen ratio equal to 106 mol/mol) in order to stimulate lipid accumulation.  
166 Moreover, an experiment without NaCl addition was included that served as control. The obtained  
167 results with regard to the impact of NaCl upon the physiological behavior of *R. toruloides* are  
168 depicted in Table 1. Generally, it should be stressed that only at increased initial NaCl concentration  
169 (e.g. NaCl at 60 g/L or higher) some negative effect upon the biomass and lipid produced was  
170 observed (Table 1). Likewise, by taking into consideration the specific growth rate for the  
171 fermentations in the media with different initial salinity imposed, similar  $\mu_{\text{max}}$  values ( $0.09 \pm 0.01 \text{ h}^{-1}$ )  
172 were recorder for all trials performed except for the fermentation presenting the highest initial NaCl  
173 quantity (=60 g/L), in which the  $\mu_{\text{max}}$  value slightly declined (see Table 1). Moreover, analysis of  
174 the supernatant at the end of all trials was performed in order to demonstrate whether secretion of  
175 glycerol or other polyols was performed due to the increasing osmotic pressure into the medium; it  
176 has been seen that up to the threshold of 1.5% (w/v) not any polyols have been identified into the  
177 medium. Thereafter, very small glycerol accumulation into the medium occurred (the maximum  
178 quantity of glycerol, c. 1.5 g/L, was obtained at the trial with initial concentration of NaCl imposed  
179 of 40 g/L). No other polyol was synthesized as response to the osmotic stress situation by *R.*  
180 *toruloides*. On the other hand, for NaCl concentrations varying from 0.5 to 2.5% (w/v), microbial  
181 growth as well as lipid production was maintained in similar levels; specifically, 88-97% of initial  
182 glucose concentration was consumed within 120-168 h of fermentation, whereas biomass  
183 production ranged between 8.2 and 8.9 g/L. In terms of lipid production, cells accumulated 60.6-  
184 62.4% (w/w) of oil. However, initial NaCl concentration of 4.0% (w/v) was found to positively  
185 affect biomass and, specifically, lipid production, yielding 9.4 and 6.7 g/L respectively.

186 Consequently, lipid quantity per DCW increased to 71.3% (w/w). Higher NaCl additions exerted  
187 inhibitory effects on yeast growth, as only 6.3 g/L of biomass were synthesized, with concomitant  
188 impact on lipid production and yield (Table 1). On the other hand, for all the above-mentioned  
189 trials, the yield of total biomass produced per glucose consumed ( $Y_{X/Glc}$ ) was *c.* 0.20 g/g, ranging  
190 between 0.19 and 0.21 g/g.

191 Taking into account the satisfactory performance of the strain at NaCl supplementation of  
192 4.0% (w/v), subsequent batch-flasks cultures were carried out with the same NaCl addition and  
193 increased glucose concentration ( $Glc_i \approx 100$  g/L). In the same manner, a control experiment without  
194 NaCl addition was included. In the absence of NaCl, elevated glucose concentrations prolonged the  
195 course of the fermentation up to 433 h. At that time, *c.* 88% of initial carbon source concentration  
196 ( $\approx 93$  g/L) was finally consumed by the strain, leading to 17.5 g/L of biomass. The  $\mu_{max}$  in the  
197 above-mentioned culture, calculated at the early exponential growth phase by fitting the equation

198  $\ln\left(\frac{X}{X_0}\right) = f(t)$  on the experimental data within this phase, was found to be  $\approx 0.08$  h<sup>-1</sup> (slightly lower

199 than that observed on the respective trial with  $Glc_i = 50$  g/L, potentially due to slight inhibition  
200 exerted by the increased initial concentration of glucose). However, biomass production per  
201 substrate consumption yield ( $Y_{X/Glc}$ ) was  $\approx 0.19$  g/g, the same value as in trials with  $Glc_i = 50$  g/L.  
202 Furthermore, the higher  $Glc_i$  concentration employed and the increased carbon-to-nitrogen ratio  
203 (C/N=211 mol/mol) seemed to enhance lipid production in terms of absolute values, reaching a  
204 maximum SCO production of 8.1 g/L. On the other hand, and despite the significant increase of  
205 lipid in absolute values compared with the respective trial in which  $Glc_i$  was adjusted to *c.* 50 g/L  
206 (see Table 1 entry 1), total lipid in DCW ( $Y_{L/X}$ ) value was lower than the one obtained in the trial  
207 with  $Glc_i = 50$  g/L (*c.* 46.3% against 55.1% w/w). The presence of 4.0% (w/v) NaCl in media with  
208 100 g/L of glucose, extended the fermentation duration to more than 500 h, a time point in which *c.*  
209 80% of the initial glucose was consumed. Biomass synthesis was slightly reduced to 16.1 g/L  
210 compared with the culture with  $Glc_i \approx 100$  g/L and no NaCl addition was performed. Nonetheless,  
211  $Y_{X/Glc}$  and  $\mu_{max}$  values were unaffected ( $= 0.20$  g/g and 0.08 h<sup>-1</sup> respectively) by the addition of salt

212 into the medium. On the other hand, lipid in terms of both absolute (g/L) and relative (% in DCW)  
213 values was noticeably higher compared with the equivalent experiment ( $\text{Glc}_i \approx 100 \text{ g/L}$ ) in which no  
214 NaCl addition occurred ( $L = 9.2 \text{ g/L}$ ,  $Y_{L/X} = 57.1\% \text{ w/w}$ ). Surprisingly enough, *R. toruloides*  
215 accumulated oil in an almost linear manner, whereas shortly after virtual exhaustion of the  
216 assimilable nitrogen from the culture medium (i.e. *c.* 50 h after inoculation) lipid in DCW almost  
217 reached its plateau (see Fig. 1).

218 Table 2 shows the FA profiles of *R. toruloides* cellular lipids, during growth on media with  
219 increasing NaCl concentrations. In every case, the predominant fatty acid of the yeast was oleic  
220 ( $\Delta^9\text{C}18:1$ ), followed by palmitic (C16:0) stearic (C18:0) and linoleic acid ( $\Delta^{9,12}\text{C}18:2$ ). The  
221 implementation of NaCl did not seem to affect the amounts of individual fatty acids in the  
222 composition of the accumulated lipids. On the contrary, the unsaturated nature of lipids increased  
223 during the course of fermentation, as indicated by the SFA/UFA ratio. This fact could be mainly  
224 attributed to the increased amounts of the unsaturated oleic and linoleic acids and the declining  
225 percentage of stearic acid that occurred in the late growth phase.

226 ii) Trials of *Rhodosporidium toruloides* on pasteurized media supplemented with NaCl.

227 Based on evidence of the tolerance against noticeable amounts of NaCl (e.g. 4.0% w/v) shown by  
228 the employed yeast strain, it was decided to further investigate the stability of microbial growth and  
229 lipid production under pasteurized conditions and assess whether the presence of 4.0% (w/v) NaCl  
230 could reduce the probability of culture contamination. It is evident that a successful accomplishment  
231 of SCO production unsterile media can reduce the cost of the process when a scale-up is envisaged.  
232 To this end, batch-flask trials were carried out with *c.* 50 and 100 g/L of glucose as substrate, in  
233 media supplemented with 4% (w/w) NaCl. At  $\text{Glc}_i \approx 50 \text{ g/L}$ , substrate exhaustion occurred around  
234 160 h after inoculation, yielding 11.7 g/L of biomass production (Table 3A). However, lipid  
235 accumulation was lower than in the experiment in which the medium had been previously subjected  
236 to heat sterilization ( $L = 5.9 \text{ g/L}$  against  $6.7 \text{ g/L}$ ). Microscopy observations revealed the presence of  
237 bacterial contamination (rods), accounting for *c.* 8% of the total microbial population. When higher

238 Glc<sub>i</sub> concentrations were employed in pasteurized media, equally some bacterial contamination  
239 occurred (c. 5% of the total microbial population). As in the trial with Glc<sub>i</sub>≈50 g/L, glucose  
240 assimilation in the pasteurized medium was more rapid than in the aseptic fermentation, possibly  
241 due to this contamination. Equally, biomass formation was enhanced in the pasteurized medium in  
242 comparison with the aseptic culture, reaching a DCW value of 17.9 g/L that contained 9.1 g/L of oil  
243 (lipid in DCW of c. 51% w/w) while in the aseptic culture the respective values were for DCW 16.1  
244 g/L and for lipid 9.2 g/L (see Table 3A). The kinetics of biomass produced, lipid accumulated and  
245 glucose assimilated for the trials with Glc<sub>i</sub>≈100 g/L are seen in Fig. 2.

246 Table 4A shows the FA profiles of the produced cellular lipids for the previously  
247 pasteurized media, in which Glc<sub>i</sub> concentration was adjusted to c. 50 and 100 g/L and constant NaCl  
248 quantity added. Despite the fact that the cultures were not axenic (as stated, some contamination by  
249 bacilli existed), the FA composition presented significant similarities with the trials in which  
250 growth and lipid accumulation occurred in previously sterilized media (see and compare Tables 2  
251 and 4A). In any case, lipid produced through the non-aseptic experiments was rich in the FA  
252  $\Delta^9$ C18:1, constituting, thus, a perfect fatty material amenable to be converted into biodiesel [2, 22].

253 iii) Batch-bioreactor cultures of *Rhodosporidium toruloides*. The next step in the  
254 experimental process involved the realization of batch-bioreactor cultures of the yeast with  
255 increasing glucose concentrations and NaCl supplementation, aiming to promote, if possible, lipid  
256 production. Previously sterilized fermentation media containing c. 50, 100 and 150 g/L of glucose  
257 and 4.0% (w/v) NaCl were used in bench top bioreactor cultures. Table 3B summarizes the  
258 quantitative data of *R. toruloides* trials in bioreactor experiments. At 50 g/L of glucose, the strain  
259 exhibited rapid substrate assimilation within 72 h. Biomass production was notably enhanced, as  
260 opposed to the respective batch-flask experiment, yielding 12.7 g/L with 8.1 g/L of oil. Increasing  
261 amounts of carbon source did not seem to drastically negatively affect the microbial metabolism,  
262 while in all cases, it is interesting to indicate that glucose was linearly consumed (Fig. 3). On the  
263 other hand, the more the Glc<sub>i</sub> concentration (and, hence, the initial molar ratio C/N of the medium)

264 increased, the more the glucose consumption rate ( $r_{Glc} = -\frac{\Delta Glc}{\Delta t}$ ) decreased; for  $Glc_i$  adjusted at *c.*  
265 50 g/L,  $r_{Glc}$  was =0.62 g/L/h decreasing to 0.54 g/L/h for  $Glc_i \approx 100$  g/L. Finally, at  $Glc_i \approx 150$  g/L,  
266  $r_{Glc}$  value eventually dropped to 0.34 g/L/h (Fig. 3). During these trials, lipid accumulation process  
267 demonstrated remarkable stability; in accordance with the trial performed in shake flasks, as  
268 depicted in Fig. 3, the evolution of lipids' kinetic profile was almost linear, regardless of the applied  
269 initial glucose concentration. Maximum biomass production was achieved at 150 g/L of glucose  
270 equal to 34.1 g/L, containing 65.1% (w/w) of oil. It is interesting to indicate that under the present  
271 culture conditions, growth was not inhibited by the increment of  $Glc_i$  concentration up to the  
272 threshold of 150 g/L; this assumption can be justified by the fact that the yields  $Y_{X/Glc}$  and  $Y_{L/Glc}$   
273 presented their higher values at the trial in which the concentration of carbon substrate had been  
274 adjusted at *c.* 150 g/L (=0.33 and 0.21 g/g respectively), being clearly the higher ones obtained in  
275 all of the previously performed trials (including fermentations in shake-flasks or bioreactor). In  
276 addition, specifically the yield  $Y_{L/Glc}$  value obtained in the bioreactor experiment with  $Glc_i \approx 150$  g/L  
277 (=0.21 g/g) was a value very close to the maximum achievable one of 0.22-0.23 g per g of  
278 consumed sugar that has been achieved so far in the international literature [1, 3, 11, 21, 23, 24],  
279 suggesting, once more, the absence of inhibitory phenomena of increased  $Glc_i$  concentrations upon  
280 the growth of *R. toruloides* under the present culture conditions.

281 Table 4B shows the FA profiles of *R. toruloides* cellular lipids, during growth on bioreactor  
282 cultures in media presenting increasing  $Glc_i$  concentrations and constant NaCl quantity added. As in  
283 the shake-flask trials, the predominant FA of the yeast was the  $\Delta^9$ C18:1, followed by the C16:0,  
284 C18:0 and  $\Delta^{9,12}$ C18:2. Moreover, as in the case of the trials with the increasing initial NaCl  
285 quantities into the medium, the increment of  $Glc_i$  concentration did not seem to have serious impact  
286 upon the FA composition of the strain, while cellular FAs were slightly more saturated at the  
287 beginning of the fermentation (Table 4B).

288 iv) Fed-batch bioreactor culture of *Rhodosporidium toruloides*. In an attempt to further  
289 investigate lipid production potential of the yeast *R. toruloides* and to reduce the time of the

290 fermentation (as seen in the previous paragraph, the more the  $\text{Glc}_i$  concentration increased, the more  
291 the time of the fermentation rose) fed-batch cultures were performed in bench top bioreactor, in  
292 media containing 4.0% (w/v) NaCl. Trials were initiated batch-wise ( $\text{Glc}_i \approx 50 \text{ g/L}$ ) and when the  
293 glucose level dropped below 10 g/L, a volume of concentrated glucose solution was aseptically  
294 introduced to the culture. In every case, it was desirable to maintain the feeding of glucose to  
295 concentrations  $\leq 50 \text{ g/L}$ , in order to increase the uptake rate of glucose. In the first cycle of the fed-  
296 batch operation (0-72 h), the  $r_{\text{Glc}}$  was  $= 0.63 \text{ g/L/h}$  and the respective specific glucose consumption  
297 rate ( $q_{\text{Glc}}$ ;  $q_{\text{Glc}} = \frac{r_{\text{Glc}}}{X_{\text{average}}}$ ) was  $\approx 0.1 \text{ g/g/h}$ . In the second cycle (72-192 h),  $r_{\text{Glc}}$  was  $\approx 0.40 \text{ g/L/h}$  and  
298  $q_{\text{Glc}}$  was  $\approx 0.06 \text{ g/g/h}$ . In the third cycle (192-272 h)  $r_{\text{Glc}}$  was  $= 0.51 \text{ g/L/h}$  and  $q_{\text{Glc}}$  was  $\approx 0.08 \text{ g/g/h}$ .  
299 During 272 h of the fermentation, feeding pulses were done twice resulting in the total consumption  
300 of c. 127 g/L of glucose (Fig. 4A). Maximum biomass achieved was 37.2 g/L with 64.5% (w/w) of  
301 accumulated oil. ( $L_{\text{max}} \approx 24 \text{ g/L}$ ). Furthermore, overall yields for lipid and biomass production per  
302 consumed substrate in fed-batch process were 0.21 and 0.33 g/g, respectively (Fig. 4B). Compared  
303 to the batch-bioreactor cultures with high initial glucose concentration ( $\text{Glc}_i \approx 150 \text{ g/L}$ ), both biomass  
304 formation and lipid accumulation were slightly improved during the fed-batch culture mode,  
305 whereas the fermentation was accomplished more rapidly in the later case than in the former one,  
306 thus the lipid volumetric productivity achieved in the fed-batch experiment was improved compared  
307 with the batch process presenting high  $\text{Glc}_i$  concentration (0.088 against 0.075 g/L/h).  
308

### 309 **Discussion**

310 The oleaginous nature of *Rhodosporidium toruloides* has been a topic of interest for many  
311 studies in the international literature. Origin of carbon or nitrogen sources, nutrient limitation and  
312 feeding strategy has been assessed for the optimization of SCO production by the particular yeast.  
313 Lipid accumulation by *R. toruloides* has been shown to improve in the presence of organic nitrogen  
314 sources [25, 26]. Although nitrogen limitation has long been recognized as a determinant factor for  
315 *de novo* lipid synthesis in oleaginous microorganisms [1-5], phosphorus- and sulfate-limitation

316 conditions have been also investigated as lipid inducing factors for *R. toruloides* strains [26, 27]. In  
317 terms of carbon sources, strains of the particular yeasts are reported to withstand carbon-rich media  
318 and under certain conditions, to promote high density cell cultures [20]. On the other hand, in  
319 earlier studies, strains of *R. toruloides* have been flask-cultured in media composed of pure stearic  
320 acid or blends of pure stearic acid, glucose and glycerol and tailor-made lipids presenting  
321 similarities with the cocoa-butter had been synthesized [8, 29]. Equally in early studies, strains of  
322 this species had been cultivated on glucose-based media in which Δ9 and Δ12 natural or artificial  
323 desaturase inhibitors had been added into the culture medium in order to suppress the desaturation  
324 reactions inside *R. toruloides* cells, so as finally, again to synthesize lipids presenting compositional  
325 similarities with the cocoa-butter [30, 31]. More recently Zhu et al. [32] have carried out a massive  
326 study based on genomic sequencing of *R. toruloides*, in an attempt to unravel lipid accumulation  
327 process on a genetic level, spent cell mass hydrolysates used as nutrients and spent water from lipid  
328 production process were used as substrates by *R. toruloides* for SCO production [33] while other  
329 species belonging to the genus *Rhodosporidium* (e.g. *R. kratochvilovae*, *R. babjevae*, *R.*  
330 *diobovatum*) have been successfully used as cell factories used for SCO on several waste streams or  
331 low-cost materials [34-36].

332 One major objective of the study was to identify the effect of NaCl addition into the culture  
333 medium upon the process of lipid accumulation of *R. toruloides* DSM 4444. To this end, batch-  
334 flask cultures of the microorganism were performed in media containing increasing amounts of  
335 NaCl up to 6.0% (w/v). The particular strain can be categorized as halotolerant, due to its ability to  
336 grow sufficient in the presence or absence of salt [13, 37]. Furthermore, NaCl concentrations up to  
337 4.0% (w/v) were found to promote optimum growth of the yeast and thus, according to Larsen [38]  
338 the strain can be designated as moderate halotolerant. This feature is commonly encountered in  
339 microorganisms isolated from marine environments, possessing unique adaptation mechanisms in  
340 high salinity conditions [13, 39-42]. A major part of the mechanisms involved in osmotic balance  
341 regulation for halotolerant microorganisms represents the production and accumulation of solutes,

342 such as glycerol, trehalose, mannitol and erythritol, etc [43]. In the current investigation and after a  
343 threshold NaCl value, small (but not negligible) glycerol quantities were found into the medium,  
344 apparently as response to the osmotic stress imposed. Under this optic, the last years there has been  
345 a number of reports in which the polymorphic yeast *Y. lipolytica* has been cultivated in media  
346 composed of high initial concentrations of (pure or biodiesel-derived) glycerol supplemented with  
347 increased initial concentrations of NaCl, and enhancement of production of erythritol, the  
348 concentration of which in some cases reached in indeed very high levels (e.g. >45 g/L or even c.  
349 200 g/L) has been reported when glycerol has been employed as fermentation substrate by wild or  
350 mutant *Y. lipolytica* strains [44-47]. Likewise, besides mannitol and erythritol, due to their high  
351 osmotic tolerance, halotolerant yeasts have been proposed as candidates for bioethanol production  
352 [48], enzyme production [49] as well as xylitol biosynthesis [50]. However, none of these  
353 microorganisms has ever been reported as oleaginous. Surprisingly enough, in the current study the  
354 addition of NaCl not only did not suppress microbial growth, but on the contrary was found to  
355 enhance lipid accumulation, as such was the case in media containing 4.0% (w/v) NaCl. Under  
356 these conditions, lipid accumulation increased to c. 29% compared to the control experiment (see  
357 Table 1). Positive correlation between salt and lipid production has been shown for oleaginous  
358 microalgae strains belonging to the genera of *Dunaliella* sp. and *Nannochloropsis* sp. [51, 52].

359 Another aspect of the study was the realization of microbial lipid production in the presence  
360 of NaCl under pasteurized conditions. In batch-flask trials with 50 or 100 g/L of initial glucose  
361 concentration, signs of bacterial contamination were noted, despite the use of a more concentrated  
362 inoculum. However, bacterial presence was less than 5% of the total microbial population and was  
363 not found to be detrimental for yeast growth and lipid formation. In the case of microbial  
364 conversions, pasteurized grape must has been used for alcoholic fermentation [53], while the effect  
365 of pasteurized whey-based medium on propionic acid production has also been evaluated [54].  
366 Additionally, the application of completely non-aseptic conditions has been tested for microbial  
367 solvent production such as ethanol and 1,3-propanediol [55-58], as means of energy and operation

368 cost reduction. However, in all of the above-mentioned cases (e.g. production of bio-ethanol and  
369 1,3-propanediol), it was the main metabolic compound-target (the bio-alcohol) that was synthesized  
370 rapidly and in high concentrations that hindered the microbial contamination with undesirable  
371 microorganisms, fact that does not happen during the SCO fermentation. As far as the SCO  
372 bioprocess is concerned, in order to perform a relatively successful non-aseptic culture, a “hurdle”  
373 should be added into the medium; i.e. Moustogianni et al. [59] have successfully produced SCO by  
374 using oleaginous Zygomycetes grown on glycerol when essential oils or antibiotics had previously  
375 been added into the fermentation medium. In any case, the current study is one of the first in the  
376 literature that deals with the production of SCO under non-aseptic conditions.

377 During batch-bioreactor trials, the yeast *R. toruloides* exhibited notable tolerance against  
378 high substrate concentrations. The particular strain grew satisfactorily without obvious substrate  
379 inhibition being exerted in media containing up to 150 g/L of glucose, while of importance was the  
380 almost linear profile of lipid accumulation, regardless of the initial substrate concentration  
381 employed into the medium. Tolerance against high substrate (e.g. sugar or glycerol) input is  
382 essential, in order to achieve high-density cultures that have been proved as effective cultivation  
383 strategy in the case of microbial lipid production. In a similar manner, Li et al. [14] demonstrated  
384 that the yeast *R. toruloides* Y4 grew well in media containing glucose up to 150 g/L, a fact directly  
385 correlated with the tolerance of the yeast against osmotic stress. On the other hand, for several  
386 oleaginous yeasts, substrate (e.g. sugar or glycerol) concentrations at c. 100 g/L or even lower (e.g.  
387 60-70 g/L) have been reported as threshold, above which microbial cell growth was repressed  
388 significantly [60-62]. In contrast, oleaginous fungi seem more resistant in high initial substrate  
389 (sugar or glycerol) quantities, since initial concentrations ranging between 80-100 g/L have been  
390 considered as ideal in order to enhance the process of lipid accumulation for the species *Mortierella*  
391 *isabellina*, *Thamnidium elegans* and *Cunninghamella echinulata* [21, 24, 63-65].

392 *R. toruloides* DSM 4444 presented remarkable cell growth and lipid production in both  
393 shake-flask and bioreactor experiments. As indicated in the previous paragraphs, strains of this

394 particular species have been employed already from early studies in order for SCO production to be  
395 performed. Initially (mid 80s) these strains had been employed as microbial cell factories in order to  
396 produce lipids that mimicked the composition of cocoa-butter [2, 4, 8, 29-31]. The last decade, with  
397 the potentiality of the replacement of edible oils by non-conventional fatty substances (e.g. yeast  
398 oils) as starting materials in order for biodiesel precursors to be synthesized has been assessed, *R.*  
399 *toruloides* has been considered as a microorganism of importance amenable to be used in the  
400 conversion of low-cost hydrophilic materials (e.g. lignocellulosic sugars, glucose, sorghum extracts,  
401 glycerol, etc) into SCOs. Characteristic results concerning production of lipid in several  
402 fermentation configurations are depicted in Table 5.

403 In the current investigation and specifically in the bioreactor experiments, conversion yields  
404 of the order of *c.* 0.21 g of lipid produced per g of glucose consumed have been seen. The  
405 stoichiometry of glucose (and similar sugars such as lactose, fructose, etc) metabolism indicates that  
406 about 1.1 moles of acetyl-CoA are generated from 100 g of glucose [1, 3, 5]. Assuming that all the  
407 acetyl-CoA produced is channeled towards lipid synthesis, the maximum theoretical yield of SCO  
408 produced per glucose consumed is around 0.32 g/g [3, 5]. This value concerning the fermentation of  
409 xylose ranges between 0.31-0.34 g/g, while with reference to glycerol, the maximum theoretical  
410 yield of SCO is around 0.30 g/g [1, 3]. However, even under ideal conditions for SCO production  
411 (e.g. highly aerated bioreactor cultures) lipid yield on glucose consumed can rarely be higher than  
412 0.22-0.23 g/g [5, 11, 23]. It may be assumed therefore that in the current investigation one of the  
413 highest conversion yields of SCO produced per sugar consumed has been achieved. As previously  
414 indicated [5, 23] cultivation in highly aerated bioreactors was considered as an important  
415 prerequisite in order to achieve such high conversion yields. However, in some cases, in shake-flask  
416 experiments equally high lipid yields can be achieved; in trials with *T. elegans* grown on sucrose in  
417 shake flasks, the conversion yield of lipid produced per sugar consumed was *c.* 0.24 g/g, while  
418 utilization of other sugars (glucose or fructose) equally resulted in exceptional conversion yields,  
419 i.e. >0.20 g/g [24]. Likewise, maximum conversion yields of the same magnitude compared with

420 growth of *T. elegans* on sucrose (c. 0.23 g/g) have been reported for *Cunninghamella echinulata*  
421 cultivated on xylose in shake-flask experiments [21].

422 Fatty acid analysis was carried out in lipids produced during *R. toruloides* batch-flask  
423 cultures with increasing NaCl content. The main fatty acids detected in yeast oil composition were  
424 oleic ( $\Delta^9$ C18:1), palmitic (C16:0), linoleic ( $\Delta^9,12$ C18:2) as well as stearic acid (C18:0). NaCl  
425 presence did not affect the concentration of individual fatty acids, whereas microbial lipid became  
426 generally more unsaturated, during lipid accumulation phase. This is attributed to the increase of  
427 oleic acid as major constituent of accumulated triglycerides [61]. The distribution of *R. toruloides*  
428 fatty acids is similar to oil profiles obtained by other oleaginous yeasts [10, 17-19, 75]. Recent  
429 studies have shown the suitability of microbial oil as starting material for biodiesel production,  
430 through its direct transformation from microbial biomass [21].

431

### 432 **Acknowledgements**

433 1) State Scholarship Foundation (Athens - Greece) is gratefully acknowledged for the scholarship  
434 attributed to Sidoine Sadjeu Tchakouteu; 2) Financial support was attributed by the research project  
435 entitled “Development of novel bioprocesses for the production of biofuels from food industry  
436 waste streams” (Acronym: “Nutri-Fuel”) (09SYN-32-621), implemented within the National  
437 Strategic Reference Framework (NSRF) 2007-2013 and co-financed by National (Greek Ministry of  
438 Higher Education and Religious Affairs - General Secretariat of Research and Technology) and  
439 Community (E.U. - European Social Fund) Funds.

440

441 The authors have declared no conflict of interest.

442

### 443 **References**

444 [1] Papanikolaou, S. and Aggelis, G., Lipids of oleaginous yeasts. Part I: Biochemistry of single cell oil production. Eur.  
445 J. Lipid Sci. Technol. 2011, 113, 1031-1051.  
446 [2] Papanikolaou, S. and Aggelis, G., Lipids of oleaginous yeasts. Part II: Technology and potential applications. Eur. J.  
447 Lipid Sci. Technol. 2011, 113, 1052-1073.

448 [3] Ratledge, C., Biochemistry, stoichiometry, substrates and economics. In: Moreton, R.S. (Ed.), Single Cell Oil,  
449 Longman Scientific & Technical, Harlow (UK) 1988, pp 33-70.

450 [4] Papanikolaou, S. and Aggelis, G., *Yarrowia lipolytica*: A model microorganism used for the production of tailor-  
451 made lipids. *Eur. J. Lipid Sci. Technol.* 2010, 112, 639-654.

452 [5] Ratledge, C. and Wynn, J., The biochemistry and molecular biology of lipid accumulation in oleaginous  
453 microorganisms. *Adv. Appl. Microbiol.* 2002, 51, 1-51.

454 [6] Sarris, D. and Papanikolaou, S., Biotechnological production of ethanol: Biochemistry, processes and technologies.  
455 Eng. Life Sci. 2016, in press, DOI: 10.1002/elsc.201400199.

456 [7] Papanikolaou, S., Chevalot, I., Komaitis, M., Aggelis, G. and Marc, I., Kinetic profile of the cellular lipid  
457 composition in an oleaginous *Yarrowia lipolytica* capable of producing a cocoa-butter substitute from industrial  
458 fats. *Antonie Van Leeuwenhoek* 2001, 80, 215-224.

459 [8] Gierhart, D.I., US Patent 4485, 173, 1984.

460 [9] Xu, J., Zhao, X., Wang, W., Du, W. and Liu, D., Microbial conversion of biodiesel byproduct glycerol to  
461 triacylglycerols by oleaginous yeast *Rhodosporidium toruloides* and the individual effect of some impurities on  
462 lipid production. *Biochem. Eng. J.* 2012, 65, 30-36.

463 [10] Sestric, R., Munch, G., Cicek, N., Sparling, R. and Levin, D.B., Growth and neutral lipid synthesis by *Yarrowia*  
464 *lipolytica* on various carbon substrates under nutrient-sufficient and nutrient-limited conditions. *Bioresour.*  
465 *Technol.* 2014, 164, 41-46.

466 [11] Ykema, A., Verbree, E.C., Kater, M.M. and Smit, H., Optimization of lipid production in the oleaginous yeast  
467 *Apotrichum curvatum* in whey permeate. *Appl. Microbiol. Biotechnol.* 1988, 29, 211-218.

468 [12] Koutinas, A., Chatzifragkou, A., Kopsahelis, N., Papanikolaou, S. and Kookos, I.K., Design and techno-economic  
469 evaluation of microbial oil production as a renewable resource for biodiesel and oleochemical production. *Fuel*  
470 2014, 116, 566-577.

471 [13] Butinar, L., Santos, S., Spencer-Martins, I., Oren, A. and Gunde-Cimerman, N., Yeast diversity in hypersaline  
472 habitats. *FEMS Microbiol. Lett.* 2005, 244, 229-234.

473 [14] Li, Y.H., Liu, Bo., Zhao, Z.B. and Bai, F.W., Optimization of culture conditions for lipid production by  
474 *Rhodosporidium toruloides*. *Chin. J. Biotechnol.* 2006, 22, 650-656.

475 [15] Matsakas, L., Bonturi, N., Alves Miranda, E., Rova, U. and Christakopoulos, P., High concentrations of dried  
476 sorghum stalks as a biomass feedstock for single cell oil production by *Rhodosporidium toruloides*. *Biotechnol.*  
477 *Biofuel.* 2015, 8, 6.

478 [16] Galafassi, S., Cucchetti, D., Pizza, F., Franzosi, G., et al., Lipid production for second generation biodiesel by the  
479 oleaginous yeast *Rhodotorula graminis*. *Bioresour. Technol.* 2012, 111, 398-403.

480 [17] Tsakona, S., Kopsahelis, N., Chatzifragkou, A., Papanikolaou, S. et al., Formulation of fermentation media from  
481 flour-rich waste streams for microbial lipid production by *Lipomyces starkeyi*. *J. Biotechnol.* 2014, 189, 36-45.

482 [18] Leiva-Candia, D.E., Tsakona, S., Kopsahelis, N., García, I.L. et al., Biorefining of by-product streams from  
483 sunflower-based biodiesel production plants for integrated synthesis of microbial oil and value-added co-products.  
484 *Bioresour. Technol.* 2015, 190, 57-65.

485 [19] Tchakouteu, S.S., Kalantzi, O., Gardeli, C., Koutinas, A.A., et al., Lipid production by yeasts growing on  
486 biodiesel-derived crude glycerol: strain selection and impact of substrate concentration on the fermentation  
487 efficiency. *J. Appl. Microbiol.* 2015, 118, 911-927.

488 [20] Li, Y.H., Zhao, Z.B. and Bai, F.W., High-density cultivation of oleaginous yeast *Rhodosporidium toruloides* Y4 in  
489 fed-batch culture. *Enzyme Microb. Technol.* 2007, 41, 312-317.

490 [21] Fakas, S., Papanikolaou, S., Batsos, A., Galiotou Panayotou, M. et al., Evaluating renewable carbon sources as  
491 substrates for single cell oil production by *Cunninghamella echinulata* and *Mortierella isabellina*. *Biomass*  
492 *Bioenerg.* 2009, 33, 573-580.

493 [22] Papanikolaou, S. and Aggelis, G., Biotechnological valorization of biodiesel derived glycerol waste through  
494 production of single cell oil and citric acid by *Yarrowia lipolytica*. *Lipid Technol.* 2009, 21, 83-87.

495 [23] Ratledge, C. and Cohen, Z., Microbial and algal lipids: Do they have a future for biodiesel or as commodity oils?  
496 *Lipid Technol.* 2008, 20, 155-160.

497 [24] Papanikolaou, S., Diamantopoulou, P., Chatzifragkou, A., Philippoussis, A. and Aggelis, G., Suitability of low-cost  
498 sugars as substrates for lipid production by the fungus *Thamnidium elegans*. *Energy Fuel.* 2010, 24, 4078-4086.

499 [25] Evans, C.T. and Ratledge, C., Influence of nitrogen metabolism on lipid accumulation in oleaginous yeasts. *J. Gen.*  
500 *Microbiol.* 1984, 130, 1693-1704.

501 [26] Evans, C.T. and Ratledge, C., Influence of nitrogen metabolism on lipid accumulation by *Rhodosporidium*  
502 *toruloides* CBS 14. *J. Gen. Microbiol.* 1984, 130, 1705-1710.

503 [27] Wu, S., Hu, C., Jin, C., Zhao, X. and Zhao, Z.B., Phosphate-limitation mediated lipid production by  
504 *Rhodosporidium toruloides*. *Bioresour. Technol.* 2010, 101, 6124-6129.

505 [28] Wu, S., Zhao, X., Shen, H., Wang, Q. and Zhao, Z.B., Microbial lipid production by *Rhodosporidium toruloides*  
506 under sulfate-limited conditions. *Bioresour. Technol.* 2011, 102, 1803-1807.

507 [29] Gierhart, D.I., US Patent 4485, 172, 1984.

508 [30] Moreton, R.S., Modification of fatty acid composition of lipid accumulating yeasts with cyclopropene fatty acid  
509 desaturase inhibitors. *Appl. Microbiol. Biotechnol.* 1985, 22, 41-45.

510 [31] Moreton, R.S. and Clode, M., US Patent 4778 630, 1988.

511 [32] Zhu, Z., Zhang, S., Liu, H., Shen, H. et al., A multi-omic map of the lipid-producing yeast *Rhodosporidium*  
512 *toruloides*. *Nature Commun.* 2012, 3, 1112.

513 [33] Yang, X., Jin, G., Gong, Z., Shen, H. et al., Recycling microbial lipid production wastes to cultivate oleaginous  
514 yeasts. *Bioresour. Technol.* 2015, 175, 91-96.

515 [34] Patel, A., Pravez, M., Deeba, F., Pruthi, V. et al., Boosting accumulation of neutral lipids in *Rhodosporidium*  
516 *kratochvilovae* HIMPA1 grown on hemp (*Cannabis sativa* Linn) seed aqueous extract as feedstock for biodiesel  
517 production. *Bioresour. Technol.* 2014, 165, 214-222.

518 [35] Patel, A., Sindhu, D.K., Arora, N., Singh, R.P. et al., Biodiesel production from non-edible lignocellulosic biomass  
519 of *Cassia fistula* L. fruit pulp using oleaginous yeast *Rhodosporidium kratochvilovae* HIMPA1. *Bioresour.*  
520 *Technol.* 2015, 197, 91-98.

521 [36] Munch, G., Sestric, R., Sparling, R., Levin, D.B. and Cicek, N., Lipid production in the under-characterized  
522 oleaginous yeasts, *Rhodosporidium babjevae* and *Rhodosporidium diobovatum*, from biodiesel-derived waste  
523 glycerol. *Bioresour. Technol.* 2015, 185, 49-55.

524 [37] Margesin, R. and Schinner, F., Biodegradation and bioremediation of hydrocarbons in extreme environments.  
525 *Appl. Microbiol. Biotechnol.* 2001, 56, 650-663.

526 [38] Larsen, H., Halophilic and halotolerent microorganisms: an overview historical perspective. *FEMS Microbiol. Rev.*  
527 1986, 39, 3-7.

528 [39] Zaky, A.S., Tucker, G.A., Daw, Z.Y. and Du, C., Marine yeast isolation and industrial application. *FEMS Yeast*  
529 *Res.* 2014, 14, 813-825.

530 [40] Kutty, S.N. and Philip, R., Marine yeasts –a review. *Yeast* 2008, 25, 465-483.

531 [41] Prista C., Loureiro-Dias M.C., Montiel, V., García, R. and Ramos, J., Mechanisms underlying the halotolerant way  
532 of *Debaryomyces hansenii*. FEMS Yeast Res. 2005, 5, 693-701.

533 [42] Rengpipat, S., Lowe, S.E. and Zeikus J.K., Effect of extreme salt concentrations on the physiology and  
534 biochemistry of *Halobacteroides acetoethylicus*. J. Bacteriol. 1988, 170, 3065-3071.

535 [43] Breuer, U. and Harms, H., *Debaryomyces hansenii*-an extremophilic yeast with biotechnological potential. Yeast  
536 2006, 23, 415-437.

537 [44] Tomaszewska, L., Rywińska, A. and Gładkowski, W., Production of erythritol and mannitol by *Yarrowia lipolytica*  
538 yeast in media containing glycerol. J. Ind. Microbiol. Biotechnol. 2012, 39, 1333-1343.

539 [45] Tomaszewska, L., Rakicka, M., Rymowicz, W., Rywinska, A., A comparative study on glycerol metabolism to  
540 erythritol and citric acid in *Yarrowia lipolytica* yeast cells. FEMS Yeast Res. 2014, 14, 966-976.

541 [46] Rywińska, A., Juszczak, P., Wojtowicz, M., Robak, M. et al., Glycerol as a promising substrate for *Yarrowia*  
542 *lipolytica* biotechnological applications. Biomass Bioenerg. 2013, 48, 148-166.

543 [47] Rywińska, A., Tomaszewska, L. and Rymowicz, W., Erythritol biosynthesis by *Yarrowia lipolytica* yeast under  
544 various culture conditions. Afr. J. Microbiol. Res. 2013, 7, 3511-3516.

545 [48] Obara, N., Ishida, M., Hamada-Sato, N. and Urano, N., Efficient bioethanol production from scrap paper shredder  
546 by a marine *Saccharomyces cerevisiae* derived C-19. Studies Sci. Technol. 2012, 1, 127-132.

547 [49] Chen, L., Chi, Z.M., Chi, Z. and Li, M., Amylase production by *Saccharomyces fibuligera* A11 in solid-state  
548 fermentation for hydrolysis of Cassava starch. Appl. Biochem. Biotechnol. 2010, 162, 252-263.

549 [50] Misra, S., Raghuwanshi, S. and Saxena, R.K., Fermentation behavior of an osmotolerant yeast *D. hansenii* for  
550 xylitol production. Biotechnol. Prog. 2012, 28, 1457-1465.

551 [51] Takagi, M., Karseno, and Yoshida, T., Effect of salt concentration on intracellular accumulation of lipids and  
552 triacylglyceride in marine microalgae *Dunaliella* cells. J. Biosci. Bioeng. 2006, 101, 223-226.

553 [52] Bartley, M.L., Boeing, W.J., Corcoran, A.A., Holguin, F.O. and Schaub, T., Effects of salinity on growth and lipid  
554 accumulation of biofuel microalga *Nannochloropsis salina* and invading organisms. Biomass Bioenerg. 2013, 54,  
555 83-88.

556 [53] Sarris, D., Kotseridis, Y., Linga, M., Galiotou-Panayotou, M., Papanikolaou, S., Enhanced ethanol production,  
557 volatile compound biosynthesis and fungicide removal during growth of a newly isolated *Saccharomyces*  
558 *cerevisiae* strain on enriched pasteurized grape musts. Eng. Life Sci. 2009, 9, 29-37.

559 [54] Anderson, T.M., Bodie, E.A., Goodman, N. and Schwartz, R.D., Inhibitory Effect of autoclaving whey-based  
560 medium on propionic acid production by *Propionibacterium shermanii*. Appl. Environ. Microbiol. 1986, 51, 427-  
561 428

562 [55] Chatzifragkou, A., Papanikolaou, S., Dietz, D., Doulgeraki, A.I. et al., Production of 1,3-propanediol by  
563 *Clostridium butyricum* growing on biodiesel-derived crude glycerol through a non-sterilized fermentation process.  
564 Appl. Microbiol. Biotechnol. 2011, 91, 101-112.

565 [56] Metsoviti, M., Zeng, A.P., Koutinas A.A. and Papanikolaou S., Enhanced 1,3-propanediol production by a newly  
566 isolated *Citrobacter freundii* strain cultivated on biodiesel-derived waste glycerol through sterile and non-sterile  
567 bioprocesses J. Biotechnol. 2013, 163, 408-418.

568 [57] Dietz, D. and Zeng, A.P. Efficient production of 1,3-propanediol from fermentation of crude glycerol with mixed  
569 cultures in a simple medium. Bioproc. Biosyst. Eng. 2014, 37, 225-233.

570 [58] Sarris, D., Matsakas, L., Aggelis, G., Koutinas, A.A. and Papanikolaou, S., Aerated vs non-aerated conversions of  
571 molasses and olive mill wastewaters blends into bioethanol by *Saccharomyces cerevisiae* under non-aseptic  
572 conditions. Ind. Crops Prod. 2014, 56, 83-93.

573 [59] Moustogianni, A., Bellou, S., Triantaphyllidou, I.E. and Aggelis, G., Feasibility of raw glycerol conversion into  
574 single cell oil by zygomycetes under non-aseptic conditions. *Biotechnol. Bioeng.* 2015, 112, 827-831.

575 [60] Zhang, J., Fang, X., Zhu, X.L, Li, Y. et al., Microbial lipid production by the oleaginous yeast *Cryptococcus*  
576 *curvatus* O3 grown in fed-batch culture. *Biomass Bioenerg.* 2011, 35, 1906-1911.

577 [61] Meesters P.A.E.P., G.N.M. Huijberts, G. Eggink. Hight-cell-density cultivation of the lipid accumulating yeast  
578 *Cryptococcus curvatus* using glycerol as a carbon source. *Appl Microbiol. Biotechnol.* 1996, 45:575-579.

579 [62] Liang, Y.N., Cui, Y., Trushenski, J. and Blackburn, J.W., Converting crude glycerol derived from yellow grease to  
580 lipids through yeast fermentation. *Bioresour. Technol.* 2010, 101, 7581-7586.

581 [63] Papanikolaou, S., Komaitis, M. and Aggelis, G., Single cell oil (SCO) production by *Mortierella isabellina* grown  
582 on high-sugar content media. *Bioresour. Technol.* 2004, 95, 287-291.

583 [64] Chatzifragkou, A., Makri, A., Belka, A., Bellou, S., et al., Biotechnological conversions of biodiesel derived waste  
584 glycerol by yeast and fungal species. *Energy*, 2011, 36, 1097-1108.

585 [65] Zikou, E., Chatzifragkou, A., Koutinas, A.A. and Papanikolaou, S., Evaluating glucose and xylose as cosubstrates  
586 for lipid accumulation and  $\gamma$ -linolenic acid biosynthesis of *Thamnidium elegans*. *J. Appl. Microbiol.* 2013, 114,  
587 1020-1032.

588 [66] Moreton, R.S., Physiology of lipid accumulating yeasts, in: Moreton, R.S. (Ed.), *Single Cell Oil*, Longman  
589 Scientific & Technical, Harlow (UK) 1988, pp. 1-32.

590 [67] Zhao, X., Wu, S., Hu, C., Wang, Q. et al., Lipid production from Jerusalem artichoke by *Rhodosporidium*  
591 *toruloides* Y4. *J. Ind. Microbiol. Biotechnol.* 2010, 37, 581-585.

592 [68] Wiebe, M.G, Koivuranta, K., Penttilä, M. and Ruohonen, L., Lipid production in batch and fed-batch cultures of  
593 *Rhodosporidium toruloides* from 5 and 6 carbon carbohydrates. *BMC Biotechnol.* 2012, 12: 26.

594 [69] Uçkun Kiran, E., Trzcinski, A. and Webb, C., Microbial oil produced from biodiesel by-products could enhance  
595 overall production. *Bioresour. Technol.* 2013, 129, 650-654.

596 [70] Wang, Z.P., Fu, W.J., Xu, H.M. and Chi, Z.M., Direct conversion of inulin into cell lipid by an inulinase-producing  
597 yeast *Rhodosporidium toruloides* 2F5. *Bioresour. Technol.* 2014, 161, 131-136.

598 [71] Bonturi, N., Matsakas, L., Nilsson, R., Christakopoulos, P. et al., Single Cell Oil producing yeasts *Lipomyces*  
599 *starkeyi* and *Rhodosporidium toruloides*: Selection of extraction strategies and biodiesel property prediction.  
600 *Energies* 2015, 8, 5040-5052.

601 [72] Bommareddy, R.R., Sabra, W., Maheshwari, G. and Zeng, A.P., Metabolic network analysis and experimental  
602 study of lipid production in *Rhodosporidium toruloides* grown on single and mixed substrates. *Microb. Cell Fact.*  
603 2015, 14, 36.

604 [73] Shen, H., Gong, Z., Yang, X., Jin, J. et al., Kinetics of continuous cultivation of the oleaginous yeast  
605 *Rhodosporidium toruloides*. *J. Biotechnol.* 2013, 168, 85-89.

606 [74] Yang, X., Jin, G., Gong, Z., Shen, H. et al., Recycling biodiesel-derived glycerol by the oleaginous yeast  
607 *Rhodosporidium toruloides* Y4 through the two-stage lipid production process. *Biochem. Eng. J.* 2014, 91, 86-91.

608 [75] Zhao, X., Kong, X., Hua, Y., Feng, B. and Zhao, Z.B., Medium optimization for lipid production through co-  
609 fermentation of glucose and xylose by the oleaginous yeast *Lipomyces starkeyi*. *Eur. J. Lipid Sci. Technol.* 2008,  
610 110, 405-412.

611

612

613 Table 1

614 Quantitative data of *Rhodosporidium toruloides* DSM 4444 originated from kinetics in media with six different initial  
615 NaCl concentrations with the same initial glucose concentration (50 g/L). Representation consumed glucose (Glc<sub>cons</sub>,  
616 g/L), produced biomass (X, g/L), lipid content (L, g/L) and lipid in dry weight (Y<sub>L/X</sub>, % w/w) when the maximum  
617 quantity of lipid (in g/L) was achieved. The maximum specific growth ( $\mu_{\max}$ , h<sup>-1</sup>), was calculated for every of the trials

618 performed by fitting the equation  $\ln\left(\frac{X}{X_0}\right) = f(t)$  within the early exponential growth phase of the respective culture.

619 Culture conditions: growth in 250-ml flasks at 185 rpm, initial pH=6.0±0.1, DO>20% (v/v), incubation temperature  
620 T=26 °C. Two lots of independent cultures were conducted by using different inocula. In all of the determinations,  
621 standard error calculated was less that 15%.

622

Entry	NaCl (%, w/v)	Time (h)	$\mu_{\max}$ (h <sup>-1</sup> )	Glc <sub>cons</sub> (g/L)	X (g/L)	L (g/L)	Y <sub>L/X</sub> (%, w/w)
1	0.0	168	0.091	48.8	8.9	4.9	55.1
2	0.5	168	0.088	43.6	8.7	5.4	62.1
3	1.0	144	0.101	48.7	8.2	5.1	62.2
4	1.5	144	0.093	44.5	8.5	5.3	62.4
5	2.5	120	0.084	47.9	8.9	5.4	60.6
6	4.0	192	0.090	48.6	9.4	6.7	71.3
7	6.0	168	0.070	30.0	6.3	2.8	44.4

623

624

625 Table 2

626 Fatty acid composition of cellular lipids of *Rhodosporidium torolooides* DSM 4444 on glucose-based media (Glc<sub>i</sub>=50  
 627 g/L), containing various NaCl concentrations. Very early (VE) growth phase is that in which the incubation time is  
 628 between 20-30 h. Early (E) growth phase is that in which the incubation time is between 50-70 h. Late growth phase is  
 629 that in which incubation time is c. 150 h. Culture conditions as in Table 1.

630

Fatty acid composition (%), w/w											*SFA/ UFA
NaCl (%, w/v)	Growth phase	C14:0	C16:0	$\Delta^9$ C16:1	C18:0	$\Delta^9$ C18:1	$\Delta^9,12$ C18:2	$\Delta^9,12,15$ C18:3	C20:0	C22:0	*SFA/ UFA
0.0	VE	1.1	28.2	0.8	11.5	48.6	7.7	2.1	-	-	0.69
	E	0.8	24.1	1.5	9.9	51.2	9.1	2.4	-	-	0.54
	L	0.6	22.2	1.4	7.1	54.5	11.5	3.0	-	-	0.43
0.5	VE	1.4	27.5	1.2	10.0	49.0	8.6	2.1	-	-	0.68
	E	1.2	25.5	0.8	12.5	48.8	7.8	2.3	0.3	0.7	0.67
	L	1.1	24.6	1.2	10.0	49.0	10.4	2.8	0.6		0.57
1.0	VE	1.6	23.8	0.9	16.7	45.7	6.6	-	-	4.3	0.87
	E	1.6	25.5	1.1	10.9	49.3	8.1	2.3	0.3	0.2	0.63
	L	1.0	23.3	0.8	8.6	52.2	11.2	2.2	0.2	0.2	0.50
1.5	VE	1.1	26.8	-	10.0	51.3	8.6	1.9	-	-	0.61
	E	1.2	25.8	0.5	11.2	50.7	7.3	2.1	0.3	0.5	0.56
	L	0.9	24.5	1.2	9.8	51.5	8.0	2.6	-	-	0.56
2.5	VE	1.2	26.9	1.2	10.5	51.3	7.0	1.6	-	-	0.63
	E	1.1	25.8	0.7	9.6	50.5	9.0	2.3	0.2	0.3	0.59
	L	0.9	24.0	1.5	9.0	53.0	8.0	2.7	-	-	0.52
4.0	VE	1.4	23.4	0.4	15.5	47.2	9.7	2.6	0.4	0.4	0.68
	E	1.2	26.3	0.4	10.3	50.4	8.2	2.4	-	0.5	0.62
	L	0.8	25.5	1.5	8.5	51.8	8.9	2.8	-	-	0.54

631 \*Ratio of saturated to unsaturated fatty acids

632

633

634 Table 3

635 Quantitative data of *Rhodosporidium toruloides* DSM 4444 originated from kinetics in shake-flask experiments in  
 636 sterilized and pasteurized media, supplemented with 4.0% (w/v) NaCl and 50 and 100 g/L initial glucose concentration  
 637 (A) and from kinetics in media containing 50, 100 and 150 g/L of glucose and 4% (w/v) NaCl in batch-bioreactor  
 638 experiments (B). Representation of initial glucose (Glc<sub>i</sub>, g/L), consumed glucose (Glc<sub>cons</sub>, g/L), produced biomass (X,  
 639 g/L), produced lipid (L, g/L), lipid in dry weight (%), lipid yield per consumed substrate (Y<sub>L/Glc</sub>, g/g) and biomass  
 640 yield per consumed substrate (Y<sub>X/Glc</sub>, g/g). Culture conditions for the shake flasks: growth in 250-mL flasks at 185 rpm,  
 641 initial pH=6.0±0.1, DO>20% (v/v), incubation temperature T=26 °C; for the bioreactor: agitation speed 200-500 rpm,  
 642 pH=6.0±0.1, DO>20% (v/v), temperature T=26 °C. Two lots of independent cultures were conducted by using different  
 643 inocula. In all of the determinations, standard error calculated was less than 10%.

644

645 A)

Culture mode and heat-treatment	Glc <sub>i</sub> (g/L)	Time (h)	Glc <sub>cons</sub> (g/L)	X (g/L)	L (g/L)	Y <sub>L/X</sub> (%,w/w)	Y <sub>L/Glc</sub> (g/g)	Y <sub>X/Glc</sub> (g/g)
Flasks, sterilized	≈50	192	48.6	9.4	6.7	71.3	0.14	0.19
Flasks, pasteurized	≈50	160	48.0	11.7	5.9	50.4	0.12	0.24
Flasks, sterilized	≈100	505	93.0	16.1	9.2	57.1	0.10	0.18
Flasks, pasteurized	≈100	311	80.0	17.9	9.1	50.8	0.11	0.22

646 B)

Culture mode	Glc <sub>i</sub> (g/L)	Time (h)	Glc <sub>cons</sub> (g/L)	X (g/L)	L (g/L)	Y <sub>L/X</sub> (%,w/w)	Y <sub>L/Glc</sub> (g/g)	Y <sub>X/Glc</sub> (g/g)
Batch-bioreactor	≈50.0	72	44.5	12.7	8.1	63.8	0.18	0.29
	≈100.0	160	90.9	25.2	14.2	56.3	0.16	0.28
	≈150.0	312	110.3	36.2	23.6	65.1	0.21	0.33

647

648

649 Table 4

650 Fatty acid composition of cellular lipids of *Rhodosporidium torolooides* DSM 4444 growing in shake flasks in previously  
 651 pasteurized media presenting increasing initial glucose (Glc<sub>i</sub>) concentration and constant NaCl concentration (A) and  
 652 fatty composition of lipids of the same microorganism growing in batch-bioreactor experiments in media presenting  
 653 increasing initial glucose (Glc<sub>i</sub>) concentration and constant NaCl concentration (B). In the flask experiments, for  
 654 Glc<sub>i</sub>≈50 g/L, very early (VE) growth phase is that in which the incubation time is between 10-20 h and late growth  
 655 phase is that in which incubation time is c. 150 h. For Glc<sub>i</sub>≈100 g/L, very early (VE) growth phase is that in which the  
 656 incubation time is between 20-30 h and late growth phase is that in which incubation time is c. 300 h. In the bioreactor  
 657 experiments, for Glc<sub>i</sub>≈50 g/L, very early (VE) growth phase is that in which the incubation time is between 10-20 h,  
 658 early (E) growth phase is that in which the incubation time is between 30-40 h and late growth phase is that in which  
 659 incubation time is c. 70 h. For Glc<sub>i</sub>≈100 g/L, very early (VE) growth phase is that in which the incubation time is  
 660 between 20-30 h, early (E) growth phase is that in which the incubation time is between 50-80 h and late growth phase  
 661 is that in which incubation time is c. 150 h.

662

663 A)

Glc <sub>i</sub> (g/L)	Growth phase	C14:0	C16:0	$\Delta^9$ C16:1	C18:0	$\Delta^9$ C18:1	$\Delta^{9,12}$ C18:2	$\Delta^{9,12,15}$ C18:3
≈50	VE	0.6	26.4	0.7	8.4	54.4	5.9	2.3
	L	1.1	23.0	0.5	7.9	57.4	6.9	2.6
≈100	VE	1.2	24.8	-	9.6	52.9	6.5	2.7
	L	1.6	22.0	1.5	8.2	55.0	6.8	2.2

664 B)

Glc <sub>i</sub> (g/L)	Growth phase	C14:0	C16:0	$\Delta^9$ C16:1	C18:0	$\Delta^9$ C18:1	$\Delta^{9,12}$ C18:2	$\Delta^{9,12,15}$ C18:3
≈50	VE	1.5	25.2	-	12.0	50.2	8.4	2.0
	E	1.0	23.8	1.1	10.1	54.7	9.3	-
	L	1.2	22.1	1.5	10.8	55.7	6.5	1.4
≈100	VE	-	23.8	-	12.2	53.7	6.7	1.5
	E	-	22.1	2.5	11.9	55.4	5.8	2.0
	L	-	20.9	3.0	9.1	54.9	8.9	2.5

665

666

667 Table 5  
668 Experimental results of *Rhodosporidium toruloides* strains producing microbial lipid during growth on several carbon  
669 sources under various fermentation configurations and their comparisons with the present study.  
670

Strain	Substrate	Culture mode	X (g/L)	$Y_{L/X}$ (% w/w)	Reference
<i>R. toruloides</i> *	Pure stearic acid	Shake flasks	11.7	35.0	Gierhart [8]
<i>R. toruloides</i> AS2.1389	Glycerol	Shake flasks	19.2	47.7	Xu et al. [9]
<i>R. toruloides</i> AS2.1389	Glycerol	Batch bioreactor	26.7	69.5	"
<i>R. toruloides</i> AS2.1389	Glucose	Shake flasks	18.3	76.0	Li et al. [14]
<i>R. toruloides</i> CCT 0783	Sweet sorghum extract	Shake flasks	41.7	33.1	Matsakas et al. [15]
<i>R. toruloides</i> DSM 444	Glycerol/sunflower meal hydrolysate blend	Shake flasks	27.9	29.0	Leiva-Candia et al. [18]
<i>R. toruloides</i> DSM 444	Glycerol/sunflower meal hydrolysate blend	Fed-batch bioreactor	37.4	51.3	"
<i>R. toruloides</i> NRRL Y-27012	Biodiesel-derived glycerol	Shake flasks	30.1	40.0	Tchakouteu et al. [19]
<i>R. toruloides</i> Y4	Glucose	Fed-batch bioreactor	106.5	67.5	Li et al. [20]
<i>R. toruloides</i> Y4	Glucose (phosphate-limited trial)	Shake flasks	20.6	51.4	Wu et al. [27]
<i>R. toruloides</i> Y4	Glucose (phosphate-limited trial)	Shake flasks	19.4	62.4	"
<i>R. toruloides</i> Y4	Glucose (sulphate-limited trial)	Shake flasks	23.0	20.8	Wu et al. [28]
<i>R. toruloides</i> Y4	Glucose (sulfate-limited trial)	Shake flasks	14.2	55.6	"
<i>R. toruloides</i> Y4	Glucose (sulfate-limited trial)	Shake flasks	17.8	55.7	"
<i>R. toruloides</i> *	Glucose/pure stearic acid	Shake flasks	9.8	46.1	Gierhart [29]
<i>R. toruloides</i> CBS14	Glucose	Shake flasks	12.3	30.8	Moreton [30]
<i>R. toruloides</i> CBS14	Glucose	Shake flasks	8.0	42.5	Moreton [66]
<i>R. toruloides</i> CBS14	Fructose	Shake flasks	7.9	25.3	"
<i>R. toruloides</i> CBS14	Glycerol	Shake flasks	5.8	34.6	"
<i>R. toruloides</i> CBS14	Glucose	Batch bioreactor	12.5	42.9	"
<i>R. toruloides</i> CBS14	Fructose	Batch bioreactor	8.7	39.8	"
<i>R. toruloides</i> CBS14	Xylose	Batch bioreactor	8.3	42.2	"
<i>R. toruloides</i> Y4	Jerusalem artichoke extracts	Shake flasks	25.5	40.0	Zhao et al. [67]
<i>R. toruloides</i> Y4	Jerusalem artichoke hydrolysates	Fed-batch bioreactor	70.1	56.5	"

<i>R. toruloides</i> CBS14	Glucose	Fed-batch bioreactor	35.0	71.4	Wiebe et al. [68]
<i>R. toruloides</i> CBS14	Glucose/xylose/arabinose blend	Fed-batch bioreactor	27.0	55.5	”
<i>R. toruloides</i> Y4	Biodiesel-derived glycerol	Batch bioreactor	35.3	46.0	Uçkun Kiran et al. [69]
<i>R. toruloides</i> 2F5	Inulin	Shake-flasks	15.8	62.1	Wang et al. [70]
<i>R. toruloides</i> 2F5	Inulin	Batch bioreactor	15.6	70.4	”
<i>R. toruloides</i> CCT 0783	Glucose/xylose blend	Batch bioreactor	13.3	42.0	Bonturi et al. [71]
<i>R. toruloides</i> DSM 444	Glucose	Batch bioreactor	~22	~40	Bommareddy et al. [72]
<i>R. toruloides</i> DSM 444	Glycerol	Batch bioreactor	~15	~57	”
<i>R. toruloides</i> AS2.1389	Glucose	Single-stage continuous bioreactor	8.7	61.8	Shen et al. [73]
<i>R. toruloides</i> AS2.1389	Glucose	Single-stage continuous bioreactor	5.7	53.1	”
<i>Rhodosporidium toruloides</i> Y4	Biodiesel-derived glycerol	Shake flasks	24.9	48.9	Yang et al. [74]
<i>R. toruloides</i> DSM 444	Glucose (NaCl-enriched culture)	Fed-batch bioreactor	37.2	64.5	Present study

671     \*: No indication of the strain

672

673

674 Fig. 1

675 Kinetics of lipid production (L; g/L) and lipid in dry cell weight ( $Y_{L/X}$ ; % w/w) during growth of *Rhodosporidium*  
676 *toruloides* DSM 4444 on glucose-based media (100 g/L) supplemented with 4% (w/v) NaCl. Culture conditions as in  
677 Table 1. Two lots of independent cultures were conducted by using different inocula. In all of the determinations,  
678 standard error calculated was less than 10%.

679

680 Fig. 2

681 Kinetics of residual glucose (Glc; g/L), biomass production (X; g/L) and lipid accumulated (L; g/L), during  
682 *Rhodosporidium toruloides* DSM 4444 growth in previously sterilized (filled symbols) and pasteurized (open symbols)  
683 media with 100 g/L of glucose and 4% (w/v) NaCl. Culture conditions as in Table 2. Two lots of independent cultures  
684 were conducted by using different inocula. In all of the determinations, standard error calculated was less than 10%.

685

686 Fig. 3

687 Kinetics of residual glucose (Glc; g/L) and lipid accumulated (L; g/L) during batch-bioreactor cultures of  
688 *Rhodosporidium toruloides* DSM 4444 in media containing 50 g/L (open symbols), 100 g/L (grey symbols) and 150 g/L  
689 (filled symbols) of glucose, supplemented with 4.0% (w/v) of NaCl. Culture conditions as in Table 4B. Two lots of  
690 independent cultures were conducted by using different inocula. In all of the determinations, standard error calculated  
691 was less than 10%.

692

693 Fig. 4

694 Kinetics of residual glucose (Glc; g/L), biomass production (X; g/L) and lipid accumulation (L; g/L) (A) and  
695 representation of biomass production (X; g/L) and lipid production (L; g/L) per substrate consumed (B) during fed-  
696 batch bioreactor cultures of *Rhodosporidium toruloides* DSM 4444, in media supplemented with 4.0% (w/v) NaCl.  
697 Culture conditions as in Table 4B. Two lots of independent cultures were conducted by using different inocula. In all of  
698 the determinations, standard error calculated was less than 10%.

699

700




Fig. 1




Fig. 2




Fig. 3




Fig. 4A




Fig. 4B