

Re-examining the U.K.'s greatest tornado outbreak: forecasting the limited extent of tornadoes along a cold front

Article

Accepted Version

Apsley, M. L., Mulder, K. J. and Schultz, D. M. (2016) Re-examining the U.K.'s greatest tornado outbreak: forecasting the limited extent of tornadoes along a cold front. *Weather and Forecasting*, 31 (3). pp. 853-875. ISSN 0882-8156 doi: 10.1175/WAF-D-15-0131.1 Available at <https://centaur.reading.ac.uk/61279/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/WAF-D-15-0131.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



AMERICAN METEOROLOGICAL SOCIETY

Weather and Forecasting

EARLY ONLINE RELEASE

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. Since it is being posted so soon after acceptance, it has not yet been copyedited, formatted, or processed by AMS Publications. This preliminary version of the manuscript may be downloaded, distributed, and cited, but please be aware that there will be visual differences and possibly some content differences between this version and the final published version.

The DOI for this manuscript is doi: 10.1175/WAF-D-15-0131.1

The final published version of this manuscript will replace the preliminary version at the above DOI once it is available.

If you would like to cite this EOR in a separate work, please use the following full citation:

Apsley, M., K. Mulder, and D. Schultz, 2016: Re-examining the U.K.'s Greatest Tornado Outbreak: Forecasting the Limited Extent of Tornadoes along a Cold Front. *Wea. Forecasting*. doi:10.1175/WAF-D-15-0131.1, in press.



**Re-examining the U.K.'s Greatest Tornado Outbreak:
Forecasting the Limited Extent of Tornadoes
Along a Cold Front**

MIRIAM L. APSLEY*, KELSEY J. MULDER, AND DAVID M. SCHULTZ

*Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Sciences, University of
Manchester, United Kingdom*

** Additional affiliation: Christ's College, University of Cambridge, United Kingdom*

Submitted as an Article to *Weather and Forecasting* 5 October 2015

Revised 12 February 2016 and 28 March 2016

Corresponding author address: Prof. David M. Schultz; Centre for Atmospheric Science; School of
Earth, Atmospheric and Environmental Sciences; University of Manchester; Simon Building, Oxford
Road; Manchester M13 9PL, United Kingdom.
E-mail: David.Schultz@manchester.ac.uk

ABSTRACT

On 23 November 1981, a strong cold front swept across the U.K., producing tornadoes from the west to the east coasts. An extensive campaign to collect tornado reports by the Tornado and Storm Research Organisation (TORRO) resulted in 104 reports, the largest U.K. outbreak. The front was simulated with a convection-permitting numerical model down to 200-m horizontal grid spacing to better understand its evolution and meteorological environment. The event was typical of tornadoes in the U.K., with convective available potential energy (CAPE) less than 150 J kg^{-1} , 0–1-km wind shear of $10\text{--}20 \text{ m s}^{-1}$, and a narrow cold-frontal rainband forming precipitation cores and gaps. A line of cyclonic absolute vorticity existed along the front, with maxima as large as 0.04 s^{-1} . Some hook-shaped misovortices bore kinematic similarity to supercells. The narrow swath along which the line was tornadic was bounded on the equatorward side by weak vorticity along the line and on the poleward side by zero CAPE, enclosing a region where the environment was otherwise favorable for tornadogenesis. To determine if the 104 tornado reports were plausible, first possible duplicate reports were eliminated, resulting in as few as 58 tornadoes to as many as 90. Second, the number of possible parent misovortices that may have spawned tornadoes is estimated from model output. The number of plausible tornado reports in the 200-m grid-spacing domain was 22 and as many as 44, whereas the model simulation was used to estimate 30 possible parent misovortices within this domain. These results suggest that 90 reports was plausible.

1. Introduction

On 23 November 1981, a strong cold front swept across the U.K., producing an unprecedented 104 reports of tornadoes (Fig. 1). Called “Britain’s greatest tornado outbreak” (Rowe and Meaden 1985), this event had many more tornadoes than the next largest event just a month earlier on 20 October 1981, which spawned only 29 tornadoes (Turner et al. 1986; Rowe 2016). For comparison, even modest outbreaks by United States standards are relatively uncommon in the U.K. Specifically, over 90% of U.K. tornado days between 1980 and 2012 had fewer than 8 tornadoes (Fig. 14 in Mulder and Schultz 2015). The November 1981 outbreak is so exceptional that it distorts the historical record and climatologies of tornadoes in the U.K. and Europe. For example, in the U.K. tornado climatology by Mulder and Schultz (2015), several figures had to be plotted with the outbreak excluded (e.g., their Figs. 5–7, 9, 12, 13), and, in their review of tornadoes across Europe, Antonescu et al. (2016) found that the large number of reports produced a bias in their synthesized results and capped the total number of reports from this outbreak at 58. Given the large number of reports distorting the climatologies and that a scientific study of this event has not been performed in nearly 35 years, we believe that the time is right to re-examine this event.

The locations of these 104 reports (Fig. 1) come from the Tornado and Storm Research Organisation (TORRO). TORRO is a U.K. not-for-profit organization responsible for collecting tornado reports from the media, from over 350 observers in the U.K., and from the public through TORRO’s website <http://www.torro.org.uk> (e.g., Elsom et al. 2001; Doe 2016). Of the 104 reports on 23 November 1981, 35 came from media reports, 30 came from the public after a call for reports on Anglia Television, and 39 were the result of TORRO’s appeal in local newspapers (Rowe 1985). The challenges of severe-weather event verification can be immense,

even when events are well observed by expert meteorologists (Speheger et al. 2002; Trapp et al. 2006). The challenges are compounded when reports are collected well after the event from primarily nonmeteorologists, as was the case in this event.

The first tornadoes of the day occurred in Anglesey, on the west coast of Wales, where there were five reports around 1000 UTC; 20 houses were damaged there, and a summer house (comparable to a small temporary building or mobile home) was turned upside down (Kemp and Morris 1982). The next seven tornado reports occurred near Aughton (near Liverpool) at 1100 UTC, and there were four more in Greater Manchester at 1200 UTC. A further 11 reports in the early afternoon came from Birmingham, Nottinghamshire, and the East Midlands (east of Birmingham), where 20 large caravans (camper vans) were blown over (Rowe and Meaden 1985). Seven tornado reports clustered near Hull in the early afternoon. Many of the reports (43), however, came from East Anglia (northeast of Cambridge) between 1300 and 1600 UTC. The last tornadoes occurred in southeast Essex (east-northeast of London) just before 1600 UTC.

Despite the large number of reports and despite occurring in conjunction with an extensive southwest–northeast-oriented cold front advancing southeastward across nearly the entirety of the U.K., these tornado reports only occurred along a narrow swath 200–250 km wide and 400 km long (Fig. 1). Thus, *the first goal of this article is to determine why the tornadoes occurred in a narrow swath along an otherwise extensive cold front.*

There are few exact details given with most of the tornado reports, so the duration of each tornado is unknown; some eyewitnesses, however, estimated lifetimes of 20–30 seconds or less (Rowe 1985). Eight tornado reports included damage track lengths between 0.3 and 4 km long. Where estimated, damage tracks are believed to have had widths of 10–20 m. The direction of travel was reported in 16 cases, with 14 coming from between west and north-northwest,

consistent with the movement of the front; the other 2 came from the south and southwest. Ninety-nine of the 104 reports were assigned values on the International Tornado Intensity Scale or T scale (Meaden 1983; Meaden et al. 2007). Compared to the Fujita (F) scale, the T scale has twice as many classifications. Conversion between the F and T scales can be performed using the equation $F \approx 0.5T$ and rounding down to the nearest integer (Brooks and Doswell 2001; Meaden et al. 2007). Figure 2 shows that most (96, or 97%) of the tornadoes were T0–T3 (or F0–F1); three tornadoes, however, reached T4 (or F2). This distribution is similar to the national distribution from the Kirk (2014) and Mulder and Schultz (2015) climatologies in which 94–95% were between T0–T3 (or F0–F1).

Because tornado reporting in this event relied on responses from a media campaign, the figure of 104 tornado reports has been controversial. Extrapolating based on the density of reports and the sparsely populated areas over which most of the cold front traveled, Rowe and Meaden (1985) suggested that the number of tornadoes may possibly have been as high as 400–500. On the other hand, only 58 of these reports (56% of the 104 reports) were later verified by TORRO experts according to the TORRO database and classified as definite; the other 46 reports were deemed to show reasonable evidence of a tornado having occurred, but not enough to be certain—these were classified as probable. Thus, the lack of confirmation of nearly half of the reports and the extreme magnitude of the outbreak in the historical context suggest that the 104 reports might be an overestimate.

In this article, we use two different approaches to investigate the tornado reports. First, we undertake a re-examination of the individual tornado reports for the possible occurrence of multiple reports of the same tornado. Second, we use a cloud-resolving model to simulate the cold front and possible parent circulations to the tornadoes. The number of parent circulations

might give us some insight into the number of tornadoes. Thus, *the second goal of this article is to re-examine the tornado reports and the meteorological conditions on that day to see if we can constrain the minimum and maximum number of tornadoes that likely occurred.*

2. Background on tornadoes along cold fronts

Tornadoes forming along cold fronts are a challenging forecasting problem. Such tornadoes are often associated with a class of convective storms occurring along cold fronts called narrow cold-frontal rainbands. Narrow cold-frontal rainbands have been described by Browning and Harrold (1970), Browning and Pardoe (1973), Carbone (1982), Hobbs and Persson (1982), Browning and Reynolds (1994), Browning and Roberts (1996), Browning et al. (1997), Jorgensen et al. (2003), and Viale et al. (2013), among many others. Narrow cold-frontal rainbands have been synthesized by conceptual models in Browning (1990) and Houze (2014, section 11.4.4). In the United States, narrow cold-frontal rainbands are a subset of what have been termed *quasi-linear convective storms* (QLCSs; Trapp et al. 2005). Trapp et al. (2005) applied this term for their investigation of tornadoes that form along such line convection (i.e., distinct from supercell convective storms).

Tornadoes along linear convective systems are challenging because they tend to have shorter lead times than tornadoes associated with supercells (Trapp et al. 2005). Even if the specific location and timing of the tornadoes cannot be predicted well in advance, predicting the general location along the line where tornadoes form would be an operationally useful tool. Indeed, Atkins et al. (2004) showed that tornadoes were more likely to form from parent mesovortices along the convective line that had greater rotation rates, implying that the strongest vortices may favor tornadogenesis.

Before discussing how tornadoes form along linear convective storms, we need to distinguish between the parent circulations that precede the tornadoes and the tornadoes themselves. One of the characteristics often observed in narrow cold-frontal rainbands is the presence of precipitation cores and gaps, aligned anticyclonically relative to the front. These core-and-gap regions have been reported for cold fronts over the eastern North Pacific Ocean (e.g., Hobbs and Biswas 1979; Hobbs and Persson 1982; Jorgensen et al. 2003), near the Alps (Hagan 1992), over eastern North America and the Atlantic Ocean (Locatelli et al. 1995; Wakimoto and Bosart 2000), and over the U.K. (e.g., James and Browning 1979; Browning and Roberts 1996). Specifically, other tornadic cold fronts in the U.K. also possessed this core-and-gap structure (e.g., Smart and Browning 2009; Clark and Parker 2014; Mulder 2015), as well as in Japan (e.g., Kobayashi et al. 2007; Sugawara and Kobayashi 2009).

The cores are often associated with heavier precipitation and relative maxima in vorticity (hereafter misovortices), whereas the gaps are associated with weaker precipitation or the absence of precipitation and relative minima in vorticity. Misovortices have diameters of 1–4 km (Fujita 1981) and have been suggested to be the parent circulation from which the tornadoes form. Different explanations have been offered to explain misovortex formation, including the release of horizontal shearing instability (e.g., Carbone 1982; Hobbs and Persson 1982; Lee and Wilhelmson 1997b; Jorgensen et al. 2003; Wheatley and Trapp 2008; Kawashima 2011), advection of hydrometeors (Locatelli et al. 1995), trapped gravity waves (Brown et al. 1999), tilting of vorticity along the cold front (Carbone 1983), or combinations of the above.

How tornadoes form along linear convective storms is less well known compared to supercellular tornadoes, primarily because detailed field observations of tornadoes forming along linear convective storms have not been collected and because of the large computational expense

of producing a tornado within a numerical model. Because of the shorter lead time and the different parent-storm morphology to supercells, Trapp et al. (1999) suggested that a different tornadogenesis mode may be responsible for tornadoes from linear convective systems than tornadoes from supercells. Carbone (1983) found that the downdraft was coincident with the tornado, suggesting the importance of tilting and a similarity with tornadogenesis in supercells. In contrast, Lee and Wilhelmson (2000) found the importance of stretching of strong initial vorticity in their simulations of nonsupercell tornadogenesis. Nevertheless, the data and simulations in this article will be insufficient to address the issue of tornadogenesis in this case. Thus, we focus on the misocyclones, the locations of the tornadoes, and an approach to forecast the occurrence of tornadoes along lines, as demonstrated for the case of Britain's greatest tornado outbreak on 23 November 1981.

3. Observations: Synoptic and mesoscale overview

At 12 UTC 22 November 1981, archived Met Office charts identified a broad region of low pressure with two centers of 994 hPa and 996 hPa centered southeast of Iceland and north of the U.K. (Fig. 3a). Twelve hours later, the cyclone consolidated with a central pressure of 986 hPa (not shown). By 12 UTC 23 November, the low had rapidly deepened another 18 hPa to 968 hPa and was moving toward Norway (Fig. 3b), making landfall by 0600 UTC 24 November with a central pressure of 959 hPa (not shown). The cyclone was associated with a sharp trough in 500-hPa geopotential height and strong geostrophic cold advection in the lower troposphere, as indicated by the 1000–500-hPa thickness contours (Fig. 4).

Associated with this cold advection was a strong cold front at the surface. Archived hourly Met Office surface maps show the front extending to the south of the cyclone across the U.K.

and its southeastward progression (Fig. 5). As the cold front crossed England and Wales, temperatures fell by 6–7°C in the first hour, and the pressure rose by as much as 4–5 hPa in the first hour after frontal passage and 3 hPa hr⁻¹ thereafter (Rowe and Meaden 1985; Fig. 5). The wind direction veered suddenly from 190°–230° before the front, a direction roughly parallel to the front, to 320°–340° after the front, a postfrontal direction nearly perpendicular to the orientation of the front (e.g., Fig. 5c). By 1800 UTC, the cold front had cleared England and moved over the North Sea (Rowe 1985; Rowe and Meaden 1985). Moderate rain preceded and was associated with the front in northwestern England at the hour ending 1200 UTC (as much as 10 mm per hour; Fig. 6). The infrared satellite image at 1325 UTC (Fig. 7) showed the low center to the north of the U.K. and the broad band of clouds associated with the cold front and a prefrontal band. As the front moved southeastward into central England, the precipitation weakened dramatically to less than 2 mm per hour during the hour ending at 1400 UTC (Fig. 6).

Unfortunately, none of the operational soundings that day were ideal for sampling the prefrontal air. The nearest proximity sounding occurred at Aughton near Liverpool, about 11 h before frontal passage at 0000 UTC (Fig. 8). This sounding exhibited only 13 J kg⁻¹ convective available potential energy (CAPE), a steep lapse rate between 850 and 700 hPa, and a strong 60 kt (31 m s⁻¹) westerly wind at 850 hPa (Fig. 8). A sounding from the NCEP–NCAR Reanalysis (Kalnay et al. 1996) for a location in western England (52.5°N, 2.5°W) at 1200 UTC 23 November had a surface-based CAPE of 147 J kg⁻¹, which is only slightly higher than the 50–100 J kg⁻¹ of CAPE from the model simulation initialized from the European Centre for Medium-Range Weather Forecasting (ECMWF) reanalyses (jump ahead to Fig. 13).

These conditions—strong cold front, small CAPE, prefrontal winds nearly parallel to the front, and postfrontal winds nearly perpendicular to the front—are consistent with weather

conditions associated with other tornado outbreaks in the U.K. (e.g., Bolton et al. 2003; Holden and Wright 2004; Clark 2009, 2013; Clark and Parker 2014; Mulder 2015). Given the synoptic situation, the morphology of the convective storm (also called its convective mode) is likely consistent with previous tornadic convective storms over the U.K., which tend to occur along cold fronts in linear convective storms. Linear convective storms account for 42% of the tornadoes and 51% of the tornado outbreaks in the U.K. (Mulder and Schultz 2015), unlike in the United States where linear storms account for only 18–25% of the tornadoes (Trapp et al. 2005; Smith et al. 2012). [In comparison, supercells produce 79% of U.S. tornadoes (Trapp et al. 2005).] Clark (2013) examined 103 convective lines in the UK and found that 27% were associated with at least one tornado, further evidence for the importance of these lines in producing tornadoes in the U.K.

Because radar data for this event (Doppler winds or even reflectivity) are unavailable, the precipitation structure of the cold front on that day is unknown. Therefore, we investigate this event further with a model simulation.

4. Model simulation: Set-up

As has been demonstrated for other cases, model simulations can be an effective tool for understanding tornadic fronts in the U.K. (e.g., Smart and Browning 2009; Groenemeijer et al. 2011; Mulder 2015). Therefore, we performed a convection-permitting simulation to construct a four-dimensionally consistent dataset to explore a likely meteorological evolution for this event. A successful simulation would be useful to interpret the conditions favorable for the tornadoes within the narrow swath and help interpret the 104 reports of tornadoes.

The simulation was performed using the Advanced Research Weather and Forecasting Model version 3.4.1 (WRF-ARW; Skamarock et al. 2008). The simulation was initialized at 0600 UTC 23 November 1981 from the ECMWF reanalysis at $0.25^\circ \times 0.25^\circ$ grid spacing interpolated onto a Lambert conformal grid. Lateral boundary conditions were provided by the ECMWF reanalyses every 6 h. Otherwise, the simulation was set up exactly the same as that in Mulder (2015) for the more modest U.K. tornado outbreak of 29 November 2011, which featured seven reported tornadoes across Wales and northern England. The simulation featured 90 vertical levels and four domains, ranging from the outermost domain with 25-km horizontal grid spacing, to three two-way nested domains of 5-km, 1-km, and 200-m horizontal grid spacing (the innermost two domains are shown in Fig. 9). Even at 200-m grid spacing, the model would have been inadequate to resolve any possible tornadoes. Instead, the innermost domain is analyzed for the existence of mesocyclones, small-scale circulations along linear convective systems that may precede tornadoes. Only output from the 1-km and 200-m domains is shown in the present article. Model output was saved for further diagnosis every 30 min for the 1-km domain and every 10 s for the 200-m domain.

The Kain–Fritsch convective parameterization (Kain and Fritsch 1990; Kain 2004) was employed on the outermost 25-km domain only. Other physical parameterizations included the five-layer thermal diffusion land-surface scheme (Skamarock et al. 2008, their section 8.4.1), Thompson et al. (2008) cloud microphysics, and Mellor–Yamada–Janjić boundary layer (Mellor and Yamada 1982; Janjić 1994, 2002). These parameterizations were chosen because Mulder (2015) found that they produced the most successful simulation of her case. Testing three different microphysical parameterizations (WRF single-moment six-class scheme; Morrison et

al. 2009; Thompson et al. 2008) did not produce different structures for the core-and-gap regions along the cold front in this case.

5. Model simulation: Mesoscale analysis

The meteorology on the 1-km domain is presented in this section. Subsequent analysis in this article occurs at 1000 UTC, around the time the first tornadoes were reported in Anglesey, and at 1400 UTC, just before the majority of tornadoes were reported in East Anglia. To illustrate the intensity of the front, surface temperature, wind, and sea-level pressure at 1000 UTC and 1400 UTC are presented in Fig. 10. The passage of the front was associated with a sharp pressure trough, temperature drop of 6°–8°C, and nearly a 90° veering of the wind (Fig. 10). The winds on either side of the front changed direction from 180°–230° on the warm side to 310°–330° on the cold side, although the wind speeds were roughly the same across the front at about 5–10 m s⁻¹. The simulation is consistent with the observations reported in section 3, except for the simulation being an hour behind the observations (cf. Figs. 5a and 10a; cf. Figs. 5c and 10b).

At 1000 UTC 23 November, simulated radar reflectivity factor reveals poorly organized precipitation along the pressure trough over most of the domain, with reflectivities of up to 45 dBZ, ahead of the wind shift along the cold front around the time the band first arrived in the U.K. (Fig. 11a). Along the cold front in the northwest part of the domain, a shorter, narrower, more organized, and more intense (45–50 dBZ) line of convection developed (Fig. 11a). As the rainband progressed across the U.K., the areal coverage of the precipitation decreased as the along-front extent of the rainband increased, consistent with the observations (cf. Figs. 6 and 11). In particular, as the line passed over the Pennine mountain range in the center of northern

England, much of the precipitation weakened and the band split into a higher reflectivity line positioned along the front, and a line of precipitation tens of km ahead of the front (Fig. 11b). The observed counterpart to the modeled prefrontal band, although present in the satellite imagery (Fig. 7 at 1325 UTC), did not appear to produce any measurable precipitation at the rain gauges (Fig. 6 during 1300–1400 UTC). Whether this is because the band was poorly forecast or the stations did not receive rain is unclear at this time. In any case, this prefrontal band is not the focus of the article as it is not associated with the formation of the tornadoes. At maturity of the convective line, cores of stronger precipitation became separated by gaps of about 10 km in length of lighter or no precipitation, similar to previously published work summarized in section 2.

The front was associated with a line of absolute vorticity maxima at 500 m above sea level, which was strongest to the north and weakest to the south (Fig. 12) because the zone of wind shift across the front broadened in association with a weaker pressure trough (Figs. 5b,c and 10b). This line of absolute vorticity maxima was between 0.005 and 0.01 s^{-1} and contained small maxima of 0.01 – 0.02 s^{-1} , as calculated on the 1-km grid. The line of vorticity moved across Britain with the cold front (Fig. 12); there were maxima in vorticity over Anglesey at 1100 UTC, near Liverpool about 1200 UTC, and in southeast England at 1500–1600 UTC, passing southeast of the U.K. by 1730 UTC. These times correspond within about an hour of reported tornado times (Rowe and Meaden 1985), which is all that can be expected given that the resolution of the tornado reports is only hourly, providing additional faith in the ability of the simulation to reproduce observed features of the front.

An examination of the three ingredients for deep moist convection (lift, moisture, and instability; e.g., Johns and Doswell 1992) shows that lift as much as several m s^{-1} was present

(not shown), associated with the strong convergence along the cold front inferred from the wind field (Fig. 11b). Moisture and instability can be diagnosed by CAPE (determined from the parcel with the maximum equivalent potential temperature in the column) (Fig. 13). At 1000 UTC, CAPE appeared as patchy areas east of the front, but generally less than 50 J kg^{-1} (Fig. 13a). By 1400 UTC, CAPE increased ahead of the front, with widespread areas over 25 J kg^{-1} and localized maxima approaching 125 J kg^{-1} , forming a slightly curved narrow (40–70 km wide) crescent of CAPE ahead of the front (Fig. 13b). (Interestingly, a second maximum of CAPE of $25\text{--}125 \text{ J kg}^{-1}$ was also present in a 20–50-km wide band about 150 km ahead of the front associated with the prefrontal rainband, although this maximum is not part of this story.) Therefore, the three ingredients for deep moist convection (i.e., instability, lift, moisture) were present along the front.

These large gradients in CAPE occurring over such short distances raise issues about the proximity soundings for U.K. tornadoes. Given the large gradients in CAPE occur over distances as small as tens of km, this raises questions about the choice of proximity sounding criteria used in Mulder and Schultz (2015) of 180 km and 3 h. Mulder and Schultz (2015) derived their criteria from previous proximity sounding studies in the United States, specifically Brooks (2009). Indeed, the prefrontal sounding for this outbreak in Fig. 8 does not meet these criteria. Other U.K. soundings on that day were even farther away from the tornadoes. Thus, the large variability in CAPE ahead of the front in this case is consistent with the recommendations for proximity sounding criteria for significant tornadoes in the United States of a range of 40–80 km and no more than 2 h (Potvin et al. 2010). Potentially noteworthy is the fact that detailed analysis of CAPE and convective inhibition near supercells in the central United States show variations of hundreds of J kg^{-1} over distances as small as a few km (e.g., Markowski et al.

2002). Therefore, perhaps our results of such strong gradients over tens of km should not be too surprising.

Given the reasonable timing and structure of the modeled front compared to the observations, we can interrogate the model output to determine the reasons that the tornado reports occurred within a relatively narrow swath along the front. Given the existence of organized deep moist convection, the potential for tornadogenesis can be explored with plots of lifting condensation level (LCL), 0–1-km wind shear, and 0–1-km storm-relative helicity. These are quantities known for their ability to discriminate tornadic from nontornadic storms in the United States (e.g., Rasmussen and Blanchard 1998; Thompson et al. 2003, 2012; Craven and Brooks 2004) and Europe (e.g., Púčik et al. 2015; Mulder and Schultz 2015).

At 1000 UTC, the lowest LCL along the front was between 600 and 1000 m (Fig. 14a). By 1400 UTC, the LCL had dropped along a similar crescent-shaped spatial distribution of low LCL (200–600 m) in the south with patches less than 200 m, and significantly higher LCL (greater than 2200 m) behind the front and to the north along the front (Fig. 14b). These results are consistent with conditions for tornadoes in the U.K. Specifically, Mulder and Schultz (2015) found that low LCL height was a statistically significant factor in predicting tornado formation in the U.K., with outbreaks having a mean LCL of about 700 m, as opposed to a null set of convective storms with lightning or hail which had an LCL of 900 m. Therefore, we would expect tornadic storms to be found along the line toward the south where the LCL is lower and the CAPE is higher.

The vertical shear of the horizontal wind over the surface to 1 km layer (i.e., 0–1-km wind shear) displayed a sharp change in magnitude across the front (Fig. 15). Just ahead of the front in the swath where the tornadoes formed, the shear was 10–20 m s^{-1} , with values over 30 m s^{-1} in

the prefrontal rainband (Fig. 15b). Behind the front, the shear was only around 5–10 m s⁻¹. Storm-relative helicity over 0–1 km also showed rather large values ahead of the front (Fig. 16). In the immediate vicinity of the front in the prefrontal environment, 0–1-km storm-relative helicity ranged from zero to several hundred m² s⁻² (Fig. 16).

Thus, despite the cold front extending across nearly the entirety of the U.K. (Figs. 5b,c), the narrow swath of tornado reports occurred in what was apparently a sweet spot for the conditions favoring deep moist convection and tornadogenesis along squall lines. Specifically, the swath of tornado reports in this case was limited on the poleward side by the rapidly increasing LCL heights and decreasing CAPE and limited on the equatorward side by the rapidly decreasing absolute vorticity along the cold front, in a prefrontal environment with adequate low-level wind shear and storm-relative helicity all along the front. Although forecasting tornadoes along linear convective systems remains a challenging forecast problem, this sweet spot may provide insight into providing more specificity for nowcasting tornado development along future linear convective systems in the U.K. or elsewhere.

6. Model simulation: Misovortex structure and evolution

The majority of tornado reports occurred within the model domain with 200-m horizontal grid spacing as the modeled front passed through this domain between 1300 UTC and 1640 UTC (Fig. 12). Analysis of vorticity, reflectivity, and surface winds from this domain exhibits more detail along the front where the majority of tornado reports occurred. This region is also where this apparent sweet spot favorable for tornadogenesis occurred.

At this higher resolution, more detail in the structure and evolution of the misovortices is apparent. Specifically, regions of larger 500-m absolute vorticity ($0.02\text{--}0.03\text{ s}^{-1}$) developed into maxima of $0.035\text{--}0.04\text{ s}^{-1}$ within the line, with 500-m updrafts of $5\text{--}10\text{ m s}^{-1}$ (e.g., Fig. 17). Pairing of absolute vorticity maxima and minima was common both within the line and in a few patches a little ahead of the line, where there was some higher reflectivity as well. Some merging and splitting of downdrafts and maxima, which has been shown to increase vorticity (Lee and Wilhelmson 1997a), was observed, as well. Background reflectivity of 35–45 dBZ occurred within the rainband, with some patches of higher reflectivity of 50–55 dBZ. Similar to the core-and-gap structures observed by Mulder (2015), the shapes of the misovortices at their maximum intensity are quite similar to each other, specifically, an updraft (usually $5\text{--}10\text{ m s}^{-1}$) located poleward of the misovortex and a downdraft ($3\text{--}6\text{ m s}^{-1}$, although some downdrafts were as large as $6\text{--}9\text{ m s}^{-1}$) located equatorward of the misovortex.

Where the rainband looked like a hook or breaking wave at its edge, the misovortex was typically located at the rear edge of the rainband in the area of lower reflectivity (10–15 dBZ), and eventually developed a hook shape (Fig. 18). Many misovortices intensified at the center of the rainband and weakened as they moved backwards relative to the rainband, leaving the misovortices on the cold side. Some evolved from a line of vorticity that curled up and split into two hooks often described as a broken-S (McAvoy et al. 2000; Clark 2011), signatures similar to the line-echo wave pattern (Nolen 1959) and the frontal type of misovortices observed modeled in squall lines (Jewett and Wilhelmson 2006). The hook-shaped echo is likely a response to the circulation around the misovortex. The kinematics of misovortices appear similar to that of supercells and may suggest that tornadoes along lines may form similar to that inside a supercell,

as suggested by Weisman and Trapp (2003). Further investigation is required to confirm whether the dynamics are similar.

To estimate how important these hook-shaped cells were in the model, all the misovortices with absolute vorticity greater than 0.02 s^{-1} were plotted every minute between 1350 and 1450 over the 200-m domain when the front was in the plotted area of Fig. 12 ($110 \times 70 \text{ km}$). Previous simulations of vortices in different storm types have produced vortices about this magnitude. For supercells, Adlerman et al. (1999) found vorticity up to 0.054 s^{-1} . For bow echoes, vorticity magnitudes ranged from 0.009 to 0.02 s^{-1} (Weisman and Trapp 2003; Trapp and Weisman 2003; Wheatley and Trapp 2008; Atkins and St. Laurent 2009). For narrow cold-frontal rainbands, Smart and Browning (2009) found vorticity up to 0.04 s^{-1} . Although the modeled vorticity magnitudes depend on the case, they also depend on model grid spacing with higher-resolution models producing higher vorticity values. We determined that 0.02 s^{-1} was a good balance between choosing a smaller value with vorticity maxima everywhere and choosing a higher value with relatively few vorticity maxima. If the numbers of vortices and vortices with hooks are calculated every ten minutes during that 60-minute period (seven times), then an average of 39 (with a standard deviation of 3) misovortices existed, of which 20.4% (with a standard deviation of 2.6%) displayed hooks at any one time. Thus, this evolution is relatively common with the model simulation.

7. Reassessment of number of reports

We can use the simulation, in conjunction with a re-examination of the reports, to re-examine this event. First, according to the TORRO database, 58 of these reports (56% of the 104 reports) were later verified by TORRO experts, and classified as definite by them; the other 46

reports were deemed to show reasonable evidence of a tornado having occurred, but not enough to be certain—these were classified as probable. So, the minimum number of credible tornadoes was deemed to be 58.

Second, the remaining 46 probable tornado reports were examined for likely duplicate reports. The following approach was followed. Each of the 46 probable reports was checked to see if it might have duplicated another report. Duplicate reports were defined in this article as those reports occurring close in space and time, generally 5 km or closer and reported at the same time. Because the reports in the TORRO database are recorded by the hour, in practice this meant tornadoes reported during the same hour. If the duplicate probable report overlapped with a definite report, then the definite report was retained and the probable report was discarded. If a definite tornado report with unknown intensity was combined with a probable tornado report with known intensity, then the intensity was assigned to the single definite report. If the duplicate probable report overlapped with another probable report, then the more trustworthy probable report was retained and the other report was discarded. Those reports that had been checked by TORRO experts or were possessing tornado tracks, direction of travel or high T-scale value were deemed to be the most trustworthy and retained. This check reduced the number of probable reports by 14 to 32. These two checks reduced the number of tornadoes on 23 November 1981 to as few as 58 and as many as 90 tornadoes (Fig. 19).

Does the simulation provide support for this many tornadoes? The innermost model domain over a section of southeast England contained a subset of 52 of the 104 reports and included part of the area targeted by Anglia Television with their 30 reports, which is why this area had a relatively high percentage of probable reports (e.g., Fig. 1). The re-examination above reduced these 52 reports to 42 tornadoes (22 definite and 20 probable tornadoes).

To produce tracks of these misovortices that might be parent circulations for tornadoes, absolute vorticity greater than 0.02 s^{-1} was plotted every minute over the 200-m domain (Fig. 20a). Taking 30 min (± 2 min because the data interval is every minute) as an approximate minimum lifetime for a parent misovortex to produce a tornado (e.g., Wakimoto and Wilson 1989; Brady and Szoke 1989), the number of misovortices produced by the model was counted. This plot was repeated for absolute vorticity maxima greater than 0.025 s^{-1} , updrafts greater than 5, 6, and 7 m s^{-1} , and downdrafts greater than 2, 3, and 4 m s^{-1} (Fig. 20). The results of counting these tracks are summarized in Table 1, which include the average and median duration of tracks lasting 30 min or more (termed *long-lived*), and the longest duration and track lengths. These results show a substantial number of long-lived tracks of various intensities (e.g., 9 misovortices of 0.025 s^{-1} or more, 23 updrafts of 5 m s^{-1} or more, 10 downdrafts of 2 m s^{-1} or more). Most of the model tracks were from the northwest (note the line of constant longitude in the panels in Fig. 20), consistent with the TORRO reports of tracks being mostly from the northwest. A few tracks from the west or southwest, however, were also present (e.g., Figs. 20a,c), which was also consistent with a few tornado reports.

Although some of the tracks of misovortices with vorticity greater than or equal to 0.02 s^{-1} on the 200-m grid are within 5 km of each other, tornado reports less than 5 km apart are more likely to represent the same tornado than ones say 20 km apart. We can never claim that our approach is perfect, but merely suggests a plausible way to filter possibly duplicate reports. Also, there was some ambiguity in how the locations of the reports were recorded (which may have been as specific as the name of a town, rather than a quantitative latitude–longitude coordinate). Such ambiguities would complicate the assessment of the duplicate reports. Finally, the tornado reports that were discounted were listed as only probable by TORRO, so

there is no risk of eliminating definite tornadoes. Thus, we are confident in the model's ability to produce a large number of misovortices that are consistent with the large number of tornado reports widespread over a large region of England and Wales.

If the tornadoes on this day developed from parent misovortices that were formed by the tilting-shear mechanism (e.g., Trapp and Weismann, 2003), we would expect horizontal vorticity to develop first, increase, be tilted vertically by an updraft–downdraft dipole, and then weaken. Thus, we would expect the parent misocyclone to have a shorter lifetime than the updraft. From Fig. 20 and Table 1, examples of tracks of updrafts ($5\text{--}10\text{ m s}^{-1}$) and tracks of vorticity greater than 0.02 s^{-1} had similar lengths. The vorticity increased to above 0.025 s^{-1} along the tracks and then toward the end of the tracks. Downdrafts of $3\text{--}6\text{ m s}^{-1}$ also appeared alongside these tracks for shorter lengths than the updrafts and of only slightly shorter lengths than the higher vorticity tracks. Counting the number of absolute vorticity maxima of 0.02 s^{-1} or more that last for 30 min or longer yields 41 misovortices, with some of the longer-lasting updrafts forming multiple misovortices. Of these 41 misovortices, 30 have updrafts of 5 m s^{-1} or more and downdrafts of 3 m s^{-1} or more each lasting longer than 4 min, meaning that there are roughly 30 possible parent circulations in the 200-m domain alone (Fig. 21). Of these 30 tracks, the average lifetime of the tracks was 47.6 min (median of 39 min), and the longest track was 175 km and lasted for 109 min. When linked with favorable environmental conditions for tornadogenesis in the model and the results of Atkins et al. (2004) who found that tornadoes were more likely to form from parent misovortices along the convective line that had greater rotation rates, the potential existed for the model misovortices to have been tornadic. Thus, these roughly 30 intense misovortices within the innermost domain are sufficient to explain the 22–44 tornado reports within this domain.

Figure 22 combines the half-hourly absolute vorticity isochrones with the regions with favorable CAPE and vorticity values, and the observed 90 tornado reports. The majority of the tornado reports (89 out of 90) were within the favorable locations (high vorticity along the cold front and nonzero CAPE). Also, there was agreement between the modeled misocyclone tracks and the locations of the tornado reports, providing additional veracity of the simulation. The possibility also existed that these misovortices could have produced multiple tornadoes each. Therefore, these statistics give an indication of the potential of high-resolution modeling to resolve features potentially responsible for the tornadoes, as convection-permitting simulations did 15 years ago for the 3 May 1999 Oklahoma–Kansas supercellular tornado outbreak (e.g., Roebber et al. 2002), and provides justification for a potentially large number of possible parent circulations for tornadogenesis in this event.

8. Conclusions

The U.K. tornado outbreak of 23 November 1981 is analyzed from a convection-permitting model simulation and a re-examination of the 104 tornado reports collected by TORRO. This case is called “Britain’s greatest tornado outbreak” (Rowe and Meaden 1985) because its 104 reports were so much greater than the next highest outbreak of 29. A synoptic situation with a strong cold front, weak CAPE (less than 125 J kg^{-1}), prefrontal winds nearly parallel to the front, and postfrontal winds nearly perpendicular to the front is consistent with weather conditions associated with other tornado outbreaks in the U.K. (Clark 2009; Clark and Parker 2014).

The model simulation produced a narrow cold-frontal rainband along a line of absolute vorticity exceeding 0.02 s^{-1} on the 200-m grid with embedded maxima of $0.035\text{--}0.04 \text{ s}^{-1}$, similar to those in previous simulations of misovortices along cold fronts in the U.K. (Smart and

Browning 2009). Misovortices along the front formed a variety of different structures and evolutions and may have been parent circulations for the tornadoes. A line of reflectivity along the cold front was characterized by precipitation cores and gaps. Updrafts of $5\text{--}10\text{ m s}^{-1}$ occurred poleward of these maxima of absolute vorticity, and weaker downdrafts of $3\text{--}6\text{ m s}^{-1}$ occurred equatorward, suggesting the potential for tilting to be involved in tornadogenesis.

The line of absolute vorticity weakened rapidly to the south in conjunction with a weakened pressure trough. Nearly all of tornadoes reported occurred within a sweet spot where the absolute vorticity was strong enough (more than 0.002 s^{-1} on the 1-km grid) and the CAPE was positive in an environment that was otherwise favorable for tornadoes (0–1-km storm-relative helicity and 0–1-km shear). This approach suggests a means by which regions favorable for tornadoes along squall lines could be forecast in the U.K. and elsewhere. The narrow (tens of km) region of positive CAPE in advance of the front also raises concerns about large distances used in determining proximity soundings in previous studies (100–200 km).

Within the model domain with 200-m horizontal grid spacing, 30 possible parent misovortices were present with the following characteristics: absolute vorticity greater than 0.02 s^{-1} , updrafts between $5\text{ and }10\text{ m s}^{-1}$ for longer than 30 min, and downdrafts between $3\text{ and }6\text{ m s}^{-1}$ were present for at least 4 min. This number of parent misovortices was comparable to the figure of 22–44 tornado reports in this area. We conclude that the number of reports in this area was potentially credible.

Reassessing the quality, timing and location of the reports allows us to place revised boundaries on the lower and upper limit of the number of tornadoes that day. A final figure was produced of 90 tornadoes: 58 definite and 32 probable, a slight reduction from the 104 total reports. This revision does not eliminate the problem of the event distorting the historical record

(Mulder and Schultz 2015; Antonescu et al., 2016). Even if the lower limit were closer to 58 reports, this event would still be the largest documented tornado outbreak in the U.K.

Acknowledgements: We thank TORRO, in particular Terence Meaden and Paul Brown, for providing the tornado data for 23 November 1981, and Jeff Trapp, David Smart, and an anonymous reviewer for their comments that have improved this article. We thank ARCHER, the U.K. National Supercomputing Service, for hosting the simulations. The historical weather maps were provided by Mark Beswick of the National Meteorological Archive and Duncan Ball of the Met Office Library, and we thank Stephen Burt for telling us about the existence of these map archives. The satellite imagery was provided by Andrew Brooks and Neil Lonie of the Dundee Satellite Receiving Station. Jonathan Fairman provided the CAPE value from the NCEP–NCAR Reanalysis. Funding for Apsley was provided by the U.K. Natural Environment Research Council through the Manchester–Liverpool Doctoral Training Programme Grant NE/L002469/1. Funding for Mulder was provided by a scholarship from the Faculty of Engineering and Physical Sciences, University of Manchester. Partial funding for Schultz was provided by the Natural Environment Research Council to the University of Manchester through Grants NE/H008225/1, NE/I005234/1, and NE/N003918/1 and by the Risk Prediction Initiative of the Bermuda Institute of Ocean Sciences through Grant RPI2.0-2016-SCHULTZ.

REFERENCES

Adlerman, E. J., K. K. Droegemeier, and R. Davies-Jones, 1999: A numerical simulation of cyclic mesocyclogenesis. *J. Atmos. Sci.*, **56**, 2045–2069.

563 Antonescu, B., D. M. Schultz, F. Lomas, and T. Kühne, 2016: Tornadoes in Europe: Synthesis of
 564 the observational datasets. *Mon. Wea. Rev.*, doi: 10.1175/MWR-D-15-0298.1.

565 Atkins, N. T., and M. St. Laurent, 2009: Bow echo mesovortices. Part I: Processes that influence
 566 their damaging potential. *Mon. Wea. Rev.*, **137**, 1497–1513.

567 Atkins, N. T., J. M. Arnott, R. W. Przybylinski, R. A. Wolf, and B. D. Ketcham, 2004: Vortex
 568 structure and evolution within bow echoes. Part I: Single-Doppler and damage analysis of the
 569 29 June 1998 derecho. *Mon. Wea. Rev.*, **132**, 2224–2242, doi: [10.1175/1520-
 570 0493\(2004\)132<2224:VSAEWB>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<2224:VSAEWB>2.0.CO;2)

571 Bolton, N., D. M. Elsom, and G. T. Meaden, 2003: Forecasting tornadoes in the United
 572 Kingdom. *Atmos. Res.*, **67–68**, 53–72, doi:10.1016/S0169-8095(03)00083-8.

573 Brady, R. H., and E. J. Szoke, 1989: A case study of non-mesocyclone tornado development in
 574 northeast Colorado: Similarities to waterspout formation. *Mon. Wea. Rev.*, **117**, 843–856,
 575 doi: 10.1175/1520-0493(1989)117<0843:ACSONT>2.0.CO;2.

576 Brooks, H. E., 2009: Proximity soundings for severe convection for Europe and the United States
 577 from reanalysis data. *Atmos. Res.*, **93**, 546–553, doi:10.1016/j.atmosres.2008.10.005.

578 Brooks, H. E., and C. A. Doswell, 2001: Some aspects of the international climatology of
 579 tornadoes by damage classification. *Atmos. Res.*, **56**, 191–201, doi: 10.1016/S0169-
 580 8095(00)00098-3.

581 Brown, M. J., J. D. Locatelli, M. T. Stoelinga, and P. V. Hobbs, 1999: Numerical modeling of
 582 precipitation cores on cold fronts. *J. Atmos. Sci.*, **56**, 1175–1196.

583 Browning, K. A., 1990: Organization of clouds and precipitation in extratropical cyclones.
 584 *Extratropical Cyclones, The Erik Palmén Memorial Volume*, C. W. Newton and E. O.
 585 Holopainen, Eds., Amer. Meteor. Soc., 129–153.

586 Browning, K. A., and T. W. Harrold, 1970: Air motion and precipitation growth at a cold front.
 587 *Quart. J. Roy. Meteor. Soc.*, **96**, 369–389.

588 Browning, K. A., and C. W. Pardoe, 1973: Structure of low-level jet streams ahead of
 589 mid-latitude cold fronts. *Quart. J. Roy. Meteor. Soc.*, **99**, 619–638.

590 Browning, K. A., and R. Reynolds, 1994: Diagnostic study of a narrow cold frontal rainband
 591 and severe winds associated with a stratospheric intrusion. *Quart. J. Roy. Meteor. Soc.*, **120**,
 592 235–257.

593 Browning, K. A., and N. M. Roberts, 1996: Variation of frontal and precipitation structure along
 594 a cold front. *Quart. J. Roy. Meteor. Soc.*, **122**, 1845–1872.

595 Browning, K. A., N. M. Roberts, and A. J. Illingworth, 1997: Mesoscale analysis of the
 596 activation of a cold front during cyclogenesis. *Quart. J. Roy. Meteor. Soc.*, **123**, 2349–2375.

597 Carbone, R. E., 1982: A severe frontal rainband. Part I. Stormwide hydrodynamic structure. *J.*
 598 *Atmos. Sci.*, **39**, 258–279.

599 Carbone, R. E., 1983: A severe frontal rainband. Part II. Tornado parent vortex circulation. *J.*
 600 *Atmos. Sci.*, **40**, 2639–2654.

601 Clark, M. R., 2009: The southern England tornadoes of 30 December 2006: Case study of a
 602 tornadic storm in a low CAPE, high shear environment. *Atmos. Res.*, **93**, 50–65,
 603 doi:10.1016/j.atmosres.2008.10.008.

604 Clark, M. R., 2011: Doppler radar observations of mesovortices within a cool-season tornadic
 605 squall line over the UK. *Atmos. Res.*, **100**, 749–764, doi:10.1016/j.atmosres.2010.09.007.

606 Clark, M. R., 2013: A provisional climatology of cool-season convective lines in the UK. *Atmos.*
 607 *Res.*, **123**, 180–196, doi:10.1016/j.atmosres.2012.09.018.

608 Clark, M. R., and D. J. Parker, 2014: On the mesoscale structure of surface wind and pressure
 609 fields near tornadic and nontornadic cold fronts. *Mon. Wea. Rev.*, **142**, 3560– 3585, doi:
 610 10.1175/MWR-D-13-00395.1.

611 Craven, J. P., and H. E Brooks, 2004: Baseline climatology of sounding derived parameters
 612 associated with deep moist convection. *Natl. Wea. Dig.*, **28**, 13–24.

613 Doe, R. K., Ed., 2016: *Extreme Weather: Forty Years of the Tornado and Storm Research*
 614 *Organization (TORRO)*. Wiley Blackwell, 327 pp.

615 Elsom, D. M., G. T. Meaden, D. J. Reynolds, M. W. Rowe, and J. D. C. Webb, 2001: Advances
 616 in tornado and storm research in the United Kingdom and Europe: The role of the Tornado
 617 and Storm Research Organisation. *Atmos. Res.*, **56**, 19–29, doi:10.1016/S0169-
 618 8095(00)00084-3.

619 Fujita, T., 1981: Tornadoes and downbursts in the context of generalized planetary scales. *J.*
 620 *Atmos. Sci.*, **38**, 1511–1534.

621 Groenemeijer, P., U. Corsmeier, and Ch. Kottmeier, 2011: The development of tornadic storms
 622 on the cold side of a front favoured by local enhancement of moisture and CAPE. *Atmos.*
 623 *Res.*, **100**, 765–781, doi:10.1016/j.atmosres.2010.10.028.

624 Hagan, M., 1992: On the appearance of a cold front with a narrow rainband in the vicinity of the
 625 Alps. *Meteor. Atmos. Phys.*, **48**, 231–248.

626 Hobbs, P. V., and K. R. Biswas, 1979: The cellular nature of narrow cold-frontal rainbands.
627 *Quart. J. Roy. Meteor. Soc.*, **105**, 723–727.

628 Hobbs, P. V., and P. O. G. Persson, 1982: The mesoscale and microscale structure and
629 organization of clouds and precipitation in midlatitude cyclones. Part V: The substructure of
630 narrow cold-frontal rainbands. *J. Atmos. Sci.*, **39**, 280–295.

631 Holden, J., and A. Wright, 2004: Tornado climatology and the development of simple prediction
632 tools. *Quart. J. Roy. Meteor. Soc.*, **130**, 1009–1021, doi: 10.1256/qj.03.45.

633 Houze, R. A., Jr., 2014: *Cloud Dynamics*, 2nd ed. Academic Press, 496 pp.

634 James, P. K., and K. A. Browning, 1979: Mesoscale structure of line convection at surface cold
635 fronts. *Quart. J. Roy. Meteor. Soc.*, **105**, 371–382.

636 Janjić, Z. I., 1994: The step-mountain eta coordinate model: Further developments of the
637 convection, viscous sublayer and turbulence closure schemes. *Mon. Wea. Rev.*, **122**, 927–
638 945, doi: 10.1175/1520-0493(1994)122<0927:TSMECM>2.0.CO;2.

639 Janjić, Z. I., 2002: Nonsingular implementation of the Mellor–Yamada level 2.5 scheme in the
640 NCEP Meso model. NCEP Office Note, No. 437, 61 pp.

641 Jewett, B. F., and R. B. Wilhelmson, 2006: The role of forcing in cell morphology and evolution
642 within midlatitude squall lines. *Mon. Wea. Rev.*, **134**, 3714–3734.

643 Johns, R. H., and C. A. Doswell III, 1992: Severe local storms forecasting. *Wea. Forecasting*, **7**,
644 588–612, doi: 10.1175/1520-0434(1992)007<0588:SLSF>2.0.CO;2.

645 Jorgensen, D. P., Z. Pu, P. O. G. Persson, and W. Tao, 2003: Variations associated with cores
646 and gaps of a Pacific narrow cold frontal rainband. *Mon. Wea. Rev.*, **131**, 2705–2729, doi:
647 10.1175/1520-0493(2003)131<2705:VAWCAG>2.0.CO;2.

648 Kain, J. S., 2004: The Kain–Fritsch convective parameterization: An update. *J. Appl. Meteor.*,
649 **43**, 170–181, doi: 10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2.

650 Kain, J. S., and J. M. Fritsch, 1990: A one-dimensional entraining/detraining plume model and
651 its application in convective parameterization. *J. Atmos. Sci.*, **47**, 2784–2802, doi:
652 10.1175/1520-0469(1990)047<2784:AODEPM>2.0.CO;2.

653 Kalnay, E., and Coauthors, 1996: The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Amer.*
654 *Meteor. Soc.*, **77**, 437–471.

655 Kawashima, M., 2011: Numerical study of horizontal shear instability waves along narrow cold
656 frontal rainbands. *J. Atmos. Sci.*, **68**, 878–903.

657 Kemp, A. K., and S. J. Morris, 1982: Line squall and minor tornadoes at Holyhead, 23
658 November 1981. *Meteor. Mag.*, **111**, 253–261.

659 Kirk, P. J., 2014: An updated tornado climatology for the UK: 1981–2010. *Weather*, **69**, 171–
660 175.

661 Kobayashi, F., Y. Sugawara, M. Imai, M. Matsui, A. Yoshida, and Y. Tamura, 2007: Tornado
662 generation in a narrow cold frontal rainband—Fujisawa tornado on April 20, 2006—. *SOLA*,
663 **3**, 21–24.

664 Lee, B. D., and R. B. Wilhelmson, 1997a: The numerical simulation of non-supercell
665 tornadogenesis. Part I: Initiation and evolution of pretornadic misocyclone and circulations
666 along a dry outflow boundary. *J. Atmos. Sci.*, **54**, 32–60.

667 Lee, B. D., and R. B. Wilhelmson, 1997b: The numerical simulation of nonsupercell
668 tornadogenesis. Part II: Evolution of a family of tornadoes along a weak outflow boundary. *J.*
669 *Atmos. Sci.*, **54**, 2387–2415.

670 Lee, B. D., and R. B. Wilhelmson, 2000: The numerical simulation of nonsupercell
 671 tornadogenesis. Part III: Parameter tests investigating the role of CAPE, vortex sheet
 672 strength, and boundary layer vertical shear. *J. Atmos. Sci.*, **57**, 2246–2261.

673 Locatelli, J. D., J. E. Martin, and P. V. Hobbs, 1995: Development and propagation of
 674 precipitation cores on cold fronts. *Atmos. Res.*, **38**, 177–206.

675 Markowski, P. M., J. M. Straka, and E. N. Rasmussen, 2002: Direct surface thermodynamic
 676 observations within the rear-flank downdrafts of nontornadic and tornadic supercells. *Mon.*
 677 *Wea. Rev.*, **130**, 1692–1721, doi: 10.1175/1520-0493(2002)130<1692:DSTOWT>2.0.CO;2.

678 McAvoy, B. P., W. A. Jones, and P. D. Moore, 2000: Investigation of an unusual storm structure
 679 associated with weak to occasionally strong tornadoes over the eastern United States.
 680 Preprints, *20th Severe Local Storms Conference*, Orlando, FL, Amer. Meteor. Soc., 182–185.

681 Meaden, G. T., 1983: The TORRO tornado intensity scale. *J. Meteor. (UK)*, **8**, 151–153.

682 Meaden, G. T., S. Kochev, L. Kolendowicz, A. Kosa-Kiss, I. Marcinonienė, M. Sioutas, H.
 683 Tooming, and J., Tyrrell, 2007: Comparing the theoretical versions of the Beaufort scale, the
 684 T-Scale and the Fujita scale. *Atmos. Res.*, **83**, 446–449, doi:10.1016/j.atmosres.2005.11.014.

685 Mellor, G. L., and T. Yamada, 1982: Development of a turbulence closure model for geophysical
 686 fluid problems. *Rev. Geophys. Space Phys.*, **20**, 851–875.

687 Morrison, H., G. Thompson, and V. Tatarskii, 2009: Impact of cloud microphysics on the
 688 development of trailing stratiform precipitation in a simulated squall line: Comparison of
 689 one- and two-moment schemes. *Mon. Wea. Rev.*, **137**, 991–1007.

690 Mulder, K. J., 2015: Tornadoes in the British Isles: Climatology, formation environments, and
 691 storm dynamics. Ph.D. dissertation, University of Manchester, 96 pp.

692 Mulder, K. J., and D. M. Schultz, 2015: Climatology, storm morphologies, and environments of
693 tornadoes in the British Isles: 1980–2012. *Mon. Wea. Rev.*, **143**, 2224–2240.

694 Nolen, R. H., 1959: A radar pattern associated with tornadoes. *Bull. Amer. Meteor. Soc.*, **40**,
695 277–279.

696 Potvin, C. K., K. L. Elmore, and S. J. Weiss, 2010: Assessing the impacts of proximity sounding
697 criteria on the climatology of significant tornado environments. *Wea. Forecasting*, **25**, 921–
698 930.

699 Púčik, T., P. Groenemeijer, D. Rýva, and M. Kolář, 2015: Proximity soundings of severe and
700 nonsevere thunderstorms in Central Europe. *Mon. Wea. Rev.*, **143**, 4805–4821.

701 Rasmussen, E. N., and D. O. Blanchard, 1998: A baseline climatology of sounding-derived
702 supercell and tornado forecast parameters. *Wea. Forecasting*, **13**, 1148–1164.

703 Roebber, P. J., D. M. Schultz, and R. Romero, 2002: Synoptic regulation of the 3 May 1999
704 tornado outbreak. *Wea. Forecasting*, **17**, 399–429.

705 Rowe, M. W., 1985: Britain’s greatest tornadoes and tornado outbreak. *J. Meteor. (UK)*, **10**,
706 212–220.

707 Rowe, M. W., 2016: Tornado extremes in the United Kingdom: The earliest, longest, widest,
708 severest, and deadliest. *Extreme Weather: Forty Years of the Tornado and Storm Research*
709 *Organization (TORRO)*. R. K. Doe, Ed., Wiley Blackwell, 77–90.

710 Rowe, M. W., and G. T. Meaden, 1985: Britain’s greatest tornado outbreak. *Weather*, **40**, 230–
711 235, doi: 10.1002/j.1477-8696.1985.tb06883.x.

712 Skamarock, W. C., J. B. Klemp, J. Dudhia, D. O. Gill, D. M. Barker, M. G. Duda, X.-Y. Huang,
 713 W. Wang, and J. G. Powers, 2008: A description of the Advanced Research WRF Version 3.
 714 NCAR technical note, NCAR/TN-475+STR, 113 pp.

715 Smart, D. J., and K. A. Browning, 2009: Morphology and evolution of cold-frontal
 716 misocyclones. *Quart. J. Roy. Meteor. Soc.*, **135**, 381–393, doi: 10.1002/qj.399.

717 Smith, B. T., R. L. Thompson, J. S. Grams, C. Broyles, and H. E. Brooks, 2012: Convective
 718 modes for significant severe thunderstorms in the contiguous United States. Part I: Storm
 719 classification and climatology. *Wea. Forecasting*, **27**, 1114–1135, doi: [10.1175/WAF-D-11-](https://doi.org/10.1175/WAF-D-11-00115.1)
 720 [00115.1](https://doi.org/10.1175/WAF-D-11-00115.1).

721 Speheger, D. A., C. A. Doswell III, and G. J. Stumpf, 2002: The tornadoes of 3 May 1999:
 722 Event verification in central Oklahoma and related issues. *Wea. Forecasting*, **17**, 362–381.

723 Sugawara, Y., and F. Kobayashi, 2009: Vertical structure of misocyclones along a narrow cold
 724 frontal rainband. *J. Meteor. Soc. Japan*, **87**, 497–503.

725 Thompson, G., P. R. Field, R. M. Rasmussen, and W. D. Hall, 2008: Explicit forecasts of winter
 726 precipitation using an improved bulk microphysics scheme. Part II: Implementation of a new
 727 snow parameterization. *Mon. Wea. Rev.*, **136**, 5095–5115, doi:10.1175/2008MWR2387.1.

728 Thompson, R. L., R. Edwards, J. A. Hart, K. L. Elmore, and P. Markowski, 2003: Close
 729 proximity soundings within supercell environments obtained from the Rapid Update Cycle.
 730 *Wea. Forecasting*, **18**, 1243–1261.

731 Thompson, R. L., B. T. Smith, J. S. Grams, A. R. Dean, and C. Broyles, 2012: Convective modes
 732 for significant severe thunderstorms in the contiguous United States. Part II: Supercell and
 733 QLCS tornado environments. *Wea. Forecasting*, **27**, 1136–1154.

734 Trapp, R. J., E. D. Mitchell, G. A. Tipton, D. W. Effertz, A. I. Watson, D. L. Andra Jr., and M.
 735 A. Magsig, 1999: Descending and non-descending tornadic vortex signatures detected by
 736 WSR-88Ds. *Wea. Forecasting*, **14**, 625–639.

737 Trapp, R. J., S. A. Tessendorf, E. S. Godfrey, and H. E. Brooks, 2005: Tornadoes from squall
 738 lines and bow echoes. Part I: Climatological distribution. *Wea. Forecasting*, **20**, 23–34, doi:
 739 10.1175/WAF-835.1.

740 Trapp, R. J., and M. L. Weisman, 2003: Low-level mesovortices within squall lines and bow
 741 echoes. Part II: Their genesis and implications. *Mon. Wea. Rev.*, **131**, 2804–2823.

742 Trapp, R. J., D. M. Wheatley, N. T. Atkins, R. W. Przybylinski, and R. Wolf, 2006: Buyer
 743 beware: Some words of caution on the use of severe wind reports in postevent assessment
 744 and research. *Wea. Forecasting*, **21**, 408–415.

745 Turner, S., Elsom, D. M., and G. T. Meaden, 1986: An outbreak of 31 tornadoes associated with
 746 a cold front in southern England on 20 October 1981. *J. Meteor. (UK)*, **11**, 37–50.

747 Viale, M., R. A. Houze Jr., and K. L. Rasmussen, 2013: Upstream orographic enhancement of a
 748 narrow cold-frontal rainband approaching the Andes. *Mon. Wea. Rev.*, **141**, 1708–1730.

749 Wakimoto, R. M., and B. L. Bosart, 2000: Airborne radar observations of a cold front during
 750 FASTEX. *Mon. Wea. Rev.*, **128**, 2447–2470.

751 Wakimoto, R. M., and J. W. Wilson, 1989: Non-supercell tornadoes. *Mon. Wea. Rev.*, **117**,
 752 1113–1140.

753 Weisman, M. L., and R. J. Trapp, 2003: Low-level meso-vortices within squall lines and bow
 754 echoes. Part I: Overview and dependence on environmental shear. *Mon. Wea. Rev.*, **131**,
 755 2779–2803.

Wheatley, D. M., and R. J. Trapp, 2008: The effect of mesoscale heterogeneity on the genesis and structure of mesovortices within quasi-linear convective systems. *Mon. Wea. Rev.*, **136**, 4220–4241.

FIGURE CAPTIONS

Figure 1. Locations of the 104 tornado reports from the TORRO database for 23 November 1981. Numbers represent their strength on the T scale; U represents unknown intensity, and half-values represent intensities between two classes (e.g., 2.5 represents T2–T3). Reports verified by TORRO (58) are classified as definite and plotted in black. Reports that have not been verified (46) are classified as probable and are plotted in red. Locations discussed in the text are labeled in blue. Locations of reports that appear to be located over water are a result of a coarse representation of geography.

Figure 2. Distribution on the T scale of intensities of the 99 tornado reports on 23 November 1981 associated with an intensity rating from the TORRO database.

Figure 3. Excerpts from Met Office Daily Weather Summary surface weather charts at (a) 1200 UTC 22 November 1981 and (b) 1200 UTC 23 November 1981. Plotted are sea level pressure contours every 4 hPa, surface fronts, surface temperatures (°C) and weather at selected cities, and occasionally wind barbs (standard notation). Crown copyright.

Figure 4. Excerpt from Met Office Daily Weather Summary 500-hPa chart at 1200 UTC 23 November 1981. Plotted are 500-hPa geopotential height (solid lines every 6 dam) and 1000–500-hPa thickness (dashed lines every 6 dam). Crown copyright.

777 Figure 5. Excerpts from the Met Office Central Forecasting Office hourly U.K. working charts at
778 (a) 1000 UTC, (b) 1200 UTC, and 1400 UTC 23 November 1981. Plotted are sea-level pressure
779 (solid lines every 2 hPa in (a) and (c) and 4 hPa in (b)), cold front (dashed line), and standard
780 station models. Crown copyright.

781 Figure 6. Hourly rainfall amounts (mm) from 213 rain gauges ending at 1200 UTC and 212 rain
782 gauges ending at 1400 UTC 23 Nov 1981.

783 Figure 7. Infrared satellite imagery (channel 5, 11.5–12.5 μm) at 1325 UTC 23 Nov 1981
784 (courtesy of Dundee Satellite Receiving Station).

785 Figure 8. Prefrontal sounding from Aughton, near Liverpool, at 0000 UTC 23 November 1981
786 (courtesy of the University of Wyoming, <http://weather.uwyo.edu/upperair/sounding.html>).

787 Figure 9. The two innermost domains used in this simulation.

788 Figure 10. Simulation of sea level pressure (hPa, blue lines), surface temperature ($^{\circ}\text{C}$, colored
789 according to scale), and surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} ,
790 respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km
791 horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.

792 Figure 11. Simulation of radar reflectivity factor (dBZ, colored according to scale) and surface
793 winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation between
794 displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000
795 UTC and (b) 1400 UTC 23 November 1981.

796 Figure 12. Simulation of absolute vorticity at 500 m ASL (s^{-1} , colored according to scale) every
797 30 min from 0930 UTC to 1730 UTC (labeled every hour) on the domain with 1-km horizontal
798 grid spacing. The red box indicates the location of the domain with 200-m horizontal grid
799 spacing.

800 Figure 13. Simulation of CAPE (J kg^{-1} , colored according to scale) and surface winds (pennant,
801 full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation between displayed wind
802 vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400
803 UTC 23 November 1981.

804 Figure 14. Simulation of lifting condensation level (LCL) (m, colored according to scale) and
805 surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation
806 between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a)
807 1000 UTC and (b) 1400 UTC 23 November 1981.

808 Figure 15. Simulation of 0–1-km vertical shear of the horizontal wind in magnitude (m s^{-1} ,
809 colored according to scale) and direction (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} ,
810 respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km
811 horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.

812 Figure 16. Simulation of 0–1-km storm-relative helicity ($\text{m}^2 \text{s}^{-2}$, colored according to scale) and
813 surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation
814 between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a)
815 1000 UTC and (b) 1400 UTC 23 November 1981.

816 Figure 17. Simulation of radar reflectivity factor (black lines every 10 dBZ), absolute vorticity at
817 500 m ASL (positive values are contoured in dark blue solid lines every 0.005 s^{-1} , starting from
818 0.01 s^{-1} ; negative values are contoured in light blue solid lines every -0.005 s^{-1} every 0.005 s^{-1}),
819 500-m updrafts (red fill above 5 m s^{-1}), and 500-m downdrafts (green fill above 2 m s^{-1}) from the
820 200-m horizontal grid spacing domain, plotted every minute from 1431:50 to 1434:50 UTC 23
821 November 1981.

822 Figure 18. Characteristic structure and evolution of a simulated misovortex within the domain at
823 200-m horizontal grid spacing, plotted every 60 s around the time that it matures: radar
824 reflectivity factor (dBZ, colored according to scale in Fig. 16), absolute vorticity at 500 m ASL
825 (black contours every 0.005 s^{-1} , starting from 0.01 s^{-1}), 500-m updraft (red contours every 5 m s^{-1})
826 1), and 500-m downdraft (pink contours every 2 m s^{-1}). Each panel is about $4 \text{ km} \times 4 \text{ km}$, and
827 the vortex is about 500 m across.

828 Figure 19. Locations of the 90 revised tornado reports from the TORRO database for 23
829 November 1981. Numbers represent their strength on the T scale. Reports verified by TORRO
830 (58) are classified as definite and plotted in black. Reports that have not been verified (32) are
831 classified as probable and are plotted in red. Locations discussed in the text are labeled in blue.
832 The red box indicates the location of the domain with 200-m horizontal grid spacing. Locations
833 of reports that appear to be located over water are a result of a coarse representation of
834 geography.

835 Figure 20. Tracks of (a) 500-m absolute vorticity (0.02 and 0.025 s^{-1}), (b) 500-m updrafts (5 , 6 ,
836 and 7 m s^{-1}), and (c) 500-m downdrafts (2 , 3 , and 4 m s^{-1}) plotted every minute from 1300 to
837 1600 UTC in the domain with 200-m horizontal grid spacing.

838 Figure 21. Simulation of 0.02 s^{-1} and 0.025 s^{-1} absolute vorticity at 500 m ASL (black contours),
839 5 m s^{-1} updrafts at 500 m (red contours), and 3 m s^{-1} downdrafts at 500 m (green contours) from
840 1300 to 1600 UTC on the domain with 200-m horizontal grid spacing.

841 Figure 22. Simulation of absolute vorticity at 500 m ASL (s^{-1} , colored according to scale) every
842 30 min from 0930 UTC to 1730 UTC on the domain with 1-km horizontal grid spacing. Purple
843 lines separate approximate areas with simulated absolute vorticity less than 0.002 s^{-1} on the 1-km
844 domain during the time of frontal passage. Blue lines separate approximate areas with simulated
845 positive CAPE during the time of frontal passage. Locations of reports that appear to be located
846 over water are a result of a coarse representation of geography. Locations of the 90 tornado
847 reports from the TORRO database for 23 November 1981. Numbers represent their strength on
848 the T scale. Reports verified by TORRO (58) are classified as definite and plotted in black.
849 Reports that have not been verified (32) are classified as probable and are plotted in red.


850

Table 1: Properties of tracks of 500-m absolute vorticity (0.02 and 0.025 s^{-1}), updrafts (5 , 6 , and 7 ms^{-1}), and downdrafts (2 , 3 , and 4 m s^{-1}) in the domain with 200-m horizontal grid spacing between 1300 and 1600 UTC. “Long-lived” refers to features lasting 30 min or more (± 2 min because the data interval is every minute). The longest duration track being listed as “101+” means that a track started within the plotting domain but continued to the edge of the domain, indicating that the track could have existed longer than 101 min. “N/A” represents no features meeting the designated criteria.

	Number of long-lived maxima	Average duration of long-lived tracks (min)	Median duration of long-lived tracks (min)	Longest duration track (min)	Longest track length (to nearest 5 km)
Vorticity ($>0.025 \text{ s}^{-1}$) tracks	9	33.2	33	38	45
Vorticity ($>0.02 \text{ s}^{-1}$) tracks	41	40.3	37	64	75
Updrafts ($> 5 \text{ m/s}$) tracks	23	55.3	48	101+	175
Updrafts ($> 6 \text{ m/s}$) tracks	5	54.2	56	76	100
Updraft ($> 7 \text{ m/s}$) tracks	0	N/A	N/A	28	35
Downdrafts ($> 2 \text{ m/s}$) tracks	10	34.2	33.5	40	75
Downdrafts ($> 3 \text{ m/s}$) tracks	0	N/A	N/A	22	35
Downdrafts ($> 4 \text{ m/s}$) tracks	0	N/A	N/A	12	20

858

Locations of 104 Tornado Reports for 23 Nov 1981



860

861

Figure 1. Locations of the 104 tornado reports from the TORRO database for 23 November

862

1981. Numbers represent their strength on the T scale; U represents unknown intensity, and

863

half-values represent intensities between two classes (e.g., 2.5 represents T2–T3). Reports

864

verified by TORRO (58) are classified as definite and plotted in black. Reports that have not

865

been verified (46) are classified as probable and are plotted in red. Locations discussed in the

866

text are labeled in blue. Locations of reports that appear to be located over water are a result of a

867

coarse representation of geography.

868




Figure 2. Distribution on the T scale of intensities of the 99 tornado reports on 23 November 1981 associated with an intensity rating from the TORRO database.






Figure 3. Excerpts from Met Office Daily Weather Summary surface weather charts at (a) 1200 UTC 22 November 1981 and (b) 1200 UTC 23 November 1981. Plotted are sea level pressure contours every 4 hPa, surface fronts, surface temperatures ($^{\circ}\text{C}$) and weather at selected cities, and occasionally wind barbs (standard notation). Crown copyright.



879


880 Figure 4. Excerpt from Met Office Daily Weather Summary 500-hPa chart at 1200 UTC 23
 881 November 1981. Plotted are 500-hPa geopotential height (solid lines every 6 dam) and 1000–
 882 500-hPa thickness (dashed lines every 6 dam). Crown copyright.

883




884

885



886


887



888

889 Figure 5. Excerpts from the Met Office Central Forecasting Office hourly U.K. working charts at
 890 (a) 1000 UTC, (b) 1200 UTC, and 1400 UTC 23 November 1981. Plotted are sea-level pressure
 891 (solid lines every 2 hPa in (a) and (c) and 4 hPa in (b)), cold front (dashed line), and standard
 892 station models. Crown copyright.


893



894

895 Figure 6. Hourly rainfall amounts (mm) from 213 rain gauges ending at 1200 UTC and 212 rain
 896 gauges ending at 1400 UTC 23 Nov 1981.

897



898

899 Figure 7. Infrared satellite imagery (channel 5, 11.5–12.5 μm) at 1325 UTC 23 Nov 1981
900 (courtesy of Dundee Satellite Receiving Station).


901

902

903

904

905



906

907

908

Figure 8. Prefrontal sounding from Aughton, near Liverpool, at 0000 UTC 23 November 1981

909

(courtesy of the University of Wyoming, <http://weather.uwyo.edu/upperair/sounding.html>).

910





Figure 9. The two innermost domains used in this simulation.

914



915

916

917

918

919

920

Figure 10. Simulation of sea level pressure (hPa, blue lines), surface temperature (°C, colored according to scale), and surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s⁻¹, respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.

920




Figure 11. Simulation of radar reflectivity factor (dBZ, colored according to scale) and surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s⁻¹, respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.





Figure 12. Simulation of absolute vorticity at 500 m ASL (s^{-1} , colored according to scale) every 30 min from 0930 UTC to 1730 UTC (labeled every hour) on the domain with 1-km horizontal grid spacing. The red box indicates the location of the domain with 200-m horizontal grid spacing.

933



934

935 Figure 13. Simulation of CAPE (J kg^{-1} , colored according to scale) and surface winds
936 (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation between
937 displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000
938 UTC and (b) 1400 UTC 23 November 1981.

939




Figure 14. Simulation of lifting condensation level (LCL) (m, colored according to scale) and surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s⁻¹, respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.




Figure 15. Simulation of 0–1-km vertical shear of the horizontal wind in magnitude (m s^{-1} , colored according to scale) and direction (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.




Figure 16. Simulation of 0–1-km storm-relative helicity ($\text{m}^2 \text{s}^{-2}$, colored according to scale) and surface winds (pennant, full barb, and half-barb denote 25, 5, 2.5 m s^{-1} , respectively; separation between displayed wind vectors is 30 km) on the domain with 1-km horizontal grid spacing at (a) 1000 UTC and (b) 1400 UTC 23 November 1981.




Figure 17. Simulation of radar reflectivity factor (black lines every 10 dBZ), absolute vorticity at 500 m ASL (positive values are contoured in dark blue solid lines every 0.005 s^{-1} , starting from 0.01 s^{-1} ; negative values are contoured in light blue solid lines every -0.005 s^{-1} every 0.005 s^{-1}), 500-m updrafts (red fill above 5 m s^{-1}), and 500-m downdrafts (green fill above 2 m s^{-1}) from the 200-m horizontal grid spacing domain, plotted every minute from 1431:50 to 1434:50 UTC 23 November 1981.



Figure 18. Characteristic structure and evolution of a simulated misovortex within the domain at 200-m horizontal grid spacing, plotted every 60 s around the time that it matures: radar reflectivity factor (dBZ, colored according to scale in Fig. 16), absolute vorticity at 500 m ASL (black contours every 0.005 s^{-1} , starting from 0.01 s^{-1}), 500-m updraft (red contours every 5 m s^{-1}), and 500-m downdraft (pink contours every 2 m s^{-1}). Each panel is about $4 \text{ km} \times 4 \text{ km}$, and the vortex is about 500 m across.

Locations of 90 Revised Tornado Reports




Figure 19. Locations of the 90 revised tornado reports from the TORRO database for 23 November 1981. Numbers represent their strength on the T scale. Reports verified by TORRO (58) are classified as definite and plotted in black. Reports that have not been verified (32) are classified as probable and are plotted in red. Locations discussed in the text are labeled in blue. The red box indicates the location of the domain with 200-m horizontal grid spacing. Locations of reports that appear to be located over water are a result of a coarse representation of geography.




Figure 20. Tracks of (a) 500-m absolute vorticity (0.02 and 0.025 s^{-1}), (b) 500-m updrafts (5 , 6 , and 7 m s^{-1}), and (c) 500-m downdrafts (2 , 3 , and 4 m s^{-1}) plotted every minute from 1300 to 1600 UTC in the domain with 200-m horizontal grid spacing.




Figure 21. Simulation of 0.02 s^{-1} and 0.025 s^{-1} absolute vorticity at 500 m ASL (black contours), 5 m s^{-1} updrafts at 500 m (red contours), and 3 m s^{-1} downdrafts at 500 m (green contours) from 1300 to 1600 UTC on the domain with 200-m horizontal grid spacing.




Figure 22. Simulation of absolute vorticity at 500 m ASL (s^{-1} , colored according to scale) every 30 min from 0930 UTC to 1730 UTC on the domain with 1-km horizontal grid spacing. Purple lines separate approximate areas with simulated absolute vorticity less than 0.002 s^{-1} on the 1-km domain during the time of frontal passage. Blue lines separate approximate areas with simulated positive CAPE during the time of frontal passage. Locations of reports that appear to be located over water are a result of a coarse representation of geography. Locations of the 90 tornado reports from the TORRO database for 23 November 1981. Numbers represent their strength on the T scale. Reports verified by TORRO (58) are classified as definite and plotted in black. Reports that have not been verified (32) are classified as probable and are plotted in red.