

*The effect of the apolipoprotein E
genotype on response to personalized
dietary advice intervention: findings from
the Food4Me randomized controlled trial*

Article

Accepted Version

Fallaize, R., Celis-Morales, C., Macready, A. L. ORCID: <https://orcid.org/0000-0003-0368-9336>, Marsaux, C. F. M., Forster, H., O'Donovan, C., Woolhead, C., San-Cristobal, R., Kolossa, S., Hallmann, J., Mavrogianni, C., Surwillo, A., Livingstone, K. M., Moschonis, G., Navas-Carretero, S., Walsh, M. C., Gibney, E. R., Brennan, L., Bouwman, J., Grimaldi, K., Manios, Y., Traczyk, I., Drevon, C. A., Martinez, J. A., Daniel, H., Saris, W. H. M., Gibney, M. J., Mathers, J. C. and Lovegrove, J. A. ORCID: <https://orcid.org/0000-0001-7633-9455> (2016) The effect of the apolipoprotein E genotype on response to personalized dietary advice intervention: findings from the Food4Me randomized controlled trial. *American Journal of Clinical Nutrition*, 104 (3). pp. 827-836. ISSN 0002-9165 doi: 10.3945/ajcn.116.135012 Available at <https://centaur.reading.ac.uk/66411/>

To link to this article DOI: <http://dx.doi.org/10.3945/ajcn.116.135012>

Publisher: American Society for Nutrition

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



The American Journal of Clinical Nutrition
Official publication of the American Society for Nutrition

The American Journal of Clinical Nutrition
AJCN/2016/135012
Version 2

The effect of APOE genotype on response to personalized dietary intervention: findings from the Food4Me randomized controlled trial

Corresponding Author: *Julie A Lovegrove*

Additional Authors: *Rosalind Fallaize* , *Carlos Celis-Morales* , *Anna L Macready* , *Cyril FM Marsaux* , *Hannah Forster* , *Clare O'Donovan* , *Clara Woodhead* , *Rodrigo San-Cristobal* , *Silvia Kolossa* , *Jacqueline Hallmann* , *Christina Mavrogianni* , *Agnieszka Surwillo* , *Katherine M Livingstone* , *George Moschonis* , *Santiago Navas-Carretero* , *Marianne C Walsh* , *Eileen R Gibney* , *Lorraine Brennan* , *Jildau Bouwman* , *Keith Grimaldi* , *Yannis Manios* , *Iwona Traczyk* , *Christian A Drevon* , *J Alfredo Martinez* , *Hannelore Daniel* , *Wim HM Saris* , *Michael J Gibney* , *John C Mathers*

Date Received: 8 Jun 2016

Trial registered: ClinicalTrials.gov Reg No. NCT01530139

Information for Authors: <http://www.ajcn.org/site/misc/ifa.xhtml>

**The effect of APOE genotype on response to personalized dietary advice
intervention: findings from the Food4Me randomized controlled trial¹⁻²**

Rosalind Fallaize³, Carlos Celis-Morales⁴, Anna L. Macready³, Cyril F.M. Marsaux⁵, Hannah Forster⁶, Clare O'Donovan⁶, Clara Woolhead⁶, Rodrigo San-Cristobal⁷, Silvia Kolossa⁸, Jacqueline Hallmann⁸, Christina Mavrogianni⁹, Agnieszka Surwillo¹⁰, Katherine M. Livingstone⁴, George Moschonis⁹, Santiago Navas-Carretero⁷, Marianne C. Walsh⁶, Eileen R. Gibney⁶, Lorraine Brennan⁶, Jildau Bouwman¹¹, Keith Grimaldi¹², Yannis Manios⁹, Iwona Traczyk¹⁰, Christian A. Drevon¹³, J. Alfredo Martinez⁷, Hannelore Daniel⁸, Wim H.M. Saris⁵, Michael J. Gibney⁶, John C. Mathers and Julie A. Lovegrove³, on behalf of the Food4Me Study

Footnotes

¹ Supported by grant 265494 from the European Commission under the Food, Agriculture, Fisheries and Biotechnology Theme of the 7th Framework Programme for Research and Technological Development

² Address correspondence to JA Lovegrove, Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, University of Reading, Whiteknights, Reading RG6 6AP, United Kingdom. Tel: +44 (0) 118 378 6418; Fax: +44 (0)118 931 0080; Email: j.a.lovegrove@reading.ac.uk

Author affiliations

³ Hugh Sinclair Unit of Human Nutrition and Institute for Cardiovascular and Metabolic Research, University of Reading, Reading; ⁴Human Nutrition Research Centre, Institute of Cellular Medicine, Newcastle University, Newcastle Upon Tyne, UK; ⁵Department of Human Biology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre +, Maastricht, The

Netherlands; ⁶UCD Institute of Food and Health, University College Dublin, Belfield, Dublin 4, Ireland; ⁷Department of Nutrition, Food Science and Physiology, University of Navarra, Pamplona, IdiSNA, Instituto de Investigación Sanitaria de Navarra, Pamplona and CIBERObn, Fisiopatología de la Obesidad y Nutrición, Instituto de Salud Carlos III, Madrid Spain; ⁸ZIEL Research Center of Nutrition and Food Sciences, Biochemistry Unit, Technische Universität München, München, Germany; ⁹Department of Nutrition and Dietetics, Harokopio University, Athens, Greece; ¹⁰National Food & Nutrition Institute (IZZ), Warsaw, Poland; ¹¹Microbiology and Systems Biology Group, TNO, Utrechtseweg 48, 3704 HE Zeist, The Netherlands; ¹²Eurogenetica Ltd. 7 Salisbury Road, Burnham-on-Sea, UK, ¹³Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway.

PubMed indexing: Fallaize, Celis-Morales, Macready, Marsaux, Forster, O'Donovan, Woolhead, San-Cristobal, Kolossa, Hallmann, Mavrogianni, Surwillo, Livingstone, Moschonis, Navas-Carretero, Walsh, Gibney, Brennan, Bouwman, Grimaldi, Manios, Traczyk, Drevon, Martinez, Daniel, Saris, Gibney, Mathers and Lovegrove

Key Words: APOE, Nutrigenomics, Food4Me, Dietary Fat, Personalized Nutrition

Abbreviations: BCT, behavioral change technique; BMI, body mass index; CHD, coronary heart disease; DBS, dried blood spot; DHA, docosahexanoic acid; EPA, eicosapentanoic acid; GLM, general linear model; LDL-C, low-density lipoprotein cholesterol; MetS, metabolic syndrome; MUFA, monounsaturated fatty acid; PA, physical activity; PN, personalized nutrition; RCT, randomised controlled trial; SFA, saturated fatty acid; TC, total cholesterol; %TE, % total energy

Running Head: APOE response to personalized diet intervention

Clinical trial registration: Clinicaltrials.gov as NCT01530139

Word count (excluding title page and references): 5959

Number of Figures: 1

Number of Tables: 7

1 **ABSTRACT (word count = 299)**

2 **Background:** The *APOE* risk allele ($\epsilon 4$) is associated with higher total cholesterol
3 (TC), amplified response to saturated fatty acid (SFA) reduction and increased CVD.
4 While knowledge of gene 'risk' may enhance dietary change, it is unclear whether $\epsilon 4$
5 carriers would benefit from gene-based personalized nutrition (PN).

6 **Objectives:** The aims of this study were to investigate **interactions between *APOE***
7 **genotype and (a) habitual dietary fat intake and (b) modulations of fat intake** on
8 **metabolic outcomes; (c) determine whether gene-based PN results in greater dietary**
9 **change compared with standard dietary advice (Level 0) and non-gene-based PN**
10 **(Levels 1-2) and (d) assess the impact of knowledge of *APOE* risk (risk: E4+, non-**
11 **risk: E4-) on dietary change following gene-based PN (Level 3).**

12 **Design:** Individuals (n=1466) recruited into the Food4Me pan-European **PN dietary**
13 **intervention** study **were randomized to four treatment arms** and genotyped for *APOE*
14 (rs429358 and rs7412). Diet and dried blood spot TC and omega-3 index were
15 determined at baseline and after 6-months intervention. **Data were analyzed using**
16 **adjusted general linear models.**

17 **Results:** Significantly higher TC concentrations were observed in E4+ participants
18 compared with E4- ($P < 0.05$). Although there were no significant differences in *APOE*
19 response to gene-based PN (E4+ vs. E4-), both groups had a greater reduction in
20 SFA (%TE) intake when compared with Level 0 (E4+, -0.72% vs. -1.95%, $P = 0.035$;
21 E4-, -0.31% vs. -1.68%, $P = 0.029$). Gene-based PN was associated with a smaller
22 reduction in SFA intake compared with non-gene-based PN (Level 2) for E4-
23 participants (-1.68% vs. -2.56%, $P = 0.025$).

24 **Conclusions:** The *APOE* $\epsilon 4$ allele was associated with greater TC. Whilst gene-
25 based PN targeted to *APOE* was more effective in reducing SFA intake than

26 standard dietary advice, there was no difference between *APOE* 'risk' and 'non-risk'
27 groups. Furthermore, disclosure of *APOE* 'non-risk' may have weakened dietary
28 response to PN.

29 INTRODUCTION

30 Coronary heart disease (CHD) is the leading cause of global mortality,
31 accounting for 1 of 5 deaths in Europe (1). Recent estimates suggest that up to 80%
32 of CHD and cerebrovascular disease could be avoided by improving diet and lifestyle
33 (2). While intervention strategies have traditionally used a 'one-size-fits-all' approach
34 to change dietary behaviour, recent evidence suggests that a personalized approach
35 may be more effective (3, 4). Moreover, there has been much interest in the use of
36 genetic information to tailor dietary advice, yet further RCTs are needed to establish
37 the benefit of such advice on sustained dietary changes (5, 6). Of particular interest
38 in relation to CHD risk is the *APOE* genotype.

39 The *APOE* gene is a key regulator of cholesterol and lipid metabolism. *APOE* is
40 polymorphic, with the common *missense* polymorphisms (rs429358 and rs7412)
41 resulting in three alleles, ϵ 2, ϵ 3 and ϵ 4, combining to form 6 haplotypes, E2/E2,
42 E2/E3, E2/E4, E3/E3, E3/E4 and E4/E4. In a sample of 5805 Caucasians, the *APOE*
43 allele frequency for ϵ 2, ϵ 3 and ϵ 4 was 0.08, 0.77 and 0.15 respectively (7). The ϵ 4
44 allele is associated with increased serum total cholesterol (TC), low-density
45 lipoprotein cholesterol (LDL-C) as well as coronary artery disease and mortality (8-
46 12). Estimates of the CHD hazard ratio for E4+ (E3/E4 and E4/E4), compared with
47 E4- (E3/E3), range from 1.06 to 1.42 (8, 9, 11, 13). There is also a growing body of
48 evidence showing that the *APOE* genotype may influence lipid response to dietary
49 fat; data from intervention studies suggest that E4+ participants may be more
50 sensitive to dietary cholesterol, total fat and, in particular, SFA modulation (14, 15).
51 Given their predisposition to CHD, ϵ 4 carriers might benefit from a lower dietary SFA
52 and blood cholesterol (16) and gene-based PN intervention. However, there is a

concern that gene-based PN may reduce motivation for dietary change in individuals without 'risky genes' and undermine current healthy eating messages (17).

The Food4Me study is a pan-European, 6-month, web-based RCT designed to assess the impact of personalizing dietary advice on change in dietary behaviour. Participants were allocated into one of four intervention groups based on standard guidelines (control), dietary intake (level 1), dietary intake and phenotype (level 2) and dietary intake, phenotype and genotype (level 3). Level 3 participants received feedback on four genes: *MTHFR*, *FADS1*, *TCF7L2*, *FTO* and *APOE*.

The aim of the present analysis was to investigate **interactions** between *APOE* genotype and **(a) habitual dietary fat intake and (b) modulations of fat intake on** metabolic outcomes in the Food4Me study, **(c)** assess whether gene-based PN led to greater changes in diet compared with standard dietary advice (control) and non-gene-based PN for E4- and E4+ participants and **(d)** assess the impact of knowledge of *APOE* risk on changes in diet and metabolic outcomes **following** gene-based PN.

PARTICIPANTS AND METHODS

The Food4Me Proof-of-Principle (PoP) study is a 6-month randomized controlled dietary **advice** intervention study conducted in 7 European research centers: University College Dublin, Ireland, University of Reading, UK, Maastricht University, the Netherlands, University of Navarra, Spain, Harokopio University, Greece, National Food and Nutrition Institute, Poland, and Technische Universität München, Germany. The study had a parallel design with 4 intervention arms and was conducted via the web to emulate a web-delivered PN service (www.food4me.org) (18). Ethics approval was granted at each center and digital informed consent was obtained prior to participation. The study was registered at

clinicaltrials.gov (ref. NCT01530139) and was developed following international regulations and the Helsinki Declaration.

Participants

A total of 1,607 participants (aged ≥ 18 years) were recruited to the Food4Me study, as detailed elsewhere (19). Exclusion criteria were: no or limited access to the Internet, following a medically prescribed diet in the past 3 months, or presence of a condition likely to alter dietary requirements e.g. Crohn's disease, coeliac disease, food allergy/intolerance, pregnancy or lactation.

Study design

A randomization scheme, incorporating both gender and age categories (< 45 years and >45 years), was used to allocate participants to one of the four Food4Me intervention groups: Level 0: standard non-personalized dietary and physical activity (PA) advice; Level 1: advice based on dietary intake and PA; Level 2: advice based on dietary intake, PA and phenotype (blood biomarkers) and Level 3: advice based on dietary intake, PA, phenotype and genotype. Detailed recruitment and study procedures are reported elsewhere (19).

Interaction with study participants was conducted remotely via the Food4Me website, by e-mail and post, using standardized operating procedures. A study welcome pack was sent to the participants via post containing: a dried blood spot (DBS) collection kit (Vitas Ltd, Oslo, Norway), an Isohelix SK-1 DNA buccal swab kit (LCG Genomics, Hertfordshire, UK), a TracmorD tri-axial accelerometer (Philips Consumer Lifestyle, The Netherlands; <http://www.directlife.philips.com>), measuring tape and standardized instructions for completion of baseline measurements (m0). On the allocated study day and following an 8-hour overnight fast, participants

collected DBS and buccal swab samples, and measured their height, weight and waist circumference (WC). Questionnaires to be completed on the same day included the validated Food4Me food frequency questionnaire (20, 21) and the validated Baecke physical activity questionnaire (22-24). Participants repeated these measurements, excluding the buccal cell sample, at 3 (m3) and 6 months (m6). The TracmorD tri-axial accelerometer (25) was worn for the entire duration of the study, and data were uploaded on a bi-weekly basis.

Dietary feedback

Following analysis of data collected at m0 and m3, participants received tailored dietary feedback (in their native language) according to their study allocation group.

The dietary feedback provided was based on a pre-defined set of algorithms incorporating dietary, anthropometric, PA, phenotypic and genotypic data where appropriate. The system was designed to ensure consistent feedback across centres and has since been successfully automatized (26). *APOE* gene variants were coded as 'risk' (a genetic variation that can be modified by diet, i.e. E3/E4 or E4/E4 (E4+)) or 'non-risk' (E2/E2, E2/E3, E3/E3 (E4-)). Alongside the risk result, Level 3 participants received the following basic information about the *APOE* genotype: "A specific variation of this gene is associated with a greater need to maintain healthy cholesterol levels. Decreasing saturated fat intake has been associated with an improvement in cholesterol and factors relating to cardiovascular health in these individuals." For Level 3 E4+ participants with high dietary SFA intake and/or high blood TC, who were being advised to lower dietary SFA, reference to 'gene risk' was also included in the advice message, i.e. "You have a genetic variation that can benefit by keeping a healthy intake of saturated fat and a normal level of blood cholesterol."

127 **Biochemical analysis**

128 Participants were asked to complete 2 DBS cards each containing 5 blood
129 spots, at m0, m3 and m6 (approximately 150 μ L blood per card). After drying the
130 blood spots at room temperature for 2-4 hours, the cards were placed in a sealed
131 aluminum bag (Whatman Foil Bags, item no. 10534321, Whatman Inc., Sanford, ME)
132 containing a drying sachet (Sorb-it, item no. 10548234, Süd-Chemie, Germany) and
133 posted back to the research center in their country. Researchers subsequently
134 shipped the DBS cards to Vitas (Vitas Ltd, Norway) for analysis of whole blood TC
135 (LC-UV) and omega-3 index [(eicosapentaenoic acid (EPA) + docosahexaenoic acid
136 (DHA)/ total fatty acids) $\times$ 100] (27). Fatty acids were measured using GC-FID.

137 **DNA extraction and genotyping**

138 Participants were instructed to rub the Isohelix SK-1 DNA buccal swab against
139 the inside of their cheek for one minute before returning it to a plastic tube containing
140 an Isohelix Dri-capsule. Upon return to the center, swabs were shipped to LCG
141 Genomics (LCG Genomics, Hertfordshire, UK) for genotypic analysis. Following DNA
142 extraction, KASPTM genotyping assays were used to provide bi-allelic scoring of
143 polymorphisms in the *APOE* gene (rs429358 and rs7412). Hardy-Weinberg
144 equilibrium for multiple alleles was analyzed, no significant deviation was observed
145 for rs7412 (0.91; $P=1.00$) whereas rs429358 displayed linkage disequilibrium (0.005;
146 $P=0.008$).

147 **Statistical analyses**

148 Data are presented as means $\pm$ SEM. Data were checked for normality of
149 distribution and skewed variables were normalised using Log₁₀ (omega-3 index) and
150 square root (TC) transformations. General linear models (GLM), adjusted for center,

gender, age and body mass index (BMI), were used to assess differences in baseline anthropometric and biochemical values between genotype groups. Habitual **nutrient intake-gene interactions** were assessed using the same GLM model but with the addition of a dietary fat $\times$ genotype interaction term; fat were dichotomised by median intake to assess the impact of the *APOE* genotype on TC and omega-3 index in participants with a similar habitual intake. Post-hoc Bonferroni tests were used to detect specific differences between groups.

Interactions between genotype and dietary **fat** on TC and omega-3 index following dietary **advice** intervention were assessed using % change in dietary fat intake, with 0% used as a reference to dichotomize participants (i.e. reduction vs. increase in fat intake), and then using the resulting groups as fixed factors in the GLM. The interaction term genotype $\times$ change in **fat** was then added to the GLM, with the change in biomarker as the response variable and the respective pre-intervention/ baseline biomarker value as a covariate. The model was adjusted for baseline variables, age, gender, center and weight change [post intervention weight (kg) – pre intervention weight (kg)].

The impact of knowledge of *APOE* risk (risk: E4+, E3E4 and E4/E4; and non-risk: E4-, E2/E2, E2/E3 and E3/E3) on change in diet and TC and omega-3 index (m6-m0) for Level 3 participants advised to lower their SFA at baseline (with high dietary SFA and/or high blood TC) were assessed using GLM. Models were adjusted for baseline variables, age, gender, center and weight change. To assess whether gene-based PN led to greater changes in diet, TC and omega-3 index (m6-m0) than standard dietary advice (Level 0) and non gene-based PN (Levels 1-2), a contrast analysis was performed. Separate analyses were conducted for E4+ (risk) and E4- (non-risk) with Level 3 as the reference group and Levels 0, 1 and 2 as the

comparison groups. As previously, participants with high dietary SFA and/or high blood TC who were advised to lower their SFA at baseline were included and analyses were adjusted for baseline variables, age, gender, center and weight change. Statistical analyses were performed using STATA (version 13.0, StataCorp, TX, USA).

RESULTS

Subject characteristics

A total of 1466 of the 1607 participants randomized into the Food4Me study were genotyped for *APOE* and included in the baseline analysis. Frequency of *APOE* genotype and *APOE* allele according to Food4Me country are presented in **Table 1**. *APOE* E2/E4 participants (n=27) were removed from subsequent analysis due to their low population frequency. Subject characteristics including anthropometry and fasted biomarkers are presented according to *APOE* genotype in **Table 2**. There was no evidence of a genotype-dependant difference in baseline anthropometry, although E4+ participants had higher TC than E4- ($P = 0.040$ for E3/E3 and $P = 0.002$ for E2 carriers).

Habitual dietary and genotype effects at baseline

The associations between dietary fat (total fat, SFA, monounsaturated fatty acids (MUFA), polyunsaturated fatty acids (PUFA) and omega-3), *APOE* genotype, dietary fat $\times$ genotype interactions and TC and omega-3 index, are reported in **Table 3**. Dietary intake was dichotomized at the median (total fat, 35.8%; SFA, 14.0%; MUFA, 13.5%; PUFA, 5.6; omega-3, 0.67%) to determine the effect of specific

199 genotypes in participants with similar habitual dietary fat intakes; presented in **Table**
200 **3** according to genotype group.

201 An independent effect of genotype was observed for dietary **fat** and TC
202 concentrations at baseline (total **fat**, $P=0.002$; SFA, $P=0.002$; MUFA, $P=0.002$;
203 PUFA, $P=0.003$ and omega-3, $P=0.004$), with the highest TC concentrations seen in
204 carriers of $\epsilon 4$ allele (E4+). Overall diet effects (SFA, $P=0.008$; MUFA, $P=0.025$;
205 PUFA, $P=0.007$ and omega-3, $P<0.001$) were observed for omega-3 index, with
206 lower dietary SFA ($11.7\% \pm 0.1$) and higher PUFA ($6.80\% \pm 0.05$) and omega-3
207 ($0.89\% \pm 0.01$) **fat** intake associated with a higher omega-3 index. Although a
208 significant MUFA $\times$ *APOE* interaction was observed for omega-3 index ($P=0.025$),
209 no differences between genotype groups and **fat** intakes were observed following
210 post-hoc analyses.

211 **Dietary and genotype effects of intervention (irrespective of group allocation)**

212 The associations between change in dietary **fat** intake (total **fat**, SFA, MUFA,
213 PUFA and omega-3), *APOE* genotype and change in **fat** $\times$ *APOE* interactions on TC
214 and omega-3 index following intervention (m6-m0) are reported in **Table 4**. Dietary
215 intake was split into participants who reduced fat intake and those who increased fat
216 intake. **Mean reductions and increases in dietary fat intakes are presented according**
217 **to genotype group.**

218 There was a significant impact of genotype on change in TC concentrations
219 following dietary **advice** intervention (total **fat**, $P=0.016$; SFA, $P=0.025$; MUFA, $P=$
220 0.019 ; PUFA, $P=0.024$ and omega-3, $P=0.027$). There were no independent effects
221 of diet on lipid biomarkers following dietary **advice** intervention, although trends were
222 observed for change in PUFA ($P=0.068$) and omega-3 **fat** intakes ($P=0.087$) on

omega-3 index. A trend was also observed for an omega-3 **fat** intake $\times$ *APOE* interaction on omega-3 index ($P=0.087$).

Effect of knowledge of *APOE* gene risk on dietary change compared with other levels of personalization

The allocation of *APOE* risk according to intervention level is shown in **Figure 1**. Participants (levels 1-3) advised to lower dietary SFA at baseline were selected for subsequent analysis. The effects of knowledge of *APOE* risk (E4+) in participants advised to reduce SFA intake at baseline on changes in diet, TC and omega-3 index (m6-m0) compared with other levels of personalization are reported in **Table 5 A** significantly greater reduction in total **fat** and SFA (%TE) was observed in E4+ participants receiving gene-based PN (Level 3) compared to those in the control group ($P=0.034$ and $P=0.035$ respectively). However, there were no differences in change in diet or biomarkers between personalized intervention groups.

The effects of knowledge of *APOE* non-risk (E4-) in participants advised to reduce SFA intake at baseline on changes in diet, TC and omega-3 index (m6-m0) compared with other levels of personalization are reported in **Table 6**. As previously, participants receiving gene-based PN had a significantly greater reduction in dietary SFA (%TE) compared with those in the control group ($P=0.029$). For total **fat** (%FE), a slight increase in intake was observed for the control group (Level 0) compared with a reduction in Level 3 (difference 2.72% TE, $P=0.006$). The opposite was observed for total carbohydrate, which reduced in the control group (Level 0) and increased in Level 3 (difference 2.15 %TE, $P=0.027$).

When comparing levels of personalization, a 0.88% greater reduction in SFA (%TE) was observed in E4- participants receiving non-gene-based PN (Level 2; PN based on diet and phenotype) compared with those E4- participants receiving gene-

based PN ($P = 0.025$). There were no significant differences between change in total fat, PUFA, MUFA, omega-3, carbohydrate and protein intake, or TC and omega-3 index for E4- carriers according to whether they received gene-based or non-gene-based PN (L3 vs. L1-2).

Effect of knowledge of *APOE* genotype on dietary change following gene-based personalized advice PN

The effect of knowledge of *APOE* risk (risk: E4+, E3/E4 and E4/E4 and non-risk: E4-, E2/E2, E2/E3 and E3/E3) in participants advised to reduce SFA intake at baseline on changes in diet, TC and omega-3 index (m6-m0) following gene-based PN (L3) are reported in **Table 7**. Approximately 30% of E4- participants receiving gene-based PN were advised to lower their SFA intake at baseline, compared with 53% of E4+ carriers (**Figure 1**). Following intervention, there were no significant differences in dietary response or change in biomarker between E4+ and E4- participants.

DISCUSSION

Key findings in the present analysis were higher TC concentrations in E4 carriers (E4+) and a nutrient intake-gene interaction between *APOE* genotype and MUFA intake for omega-3 index at baseline. Following intervention, gene-based PN resulted in significantly greater reductions in total fat and SFA (%TE) compared with standard dietary advice (control), irrespective of gene risk. For E4- ('non-risk') participants advised to lower SFA intake, gene-based PN resulted in smaller changes in dietary SFA intake at month 6 than non-gene-based PN (Level 2).

Although the *APOE* rs429358 distribution was not in Hardy-Weinberg equilibrium, the haplotype frequencies observed in the Food4Me cohort (ϵ 2, 6.5; ϵ 3, 79.3; ϵ 4, 14.2) were similar to those reported in previous studies of European populations (28). In contrast to previous observations (29, 30), there was no clear geographical cline in ϵ 4 frequency.

DBS TC differed according to *APOE* genotype with significantly higher TC observed in E4+ participants compared with E4-. The difference in TC between E4+ and those who were E4-: E3/E3 in the present study (0.15 mmol/L) was similar to previous data (0.16-0.36 mmol/L) in a large meta-analysis of 54,377 participants (31).

At baseline, there was a significant nutrient intake-gene interaction between total MUFA intake and *APOE* on long-chain omega-3 index, a reliable biomarker of omega-3 status, and dietary omega-3 PUFA, EPA and DHA intake (32, 33). Furthermore, there is a dose-dependent inverse association between omega-3 index and CHD mortality (33), with an index $\geq 8\%$ offering the most cardio-protective effects and an index $\leq 4\%$ being associated with the greatest risk of CHD mortality (27). Thus, the omega-3 index may be a risk factor for CHD (34). In the Food4Me study, a higher omega-3 index was associated with lower SFA and higher PUFA and dietary omega-3 intake. In a study investigating the determinants of omega-3 index in a Mediterranean population, there were significant associations between EPA and DHA intakes and omega-3 index ($P < 0.001$) and a trend for an inverse association between dietary SFA and omega-3 index ($P = 0.095$) (35).

It has been suggested that gene-based dietary information is more understandable and useful than general dietary guidelines (36) and may enhance motivation to change (37). In a 2010 systematic review, a beneficial effect of genome-based risk estimates on dietary behavior was reported (pooled OR for 2

RCT 2.24, 95% CI 1.17 to 4.27, $P = 0.01$, $I^2 = 0\%$); but no benefit of genome-based risk estimates on intention to change dietary behavior was observed (5). Furthermore, in a Canadian RCT, knowledge of *ACE* gene risk resulted in a significantly greater reduction in sodium intake compared with non-gene based advice (-287 ± 114 vs. 130 ± 118 mg/day, $P = 0.008$) at 12-month follow-up (38). Change in sodium intake by participants carrying the 'non-risk' *ACE* genotype (-244 mg/day) was not significantly different ($P = 0.11$) compared with the control group. In our present study, gene-based PN promoted significantly greater reductions in the intake of total fat and SFA than standard dietary advice (control), for both risk (E4+) and non-risk (E4-) participants advised to lower SFA. However, there were no significant differences in change of diet, TC or omega-3 index between *APOE* risk groups (E4+ and E4-) receiving gene-based PN. In the REVEAL study, which investigated the impact of knowledge of Alzheimer's disease (AD) risk (estimated using *APOE* genotype and family history to generate a numerical risk) on dietary behaviors, E4+ participants were significantly more likely to endorse AD-specific health behavior change than E4- participants at 12 months follow-up (39). A similar result was observed in a study investigating the impact of knowledge of *FTO* genotype on readiness to control weight; whereby individuals with higher 'risk' (AA or AT) displayed greater willingness to change than those with lower risk (TT) ($P = 0.051$) (40).

Whilst there was no additional benefit of gene-based PN for E4+ participants in the Food4Me study, knowledge of 'non-risk' (E4-) resulted in a lower reduction in SFA intake at 6 months compared with E4- participants receiving non-gene-based PN (Level 2) who were not informed of their *APOE* risk (-1.68% vs. -2.56%). Providing 'no-risk' genotypic results may reduce motivation to follow dietary advice

(41). A potential reason for the lack of response in Food4Me E4 carriers is the absence of a specific behavior change technique (BCT) involving information on the consequences of a specific behavior related to genotype. A key BCT in the CALO-RE taxonomy (a 40-item taxonomy to improve PA and healthy eating behaviors) is to “provide information of the consequences of the behavior to the individual”. In the context of *APOE* genotype, a consequence of carrying the $\epsilon 4$ allele would be increased CVD risk (31) and the corresponding risk-reducing behavior would be lowering SFA intake. In the present study, *APOE* risk information conveyed to participants was framed positively viz : “you have a genetic variation that can benefit by keeping a healthy intake of saturated fat and a normal level of blood cholesterol.” The lack of an explicit link to an adverse consequence of E4+ status, e.g. higher CVD risk, may have reduced the efficacy of this advice. In the REVEAL study, participants were informed that the E4 allele was associated with an increased risk of Alzheimer’s disease prior to gene disclosure (39). Whilst genotypic testing for polygenic disease risk may result in a fatalistic attitude (37), information on consequences of personal characteristics (e.g. genotype) and fear arousal can be useful aids in enhancing behavior change (42). In a meta-analysis of fear arousal techniques, stronger fear messages promoted greater intention and behavior change in public health campaigns, provided that the threat was perceived to be severe, personally relevant, and that the individual could take specific action to mitigate their risk (43). In a Finnish RCT, knowledge of personal *APOE* risk resulted in greater short-term improvements in dietary quality, WC and serum triacylglycerol, when participants were informed of the link between dietary fat, cholesterol and CVD risk in an oral communication session (44). Furthermore, E4+ individuals significantly

improved fat quality at 6-months ($P < 0.01$), whereas there was no difference in fat quality in the E4- or control groups (44).

A limitation of internet-delivered PN (as used in our Food4Me study) is the reduced opportunity to employ BCT in response to verbal and non-verbal cues (e.g. body-language, facial expressions). Recent focus group data also revealed a lack of understanding amongst consumers of the use of genetic information to tailor dietary advice, and opinions regarding gene-based PN were mostly negative (45). **Given that understanding and 'knowledge' of specific gene-based PN advice was not evaluated in the Food4Me study, it is not possible to ascertain if this contributed to the lack of effect observed.** The Food4Me study was designed to assess the impact of three levels of personalization on dietary change and was not specifically targeted to the *APOE* genotype. Furthermore, although participants were informed that they had a 'risky' gene variant that would benefit from dietary change, advice was not stratified according to specific genotype groups (e.g. differing advice for E2/E3 and E3/E3). Strengths of this study include using the internet to assess and deliver dietary advice, prospective genotyping, a larger sample size than reported previously (39, 44, 46), the measurement of actual dietary change, as distinct from intention to change, and the availability of relevant blood-based biomarkers of **fat status** (obtained from unsupervised sampling). As such, the Food4Me study provides robust evidence of the impact of knowledge of *APOE* risk on adherence to dietary advice.

CONCLUSION

APOE status was significantly associated with TC at baseline with highest concentrations in E4+ participants. Whilst gene-based PN targeted to *APOE* was more effective in reducing SFA intake than standard dietary advice, there was no

added benefit of knowledge of *APOE* 'risk' on dietary change. Furthermore, it appears that disclosure of genotypic 'non-risk' status may have weakened the dietary response to PN. Future research should explore ways in which this detrimental response to gene-based PN can be mitigated.

ACKNOWLEDGEMENTS

The authors' responsibilities were as follows: MG, JCM, JAL, ERG, LB, WHMS, HD, CAD, JAM, YM and IT, contributed to the research design. RF, ALM, CCM, CFMM, HF, SNC, SK, LT, AS, MCW and JCM conducted the intervention. RF and CCM performed the statistical analysis for the manuscript. RF and JAL drafted the paper. All authors contributed to a critical review of the manuscript during the writing process. All authors approved the final version to be published. CAD is a founder, board member, stockowner and a consultant for Vitas Ltd. No other authors declare a conflict of interest. This work was funded by the EU FP7 Food4Me (KBBE.2010.2.3-02, Project no. 265494).

REFERENCES

1. Nichols M, Townsend N, Scarborough P, Rayner M. Cardiovascular disease in Europe 2014: epidemiological update. *Eur Heart J* 2014;35(42):2950-9.
2. Alwan A, Armstrong T, Bettcher D, Branca F, Chisholm D, Ezzati M, Garfield R, MacLean D, Mathers C, Mendis S, et al. Global status report on noncommunicable diseases 2010. Geneva: World Health Organization, 2011.
3. Kreuter MW, Wray RJ. Tailored and targeted health communication: strategies for enhancing information relevance. *Am J Health Behav* 2003;27(Supplement 3):S227-S32.

4. Celis-Morales C, Lara J, Mathers JC. Personalising nutritional guidance for more effective behaviour change. *Proc Nutr Soc* 2014;1-9.
5. Marteau TM, French DP, Griffin SJ, Prevost AT, Sutton S, Watkinson C, Attwood S, Hollands GJ. Effects of communicating DNA - based disease risk estimates on risk - reducing behaviours. *The Cochrane Library* 2010. doi: 10.1002/14651858.CD007275.pub2.
6. San-Cristobal R, Milagro FI, Martínez JA. Future challenges and present ethical considerations in the use of personalized nutrition based on genetic advice. *Journal of the Academy of Nutrition and Dietetics* 2013;113(11):1447-54.
7. Davignon J, Gregg RE, Sing CF. Apolipoprotein E polymorphism and atherosclerosis. *Arterioscler Thromb Vasc Biol* 1988;8(1):1-21. doi: 10.1161/01.ATV.8.1.1
8. Song Y, Stampfer MJ, Liu S. Meta-analysis: apolipoprotein E genotypes and risk for coronary heart disease. *Ann Intern Med* 2004;141(2):137-47. doi: 10.7326/0003-4819-141-2-200407200-00013
9. Bennet AM, Di Angelantonio E, Ye Z, Wensley F, Dahlin A, Ahlbom A, Keavney B, Collins R, Wiman B, de Faire U, et al. Association of apolipoprotein E genotypes with lipid levels and coronary risk. *JAMA* 2007;298(11):1300-11. doi: 10.1001/jama.298.11.1300
10. Waterworth DM, Ricketts SL, Song K, Chen L, Zhao JH, Ripatti S, Aulchenko YS, Zhang W, Yuan X, Lim N, et al. Genetic variants influencing circulating lipid levels and risk of coronary artery disease. *Arterioscler Thromb Vasc Biol* 2010;30(11):2264-76. doi: 10.1161/ATVBAHA.109.201020

11. Wilson PW, Schaefer EJ, Larson MG, Ordovas JM. Apolipoprotein E alleles and risk of coronary disease. A meta-analysis. *Arterioscler Thromb Vasc Biol* 1996;16(10):1250-5. doi: 10.1161/01.ATV.16.10.1250
12. Povel CM, Boer JM, Imholz S, Dollé ME, Feskens EJ. Genetic variants in lipid metabolism are independently associated with multiple features of the metabolic syndrome. *Lipids Health Dis* 2011;10:118. doi: 10.1186/1476-511X-10-118
13. Ward H, Mitrou PN, Bowman R, Luben R, Wareham NJ, Khaw K-T, Bingham S. APOE genotype, lipids, and coronary heart disease risk: a prospective population study. *Arch Intern Med* 2009;169(15):1424-9. doi: 10.1001/archinternmed.2009.234
14. Masson LF, McNeill G, Avenell A. Genetic variation and the lipid response to dietary intervention: a systematic review. *Am J Clin Nutr* 2003;77(5):1098-111.
15. Carvalho-Wells AL, Jackson KG, Lockyer S, Lovegrove JA, Minihane AM. APOE genotype influences triglyceride and C-reactive protein responses to altered dietary fat intake in UK adults. *Am J Clin Nutr* 2012;96(6):1447-53. doi: 10.3945/ajcn.112.043240
16. Ordovas JM, Lopez-Miranda J, Mata P, Perez-Jimenez F, Lichtenstein AH, Schaefer EJ. Gene-diet interaction in determining plasma lipid response to dietary intervention. *Atherosclerosis* 1995;118:S11-S27.
17. Lovegrove JA, Gitau R. Personalized nutrition for the prevention of cardiovascular disease: a future perspective. *Journal of Human Nutrition and Dietetics* 2008;21:306-16.
18. Food4Me. Internet: <http://www.food4me.org> (accessed March 2016).

19. Celis-Morales C, Livingstone KM, Marsaux CF, Forster H, O'Donovan CB, Woolhead C, Macready AL, Fallaize R, Navas-Carretero S, San-Cristobal R. Design and baseline characteristics of the Food4Me study: a web-based randomised controlled trial of personalised nutrition in seven European countries. *Genes Nutr* 2015;10(1):1-13. doi: 10.1007/s12263-014-0450-2
20. Forster H, Fallaize R, Gallagher C, O'Donovan CB, Woolhead C, Walsh MC, Macready AL, Lovegrove JA, Mathers JC, Gibney MJ. Online Dietary Intake Estimation: The Food4Me Food Frequency Questionnaire. *J Med Internet Res* 2014;16(6). doi: 10.2196/jmir.3105
21. Fallaize R, Forster H, Macready AL, Walsh MC, Mathers JC, Brennan L, Gibney ER, Gibney MJ, Lovegrove JA. Online dietary intake estimation: reproducibility and validity of the Food4Me Food Frequency Questionnaire against a 4-day weighed food record. *J Med Internet Res* 2014;16(8). doi: 10.2196/jmir.3355
22. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *The American journal of clinical nutrition* 1982;36(5):936-42.
23. Montoye HJ, Kemper HC, Saris WH, Washburn RA. Measuring physical activity and energy expenditure: Human Kinetics Champaign, IL, 1996.
24. Philippaerts RM, Westerterp KR, Lefevre J. Doubly labelled water validation of three physical activity questionnaires. *Int J Sports Med* 1999;20(5):284-9.
25. Bonomi AG, Plasqui G, Goris AH, Westerterp KR. Estimation of Free - Living Energy Expenditure Using a Novel Activity Monitor Designed to Minimize Obtrusiveness. *Obesity* 2010;18(9):1845-51. doi: 10.1038/oby.2010.34

26. Forster H, Walsh MC, O'Donovan CB, Woolhead C, McGirr C, O'Riordan R, Celis-Morales C, Fallaize R, Macready AL, Marsaux CF, et al. A Dietary Feedback System for the Delivery of Consistent Personalized Dietary Advice in the Web-Based Multicenter Food4Me Study. *J Med Internet Res* 2016;18(6):e150. doi: 10.2196/jmir.5620
27. Harris WS, von Schacky C. The Omega-3 Index: a new risk factor for death from coronary heart disease? *Prev Med* 2004;39(1):212-20. doi: 10.1016/j.ypmed.2004.02.030
28. Schiele F, De Bacquer D, Vincent-Viry M, Beisiegel U, Ehnholm C, Evans A, Kafatos A, Martins MC, Sans S, Sass C. Apolipoprotein E serum concentration and polymorphism in six European countries: the ApoEurope Project. *Atherosclerosis* 2000;152(2):475-88. doi: 10.1016/S0021-9150(99)00501-8
29. Eichner JE, Dunn ST, Perveen G, Thompson DM, Stewart KE, Stroehla BC. Apolipoprotein E polymorphism and cardiovascular disease: a HuGE review. *Am J Epidemiol* 2002;155(6):487-95. doi: 10.1093/aje/155.6.487
30. Tiret L, de Knijff P, Menzel H-J, Ehnholm C, Nicaud V, Havekes LM. ApoE polymorphism and predisposition to coronary heart disease in youths of different European populations. The EARS Study. European Atherosclerosis Research Study. *Arterioscler Thromb Vasc Biol* 1994;14(10):1617-24. doi: 10.1161/01.ATV.14.10.1617
31. Khan TA, Shah T, Prieto D, Zhang W, Price J, Fowkes GR, Cooper J, Talmud PJ, Humphries SE, Sundstrom J. Apolipoprotein E genotype, cardiovascular biomarkers and risk of stroke: Systematic review and meta-analysis of 14 015

- stroke cases and pooled analysis of primary biomarker data from up to 60 883 individuals. *Int J Epidemiol* 2013;42(2):475-92. doi: 10.1093/ije/dyt034
32. Andersen LF, Solvoll K, Dreven CA. Very-long-chain n-3 fatty acids as biomarkers for intake of fish and n-3 fatty acid concentrates. *The American journal of clinical nutrition* 1996;64(3):305-11.
 33. Harris WS. The omega-3 index as a risk factor for coronary heart disease. *Am J Clin Nutr* 2008;87(6):1997S-2002S.
 34. von Schacky C. Omega-3 index and cardiovascular health. *Nutrients* 2014;6(2):799-814. doi: 10.3390/nu6020799
 35. Sala-Vila A, Harris WS, Cofán M, Pérez-Heras AM, Pintó X, Lamuela-Raventós RM, Covas M-I, Estruch R, Ros E. Determinants of the omega-3 index in a Mediterranean population at increased risk for CHD. *Br J Nutr* 2011;106(03):425-31. doi: 10.1017/S0007114511000171
 36. Nielsen DE, El-Sohemy A. A randomized trial of genetic information for personalized nutrition. *Genes Nutr* 2012;1-8. doi: 10.1007/s12263-012-0290-x
 37. Joost HG, Gibney MJ, Cashman KD, Görman U, Hesketh JE, Mueller M, van Ommen B, Williams CM, Mathers JC. Personalised nutrition: status and perspectives. *Br J Nutr* 2007;98(1):26-31. doi: 10.1017/S0007114507685195
 38. Nielsen DE, El-Sohemy A. Disclosure of genetic information and change in dietary intake: a randomized controlled trial. *PloS one* 2014;9(11):e112665. doi: 10.1371/journal.pone.0112665
 39. Chao S, Roberts JS, Marteau TM, Silliman R, Cupples LA, Green RC. Health behavior changes after genetic risk assessment for Alzheimer disease: The REVEAL Study. *Alzheimer Dis Assoc Disord* 2008;22(1):94. doi: 10.1097/WAD.0b013e31815a9dcc

40. Meisel SF, Wardle J. Responses to FTO genetic test feedback for obesity in a sample of overweight adults: a qualitative analysis. *Genes Nutr* 2014;9(1):1-4.
41. Hunter DJ, Khoury MJ, Drazen JM. Letting the Genome out of the Bottle — Will We Get Our Wish. *N Engl J Med* 2008;358:105-7. doi: 10.1056/NEJMp0708162
42. Wilson BJ. Designing media messages about health and nutrition: what strategies are most effective? *J Nutr Educ Behav* 2007;39(2):S13-S9. doi: 10.1016/j.jneb.2006.09.001
43. Witte K, Allen M. A meta-analysis of fear appeals: Implications for effective public health campaigns. *Health Educ Behav* 2000;27(5):591-615. doi: 10.1177/109019810002700506
44. Hietaranta-Luoma HL, Tahvonen R, Iso-Touru T, Puolijoki H, Hopia A. An Intervention Study of Individual, apoE Genotype-Based Dietary and Physical-Activity Advice: Impact on Health Behavior. *J Nutrigenet Nutrigenomics* 2014;7(3):161-74. doi: 10.1159/000371743
45. Berezowska A, Fischer ARH, Ronteltap A, Kuznesof S, Macready AL, Fallaize R, van Trijp HCM. Understanding consumer evaluations of personalised nutrition services in terms of the privacy calculus: a qualitative study. *Public Health Genomics* 2014;17:127-40. doi: 10.1159/000358851
46. Hietaranta-Luoma HL, Åkerman K, Tahvonen R, Puolijoki H, Hopia A. Using Individual, ApoE Genotype-Based Dietary and Physical Activity Advice to Promote Healthy Lifestyles in Finland—Impacts on Cardiovascular Risk Markers. *Open Journal of Preventive Medicine* 2015;5(05):206. doi: 10.4236/ojpm.2015.55024

TABLE 1. Frequency of *APOE* genotype and *APOE* allele by Food4Me center (n=1466)

	All	Ireland	UK	The Netherlands	Germany	Poland	Spain	Greece
Genotype (n, %)								
E2/E2	6 (0.4)	1 (0.5)	0 (0.0)	3 (1.4)	0 (0.0)	2 (1.0)	0 (0.0)	0 (0.0)
E2/E3	152 (10.4)	14 (6.5)	22 (10.6)	28 (12.7)	21 (10.2)	29 (14.4)	22 (10.4)	16 (7.7)
E2/E4	27 (1.8)	3 (1.4)	6 (2.9)	3 (1.4)	7 (3.4)	4 (2.0)	1 (0.5)	3 (1.4)
E3/E3	922 (62.9)	133 (62.1)	132 (64.1)	124 (56.4)	125 (61.0)	125 (62.1)	139 (65.6)	144 (69.2)
E3/E4	330 (22.5)	57 (26.6)	43 (20.8)	58 (26.4)	48 (23.4)	38 (18.9)	46 (21.7)	40 (19.2)
E4/E4	29 (2.0)	6 (2.8)	3 (1.5)	4 (1.8)	4 (2.0)	3 (1.5)	4 (1.9)	5 (2.4)
Total	1466 (100)	214 (100)	206 (100)	220 (100)	205 (100)	201 (100)	212 (100)	208 (100)
E2 carriers ¹	158 (10.8)	15 (7.0)	22 (10.7)	31 (14.1)	21 (10.2)	31 (15.4)	22 (10.4)	16 (7.7)
E4 carriers ¹	359 (24.5)	63 (29.4)	46 (22.3)	62 (28.2)	52 (25.4)	41 (20.4)	50 (23.6)	45 (21.6)
Allele frequency (%)								
ε2	6.5	4.4	6.5	8.4	6.8	8.9	5.4	4.6
ε3	79.3	78.7	76.2	75.9	77.8	76.0	81.6	82.7
ε4	14.2	16.8	17.4	15.7	15.3	15.1	13.0	12.7

¹Genotype groups combined; E2 carriers represent E2/E2 and E2/E3, E4 carriers represent E4/E3 and E4/E4

TABLE 2. Anthropometric characteristics and fasted blood biomarkers by *APOE* genotype in European adults in the Food4Me study¹

	APOE genotype ¹				P ²
	E4-			E4+	
	All (n=1439)	E2 carriers (n=158)	E3/E3 (n=922)	E4 carriers (n=359)	
Gender ratio (M/F)	611/846				
Age (y)	40 ± 0.4	40 ± 1	40 ± 0.4	40 ± 0.7	0.630
BMI (kg/m ²)	25.5 ± 0.13	25.7 ± 0.4	25.4 ± 0.2	25.5 ± 0.3	0.704
Weight (kg)	74.6 ± 0.44	76.8 ± 1.4	74.3 ± 0.5	75.4 ± 0.8	0.608
Waist circumference (m)	0.86 ± 0.004	0.87 ± 0.01	0.86 ± 0.005	0.85 ± 0.01	0.693
Height (m)	1.71 ± 0.003	1.73 ± 0.01	1.71 ± 0.003	1.72 ± 0.005	0.252
Cholesterol (mmol/L)	4.59 ± 0.03	4.42 ± 0.08 ^a	4.55 ± 0.03 ^a	4.70 ± 0.05 ^b	0.002
Omega 3 index	5.68 ± 0.03	5.81 ± 0.10	5.66 ± 0.04	5.74 ± 0.06	0.341

¹ Data are means ± SEM² Data were analyzed by GLM with adjustment for age, gender, center and BMI. Where *P* for genotype < 0.05, a Bonferroni post-hoc test was applied to determine between-group effects. Superscript letters ^a and ^b denote significant differences between genotype groups, *P* < 0.05.

TABLE 3. Effect of *APOE* genotype and dietary fat intake (total and fat classes)¹ on metabolic markers measured in dried blood spots at baseline in the Food4Me intervention study²

	E4-				E4+		Diet	<i>P</i> ³	Diet × Genotype
	E2 carriers (n=158)		E3/E3 (n=922)		E4 carriers (n=359)				
	Low Intake	High Intake	Low Intake	High Intake	Low Intake	High Intake			
Total fat	(n=80)	(n=78)	(n=452)	(n=470)	(n=188)	(n=171)			
Total fat (%TE)	31.7 ± 0.4	39.9 ± 0.4	31.3 ± 0.2	40.6 ± 0.2	31.3 ± 0.3	40.6 ± 0.3			
Cholesterol (mmol/L)	4.37 ± 0.11	4.48 ± 0.11	4.45 ± 0.04	4.64 ± 0.04	4.66 ± 0.07	4.73 ± 0.07	0.251	0.002	0.435
Omega-3 index	5.81 ± 0.10	5.81 ± 0.13	5.66 ± 0.06	5.64 ± 0.06	5.79 ± 0.09	5.68 ± 0.09	0.989	0.344	0.456
SFA	(n=77)	(n=81)	(n=456)	(n=466)	(n=187)	(n=172)			
SFA (%TE)	11.7 ± 0.2	16.7 ± 0.2	11.7 ± 0.1	16.7 ± 0.1	11.6 ± 0.1	16.4 ± 0.1			
Cholesterol (mmol/L)	4.40 ± 0.11	4.44 ± 0.11	4.49 ± 0.04	4.61 ± 0.04	4.66 ± 0.07	4.73 ± 0.07	0.413	0.002	0.789
Omega-3 index	5.86 ± 0.14	5.76 ± 0.13	5.72 ± 0.06	5.58 ± 0.06	5.88 ± 0.09	5.57 ± 0.09	0.008	0.343	0.573
MUFA	(n=84)	(n=74)	(n=451)	(n=471)	(n=185)	(n=174)			
MUFA (%TE)	11.7 ± 0.2	15.5 ± 0.2	11.4 ± 0.1	16.1 ± 0.1	11.5 ± 0.1	16.1 ± 0.2			
Cholesterol (mmol/L)	4.40 ± 0.10	4.45 ± 0.11	4.49 ± 0.04	4.60 ± 0.04	4.98 ± 0.07	4.80 ± 0.07	0.078	0.002	0.470
Omega-3 index	5.67 ± 0.13	5.97 ± 0.14	5.71 ± 0.06	5.60 ± 0.06	5.86 ± 0.09	5.60 ± 0.09	0.025	0.280	0.025
PUFA	(n=86)	(n=72)	(n=460)	(n=462)	(n=174)	(n=185)			
PUFA (%TE)	4.7 ± 0.1	6.8 ± 0.1	4.6 ± 0.1	6.8 ± 0.1	4.7 ± 0.1	6.7 ± 0.1			
Cholesterol (mmol/L)	4.38 ± 0.10	4.47 ± 0.11	4.51 ± 0.04	4.59 ± 0.04	4.69 ± 0.07	4.69 ± 0.07	0.445	0.003	0.614
Omega-3 index	5.65 ± 0.13	6.00 ± 0.14	5.52 ± 0.06	5.77 ± 0.06	5.62 ± 0.09	5.84 ± 0.09	0.007	0.291	0.803
Omega-3	(n=80)	(n=78)	(n=485)	(n=437)	(n=155)	(n=204)			
Omega-3 (%TE)	0.55 ± 0.01	0.90 ± 0.03	0.55 ± 0.01	0.89 ± 0.01	0.55 ± 0.01	0.89 ± 0.02			
Cholesterol (mmol/L)	4.43 ± 0.11	4.41 ± 0.11	4.50 ± 0.04	4.61 ± 0.05	4.64 ± 0.08	4.74 ± 0.07	0.068	0.004	0.820
Omega-3 index	5.50 ± 0.13	6.12 ± 0.08	5.34 ± 0.05	5.99 ± 0.06	5.30 ± 0.09	6.07 ± 0.08	<0.001	0.546	0.463

¹ Intakes of fat were dichotomised at the median: total fat, 35.8% (low intake, 31.4% $\pm$ 0.1; high intake 40.5% $\pm$ 0.1); SFA, 14.0% (low intake, 11.7% $\pm$ 0.1; high intake 16.6% $\pm$ 0.1); MUFA, 13.5% (low intake, 11.5% $\pm$ 0.1; high intake 16.0% $\pm$ 0.1); PUFA, 5.6% (low intake, 4.67% $\pm$ 0.02; high intake 6.80% $\pm$ 0.05); omega-3, 0.67% (low intake, 0.55% $\pm$ 0.01; high intake 0.89% $\pm$ 0.01)

² Genotype groups combined; E2 carriers represent E2/E2 and E2/E3, E4 carriers represent E4/3 and E4/E4; %TE, % total energy; low intake, less than median fat intake; high intake, greater than median fat intake; data are mean $\pm$ SEM

³ Data were analysed by GLM with adjustment for centre, gender, age and BMI. Where *P* for diet x genotype < 0.05, a Bonferroni post-hoc test was applied to determine between-group effects (significant differences were not detected post-hoc)

TABLE 4. Effect of *APOE* genotype and change in dietary fat intake (total and **fat** classes)¹ on changes in metabolic markers measured in dried blood spots between baseline and month 6 for participants in the Food4Me intervention study²

	E4-				E4+		Diet	<i>P</i> ³	
	E2 carriers (n=132)		E3/E3 (n=794)		E4 carriers (n=315)				
	Decreased Intake	Increased Intake	Decreased Intake	Increased Intake	Decreased Intake	Increased Intake			
Total fat	(n=72)	(n=60)	(n=424)	(n=370)	(n=178)	(n=137)			
Total fat (%TE)	-4.49 ± 0.42	3.90 ± 0.41	-4.91 ± 0.19	3.93 ± 0.18	-4.76 ± 0.29	4.16 ± 0.34			
Cholesterol (mmol/L)	-0.26 ± 0.12	-0.24 ± 0.13	-0.18 ± 0.05	-0.21 ± 0.05	-0.26 ± 0.08	-0.03 ± 0.09	0.527	0.016	0.313
Omega-3 index	0.24 ± 0.15	-0.08 ± 0.16	0.26 ± 0.06	0.25 ± 0.06	0.40 ± 0.09	0.15 ± 0.11	0.808	0.136	0.384
SFA	(n=86)	(n=46)	(n=484)	(n=310)	(n=206)	(n=109)			
SFA (%TE)	-2.56 ± 0.21	2.01 ± 0.23	-2.68 ± 0.10	1.75 ± 0.08	-2.48 ± 0.14	2.13 ± 0.19			
Cholesterol (mmol/L)	-0.32 ± 0.11	-0.14 ± 0.14	-0.21 ± 0.05	-0.17 ± 0.06	-0.18 ± 0.07	-0.11 ± 0.10	0.982	0.025	0.941
Omega-3 index	0.24 ± 0.14	-0.14 ± 0.17	0.33 ± 0.06	0.14 ± 0.07	0.39 ± 0.09	0.10 ± 0.12	0.986	0.069	0.377
MUFA	(n=64)	(n=68)	(n=397)	(n=397)	(n=165)	(n=150)			
MUFA (%TE)	-1.88 ± 0.18	1.65 ± 0.17	-2.10 ± 0.10	2.00 ± 0.10	-2.19 ± 0.15	2.13 ± 0.17			
Cholesterol (mmol/L)	-0.29 ± 0.13	-0.21 ± 0.12	-0.21 ± 0.05	-0.19 ± 0.05	-0.29 ± 0.08	-0.01 ± 0.08	0.392	0.019	0.583
Omega-3 index	0.25 ± 0.15	-0.04 ± 0.15	0.23 ± 0.06	0.28 ± 0.06	0.36 ± 0.10	0.21 ± 0.10	0.547	0.309	0.373
PUFA	(n=58)	(n=74)	(n=357)	(n=437)	(n=153)	(n=162)			
PUFA (%TE)	-0.83 ± 0.10	1.12 ± 0.11	-1.06 ± 0.06	1.13 ± 0.06	-0.93 ± 0.07	1.13 ± 0.09			
Cholesterol (mmol/L)	-0.28 ± 0.13	-0.23 ± 0.12	-0.12 ± 0.05	-0.26 ± 0.05	-0.23 ± 0.08	-0.09 ± 0.08	0.611	0.024	0.148
Omega-3 index	-0.004 ± 0.16	0.18 ± 0.14	0.18 ± 0.07	0.32 ± 0.06	0.41 ± 0.10	0.17 ± 0.10	0.068	0.467	0.303
Omega-3	(n=53)	(n=79)	(n=294)	(n=500)	(n=129)	(n=186)			

Omega-3 (%TE)	-0.12 ± 0.02	0.18 ± 0.02	-0.14 ± 0.01	0.22 ± 0.02	-0.13 ± 0.01	0.15 ± 0.03			
Cholesterol (mmol/L)	-0.15 ± 0.14	-0.32 ± 0.11	-0.23 ± 0.06	-0.18 ± 0.05	-0.18 ± 0.09	-0.14 ± 0.08	0.738	0.027	0.738
Omega-3 index	0.02 ± 0.17	0.14 ± 0.14	0.02 ± 0.07	0.39 ± 0.06	0.24 ± 0.11	0.32 ± 0.09	0.087	0.412	0.087

¹ 0% change in fat intake used as a reference to dichotomize participants i.e. comparison of reduction vs. increase in fat intake; total fat (decrease, -4.82% ± 0.15; increase 3.98% ± 0.15), SFA (decrease, -2.62% ± 0.08; increase 1.84% ± 0.08), MUFA (decrease, -2.10% ± 0.07; increase 1.99% ± 0.08), PUFA (decrease, -1.00% ± 0.04; increase 1.13% ± 0.04), omega-3 (decrease, -0.14% ± 0.01; increase 0.22% ± 0.02)

² Genotype groups combined; E2 carriers represent E2/E2 and E2/E3, E4 carriers represent E4/3 and E4/E4; %TE, % total energy; increased intake, greater than 0% change in fat intake; decreased intake, less than 0% change in fat intake; data are mean change ± SEM (m6 - m0)

³ Data were analysed by GLM with adjustment for baseline values, centre, gender, age and change in weight (m6 - m0).

TABLE 5. Effect of knowledge of *APOE* risk (E4+) on change in dietary intake between baseline and month 6 for participants in the Food4Me intervention study¹

	Control	Personalized intervention arms				<i>P</i> ²		
	Level 0 (L0) <i>APOE</i> risk (n=77)	Level 1 (L1) <i>APOE</i> risk (n=47)	Level 2 (L2) <i>APOE</i> risk (n=35)	Level 3 (L3) <i>APOE</i> risk (n=40)	L3 vs. Control (L0)	L3 vs. L1	L3 vs. L2	
Total fat (%TE)	0.37 ± 0.65	-3.03 ± 0.79	-1.63 ± 1.00	-3.07 ± 0.86	0.034	0.970	0.317	
SFA (%TE)	-0.72 ± 0.35	-2.53 ± 0.37	-1.58 ± 0.56	-1.95 ± 0.45	0.035	0.335	0.537	
MUFA (%TE)	0.37 ± 0.32	-0.71 ± 0.35	-0.41 ± 0.42	-1.05 ± 0.36	0.073	0.467	0.303	
PUFA (%TE)	-0.04 ± 0.13	0.20 ± 0.19	0.30 ± 0.23	0.01 ± 0.23	0.718	0.965	0.720	
Omega-3 (%TE)	0.04 ± 0.03	0.08 ± 0.03	0.08 ± 0.03	0.08 ± 0.03	0.899	0.900	0.990	
Carbohydrate (%TE)	-0.89 ± 0.76	1.89 ± 0.85	0.11 ± 0.98	1.55 ± 0.92	0.127	0.945	0.130	
Protein (%TE)	0.38 ± 0.43	0.40 ± 0.43	0.49 ± 0.49	1.37 ± 0.40	0.392	0.245	0.226	
BMI (kg/m ²)	-0.25 ± 0.13	-0.35 ± 0.15	-0.04 ± 0.19	-0.44 ± 0.18	0.231	0.590	0.086	
Cholesterol (mmol/L)	-0.32 ± 0.11	-0.04 ± 0.16	-0.39 ± 0.15	-0.19 ± 0.16	0.240	0.663	0.228	
Omega-3 index	-0.04 ± 0.11	0.29 ± 0.16	0.38 ± 0.16	0.14 ± 0.16	0.545	0.610	0.240	

¹ E4-, E2/E2, E2/E3 and E3/E3; E4+, E3/E4 and E4/E4; %TE, % total energy; data are mean change ± SEM (m6 - m0)

² Data were analysed by GLM with adjustment for baseline values, centre, gender, age and change in weight (m6 - m0).

TABLE 6. Effect of knowledge of *APOE* non-risk (E4-) on change in dietary intake between baseline and month 6 for participants in the Food4Me intervention study¹

	Control	Personalized intervention arms			<i>P</i> ²		
	Level 0 (L0) <i>APOE</i> non-risk (n=225)	Level 1 (L1) <i>APOE</i> non-risk (n=145)	Level 2 (L2) <i>APOE</i> non-risk (n=119)	Level 3 (L3) <i>APOE</i> non-risk (n=72)	L3 vs. Control (L0)	L3 vs. L1	L3 vs. L2
Total fat (%TE)	0.31 ± 0.37	-2.63 ± 0.47	-3.42 ± 0.51	-2.41 ± 0.66	0.006	0.280	0.381
SFA (%TE)	-0.31 ± 0.20	-1.88 ± 0.25	-2.56 ± 0.27	-1.68 ± 0.35	0.029	0.119	0.025
MUFA (%TE)	0.32 ± 0.17	-0.75 ± 0.22	-0.87 ± 0.24	-0.64 ± 0.31	0.012	0.382	0.601
PUFA (%TE)	0.25 ± 0.11	-0.01 ± 0.14	0.04 ± 0.15	-0.18 ± 0.19	0.053	0.273	0.119
Omega-3 (%TE)	0.13 ± 0.03	0.02 ± 0.04	0.05 ± 0.05	0.06 ± 0.06	0.278	0.442	0.903
Carbohydrate (%TE)	-1.22 ± 0.45	1.65 ± 0.55	1.92 ± 0.61	0.93 ± 0.79	0.027	0.211	0.558
Protein (%TE)	0.85 ± 0.21	0.77 ± 0.26	0.80 ± 0.28	1.17 ± 0.36	0.997	0.346	0.634
BMI (kg/m ²)	-0.28 ± 0.08	-0.44 ± 0.09	-0.41 ± 0.10	-0.51 ± 0.13	0.970	0.711	0.364
Cholesterol (mmol/L)	-0.27 ± 0.07	-0.22 ± 0.08	-0.39 ± 0.09	-0.41 ± 0.12	0.855	0.959	0.560
Omega-3 index	0.27 ± 0.07	0.11 ± 0.09	0.26 ± 0.09	0.18 ± 0.12	0.536	0.700	0.464

¹ E4-, E2/E2, E2/E3 and E3/E3; E4+, E3/E4 and E4/E4; %TE, % total energy; data are mean change ± SEM (m6 - m0)

² Data were analysed by GLM with adjustment for baseline values, centre, gender, age and change in weight (m6 - m0).

TABLE 7. Effect of knowledge of *APOE* genotype on change in dietary intake between baseline and month 6 for participants receiving gene-based personalized nutrition (Level 3) in the Food4Me intervention study¹

	Level 3 (L3)		P²
	<i>APOE</i> non-risk (E4-) (n=72)	<i>APOE</i> risk (E4+) (n=40)	
Total fat (%TE)	-2.41 ± 0.64	-3.07 ± 0.86	0.433
SFA (%TE)	-1.68 ± 0.33	-1.95 ± 0.45	0.348
MUFA (%TE)	-0.64 ± 0.28	-1.05 ± 0.36	0.307
PUFA (%TE)	-0.18 ± 0.17	0.01 ± 0.23	0.223
Omega-3 (%TE)	0.06 ± 0.02	0.08 ± 0.03	0.392
Carbohydrate (%TE)	0.93 ± 0.68	1.55 ± 0.92	0.421
Protein (%TE)	1.17 ± 0.30	1.37 ± 0.40	0.502
BMI (kg/m ²)	-0.51 ± 0.13	-0.44 ± 0.18	0.229
Cholesterol (mmol/L)	-0.41 ± 0.12	-0.19 ± 0.16	0.203
Omega-3 index	0.18 ± 0.12	0.14 ± 0.16	0.777

¹ E4-, E2/E2, E2/E3 and E3/E3; E4+, E3/E4 and E4/E4; %TE, % total energy; data are mean change ± SEM (m6 - m0)

² Data were analysed by GLM with adjustment for baseline values, centre, gender, age and change in weight (m6 - m0).

Figure 1: Consort diagram of participants randomized into the Food4Me Proof of Principle Study * Total number of participants reporting one or more exclusion criteria. **Parentheses** indicate the percentage of each group who received advice to reduce SFA intake at month 0.

