

Peptide hormones and lipopeptides: from self-assembly to therapeutic applications

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Hutchinson_et_al-2017-Journal_of_Peptide_Science.pdf

Hutchinson, J. A., Burholt, S. and Hamley, I. W. ORCID: <https://orcid.org/0000-0002-4549-0926> (2017) Peptide hormones and lipopeptides: from self-assembly to therapeutic applications. *Journal of Peptide Science*, 23 (2). pp. 82-94. ISSN 1075-2617 doi: 10.1002/psc.2954 Available at <https://reading-pure-test.eprints-hosting.org/69042/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/psc.2954>

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Peptide hormones and lipopeptides: from self-assembly to therapeutic applications

J. A. Hutchinson, S. Burholt and I. W. Hamley*

This review describes the properties and activities of lipopeptides and peptide hormones and how the lipidation of peptide hormones could potentially produce therapeutic agents combating some of the most prevalent diseases and conditions. The self-assembly of these types of molecules is outlined, and how this can impact on bioactivity. Peptide hormones specific to the uptake of food and produced in the gastrointestinal tract are discussed in detail. The advantages of lipidated peptide hormones over natural peptide hormones are summarised, in terms of stability and renal clearance, with potential application as therapeutic agents. © 2017 The Authors Journal of Peptide Science published by European Peptide Society and John Wiley & Sons Ltd.

Keywords: peptide hormones; lipopeptides; self-assembly; bioactivity; peptide; therapeutics

Peptide and Lipopeptide Self-assembly

The self-assembly of peptides in solution is driven by a combination of hydrogen bonding, electrostatic and other (e.g. π -stacking) interactions that also contribute to the self-assembly of lipopeptides [1]. However, in the latter class of system, hydrophobic interactions are important. The self-assembly of peptides and lipopeptides can be influenced by many variables including concentration, pH, ionic strength of solution or temperature [2].

Many self-assembling molecules are amphiphilic, meaning they have both hydrophobic and hydrophilic character. They generally self-assemble above a critical concentration, known as the critical aggregation concentration (CAC). Amphiphilic molecules such as lipids, peptides and proteins serve as building blocks for the construction of functional assemblies *in vivo*, e.g. the cytoskeleton and extracellular matrix. Lipids are one of the simplest amphiphilic structures and are composed of a hydrophilic polar head group and a hydrophobic tail. Peptides and proteins, however, are distinct in the way in which amphiphilicity is displayed because when folded, they can display regions that are either hydrophobic or hydrophilic. An example of this is an α -helix, as it could contain a section of hydrophobic residues along one face and a hydrophilic section of residues on the opposite face. For β -sheet structures, the peptide chain can be composed of alternating hydrophilic and hydrophobic residues, so that the side chains of the residues are displayed on opposite faces of the sheet.

In aqueous environments, amphiphilic molecules associate through non-covalent interactions to form ordered assemblies of different sizes, from nanometres to microns [3]. These self-assembled structures include spherical and worm-like micelles, vesicles, fibrils and nanotubes (Figure 1). Micelles consist of a hydrophobic inner core surrounded by a hydrophilic outer shell that is exposed to water, and they can be spheres, discs or worm-like structures [4]. Micelles form spontaneously when the concentration is above a critical micelle concentration [CMC] and temperature [5]. The CMC is a subcategory of CAC, which is a more general term for the aggregation into many different structures.

Amphiphiles with an intermediate level of hydrophobicity can assemble into bilayer vesicles. Vesicles are spherical, hollow,

lamellar structures with an aqueous core. The hydrophobic moieties form the inner section of the bilayer, and the hydrophilic parts are exposed to the aqueous environment [6].

Self-assembling peptides are peptides that undergo spontaneous assembly into ordered nanostructures. This is observed depending on the hydrophilic/lipophilic balance of the molecules, as well as the interactions between the peptide units [7]. In one case of self-assembling peptides, hydrogen bonding between backbones plays an important role by forcing the peptide monomers to pack longitudinally into β -sheets. The inter-sheet interactions between the side chains of the peptides regulate lateral packing, and the stronger these interactions, the better the lateral packing [8]. A further type of self-assembling peptide forms coiled coil structures, formed from aggregated α -helices [9].

Among all organic building blocks, peptides are very promising platforms because of their ease of synthesis, chemical diversity and their similar biological properties to proteins. In addition to this, peptides are very useful components in creating self-assembled nanostructures because of their biocompatibility, biodegradability and biofunctionality [10].

The number, type and sequence of amino acids determines the self-assembly of peptides, and depending on the amino acid sequence, the peptide can form a variety of different structures. As a result of this, peptides provide a unique platform for the design of nanomaterials with controllable structural features. Self-assembled peptide nanostructures have demonstrated potential use for many biomedical applications such as drug delivery, tissue engineering and antimicrobial agents, to name a few, and several of these applications are discussed in detail in the following.

* Correspondence to: I. W. Hamley, Department of Chemistry, University of Reading, Whiteknights, Reading RG6 6AD, UK. E-mail: i.w.hamley@reading.ac.uk

Department of Chemistry, University of Reading, Whiteknights, Reading RG6 6AD, UK


This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Biography

Jessica Hutchinson received her MChem degree from Northumbria University in 2014. She is currently a PhD student at the University of Reading under the guidance of Professor Ian W. Hamley. Her work is focused on the self-assembly and bioactivity of lipopeptides, in particular the lipidated gastrointestinal peptide hormone PYY_{3–36}, and fragments of it.

**Biography**

Samuel Burholt received his MChem with Forensic Science from the University of Reading in 2015. He is now entering his second year of a PhD with supervision from Professor Ian Hamley at the University of Reading. His current work is on the self-assembly of lipopeptides and incorporating them within gel networks. The main lipopeptide of interest is the lipidated version of the appetite-reducing peptide PYY_{3–36}.

**Biography**

Professor Ian W. Hamley is Diamond Professor of Physical Chemistry at the University of Reading. He has more than 20 years' experience of research on different types of soft materials, including peptides, polymers, liquid crystals and surfactants. He obtained his PhD from the University of Southampton in 1991 and then undertook postdoctoral research at AMOLF, Amsterdam, and the University of Minnesota. In 1993, he returned to a lectureship at the University of Durham, UK, and moved to the University of Leeds in 1995 where he was promoted to a professorship in 2004. He relocated to the University of Reading in 2005. He received a Royal Society-Wolfson Research Merit Award in 2011 and the RSC Peter Day Award for Materials Chemistry (2016). His research programme and peptide and peptide-conjugate-based materials focuses on their self-assembly behaviour and its relationship to bioactivity. He has published over 350 journal publications (h-index 58) and several edited and authored books.

**Peptide Amphiphiles**

Peptide amphiphiles (PAs) may comprise sequences of hydrophobic and hydrophilic peptides or hydrophilic peptides attached to lipid chains. The class of PA termed lipopeptide consists of one or more lipid chains attached to hydrophilic peptide sequences containing charged residues [2,7]. They are a class of molecules

that combine the structural features of amphiphilic surfactants with the functions of bioactive peptides, and they are known to assemble into a variety of nanostructures [11,12]. It is proposed that PAs designed to form bioactive fibrils should be composed of four key structural features (Figure 2): A hydrophobic domain that is typically an alkyl chain attached to a peptide sequence, which favours intermolecular hydrogen bonding. Then there is a charged amino acid domain that enhances solubility in water. The final structural feature that makes up a peptide amphiphile is a bio-derived or bio-inspired epitope that allows interaction with cells or proteins [13].

Molecules that contain both polar and non-polar elements often undergo self-assembly, which allows the hydrophilic moieties to be exposed to the aqueous environment and the hydrophobic moieties to be shielded from the aqueous media. There is normally a distinct relationship between the amphiphilic character of a peptide and its function [12]. For example, amphiphilic peptides fold into helices or sheets to allow the non-polar residues to interact with the lipid chains in the interior of the cell membrane and to allow the polar residues to be exposed to the aqueous environment. This self-assembly allows the peptide molecules to optimise their interaction with the surroundings.

Peptide amphiphiles have great potential in biomedical applications [1,11,13–18] and can be utilised to act as therapeutic agents to treat diseases by delivering drugs. They can then be metabolised into lipids and amino acids, which are then easily cleared by the kidneys [19]. In therapeutic applications, the hydrophobic tail assists transport across the cell membrane, and the peptide epitope can then be used to target a specific cell via a ligand-receptor complex [20].

Lipopeptides are able to form supramolecular nanostructures such as fibrils, micelles and vesicles. Lipopeptide biosurfactants (surfactants of biological origin) are produced by a wide variety of bacteria, fungi and yeast [7,21]. They are surface active compounds that have the ability to decrease the surface and interfacial tension, allowing them to disrupt biological activity as part of the organism's host defence mechanism [21].

Applications of Self-assembled PAs and Lipopeptides

Lipopeptides have a wide range of applications such as use as antimicrobial agents and in immune disease therapies, cosmeceuticals and also fungicides, all of which are explained in more detail in the succeeding sections.

Biosurfactants with Antimicrobial and Antifungal Applications

There are many types of lipopeptides; among the most popular are the classes of surfactins, iturins and fengycins, which are produced by the *Bacillus subtilis* family [21–23]. *B. subtilis* strains produce a wide range of lipopeptides that are potent biosurfactants and have specific antimicrobial and antiviral activities [21]. The fact that surfactins are biosurfactants means that they have diverse functional properties such as low toxicity, biodegradability and a higher tolerance towards variation of temperature and pH [21]. Iturins are pore-forming lipopeptides with antifungal activity, and this is dependent on the interaction with the cytoplasmic membrane of the target cells [21,22,24]. Finally, fengycins are another class of biosurfactant with antifungal properties [23,24].




Figure 1. Molecular interactions and possible self-assembled structures of typical lipopeptides.




Figure 2. The four domains in a PA molecule required for self-assembly into β -sheet fibrils with a coating of bioactive epitopes [11].

Toll-like Receptor Agonists

Lipopeptides, such as those incorporating the CSK₄ peptide motif and one, two or three palmitoyl (hexadecyl) chains [25], can also act as toll-like receptor (TLR) agonists, which have important applications in the treatment of disease. TLRs are transmembrane proteins that are very important in the immune system and as a result are therapeutic targets to treat disease.

These receptor agonists respond to invading pathogens by recognising specific pathogen-associated microbial patterns (PAMPs) or danger-associated molecular patterns (DAMPs), which are primarily produced by microbial pathogens [26]. PAMPs can contain a variety of different components including lipopolysaccharide, peptidoglycan, lipopeptide and bacterial DNA. DAMPs can be intracellular proteins or proteins from the extracellular matrix [27,28].

Toll-like receptors have a common structure being type-1 transmembrane proteins. This includes having an extracellular domain formed from leucine-rich repeats (LRRs) and a cytoplasmic tail containing an area known as the TLR domain. X-ray crystallography reveals that the LRR domain has a horseshoe-like shape [29], traversing the membrane. Currently, 10 TLRs have been found in humans, each with a different role and target [30].

Toll-like receptor agonists are compounds that stimulate these receptors to modulate the interaction with one of the PAMPs/DAMPs described earlier. The development of synthetic or natural agonists is an interesting avenue of research to treat a multitude of conditions that includes, but is not limited to; advanced melanoma [31], alcoholic chronic liver disease [32], asthma [33], neuropathic pain [34] and restenosis (re-narrowing of the blood vessels) [35].

Skincare

Peptide amphiphiles are employed in skincare products with the claimed ability to help stimulate collagen production. A commercially available lipopeptide with the trade name Matrixyl™ [C₁₆-Lys-Thr-Thr-Lys-Ser (KTTKS)] has been used in anti-wrinkle creams. This lipopeptide has been shown to self-assemble into a β -sheet tape-like superstructure (Figure 3). Small-angle X-ray scattering further established a bilayer structure with a spacing of 5.3 nm [36].

The mechanism for this stimulation of collagen expression is not fully understood, but it has been reported that the KTTKS polypeptide increases the skin's extracellular matrix (ECM) production




Figure 3. Confocal microscopy image of Matrixyl™ fibre superstructure. (Labelled with the dye rhodamine B, 0.0014 w% Matrixyl™ in water) [36].

[37,38]. ECM is the outer region of a cell that supports the cells and those around it. The conjugation of the KTTKS peptide motif to a palmitoyl (C_{16}) chain has been shown to enhance skin permeability, and the self-assembled structure (Figure 3) presents a peptide-rich surface on the nanopipes, these features contributing towards increased collagen production making it a highly sought after skincare additive. This lipopeptide's collagen-stimulating activity occurs even at low concentration, close to its [37]. Other lipopeptides, such as C_{16} -GHK or C_{16} -KT, also have been reported to have collagen-stimulating effects, and their self-assembly has been examined [39].

Tissue Scaffolds

Various groups have looked into the use of PAs and lipopeptides as scaffolds for the production of tissue or for other applications in regenerative medicine. In one study, the linear RGD amino acid sequence was conjugated with dialkyl chains. This scaffold allowed for spreading of melanoma and endothelial cells when it was carboxyl-coupled, but spreading was not seen with amino-coupled dialkyl chains [40]. Another group managed to encapsulate cells within nanofibrils made up of PAs with the Ile-Lys-Val-Ala-Val (IKVAV) peptide group (Figure 4). These PAs were shown to be effective at treating spinal cord injuries in mice. This was due to the nanofibrils inhibiting scar formation, which helped lessen the injury [41]. This is a fascinating and broad topic that has been




Figure 4. Self-assembly of IKVAV functionalized PAs into fibrils [42].

reviewed in much detail elsewhere by the Stupp group and is therefore outside the scope of the present review [42–47].

Antimicrobial Materials

There are a great number of studies on peptides with antimicrobial properties [48–50]. A common feature for antimicrobial activity is the presence of cationic residues such as lysine or more especially arginine as these can interact with cell walls. Lipidation of peptides has been shown to improve the uptake of the peptide into the cell [51]. This was investigated with a range of Gram-negative/positive bacteria and two fungal strains. Uptake into the cell wall and then subsequent disruption of the cell membrane was found out to be the mode of action, which caused bacterial leakage and cell death [51]. A peptide containing a trans-activating transcriptional activator (TAT) sequence along with an amphiphilic Gly–Arg sequence conjugated to cholesterol was also found to be an effective antimicrobial agent [52]. This peptide was shown to work by the same process of incorporation into the cell membrane followed by its disruption. The paper also showed that the PA can also be transported across the blood–brain barrier [52]. One of the most well-known lipopeptides used in treating infections is daptomycin (Figure 5) [53].

Daptomycin is able to combat systemic and life-threatening infections and is trademarked under the name Cubicin. It comprises a cyclic peptide group joined by an amide linkage to a lipid chain. The mechanism by which daptomycin acts on bacteria has been carefully examined [54]. It works by insertion of the lipid chain into the cell wall. Daptomycin molecules then aggregate, deforming the curvature of the membrane, causing holes to form and leading to the leakage of ions from the cell. This then causes a serious depolarization resulting in the inhibition of various synthesis processes including those of DNA, protein and RNA. This combines to cause cell death. The self-assembly of daptomycin was studied outside of the cell and it was shown to aggregate into micelles [53]. This aggregation process may be correlated to its excellent antimicrobial activity. A recent review on the self-assembly of lipopeptides including therapeutic lipopeptides is available [7].

Drug Delivery

Lipopeptides are interesting vehicles for drug delivery with several modes of activity. In one, the self-assembled structure of the PA is exploited, i.e. the hydrophobic area inside a micelle or bilayer structure is used to house a hydrophobic drug molecule. It is important to distinguish between PAs with inherent therapeutic potential and those that are merely vehicles or devices. In the former case, many types of bioactive peptides have been lipidated to create active pharmaceutical ingredients (APIs) (e.g. Table 1). In the latter




Figure 5. Daptomycin chemical structure, redrawn from [53].

Table 1. Peptide therapeutics on the market

Trade name	Peptide	Company	Molecular properties	Related reference
Copaxone	Glatiramer	Teva	Four amino acids (L-glutamic acid, L-alanine, L-lysine and L-tyrosine) in a defined molar ratio	[105,106]
Lupron	Leuprolide	Abbott	Synthetic nonapeptide analogue of naturally occurring gonadotropin-releasing hormone (GnRH or LH-RH)	[107,108]
Vicoza	Liraglutide	Novo	97% homologous to native human GLP-1 (7–37) by substituting arginine for lysine at position 34 and addition of a fatty acid chain	[109]
Zoladex	Goserelin	AZ	Natural LHRH/GnRH decapeptide with two substitutions to inhibit rapid degradation.	[110,111]
Sandostatin	Octreotide	Novartis	Longer acting synthetic octapeptide analogue of naturally occurring somatostatin	[112,113]
Forteo	Teriparatide	Lilly/Amylin	Recombinant form of parathyroid hormone consisting of the first (N-terminus) 34 amino acids, which is the bioactive portion of the hormone	[114,115]
Byetta	Exenatide	Lilly/Amylin	Synthetic version of exendin-4, a hormone found in the saliva of the Gila monster	[116,117]
Cubicin	Daptomycin	Cubist	Cyclic lipopeptide, consisting of 13 amino acids, 10 of which are arranged in a cyclic fashion and three on an exocyclic tail	[118,119]
Integrilin	Eptifibatide	Merck	Cyclic heptapeptide composed with S–S bridge, two unnatural building blocks and amide	[120,121]
Angiomax/angiox	Bivalirudin	Medicines	20-amino acid polypeptide	[122,123]
Fortical	Calcitonin	Upsher-Smith	32-amino acid polypeptide	[124–126]
Somatuline	Lanreotide	Ipsen	Cyclic peptide that is a long-acting analogue of somatostatin.	[127,128]

case, the self-assembled structure allows for the transport of the drug in highly aqueous environments to the target cells. An example of this is a TAT_{48–60} fragment conjugated with one, two or four attached octanoic acid groups [55]. These TAT PAs form β -sheet structures. It was shown that the PA with four octanoic acid groups could encapsulate and retain the hydrophobic drug paclitaxel. This was deemed to be due to the high hydrophobicity of the octanoic acid groups. This encapsulation was highly efficient ($6.8 \pm 0.4\%$), compared with previous nanoscale delivery vehicles for which encapsulation rarely exceeds 5%, making it a favoured drug delivery system [55]. Another strategy to create PAs for drug delivery is the lipidation of receptor-specific peptide head groups. The peptide head groups bind selectively to the receptor, while the lipid group allows the PA to cross cell membranes and also increases bioactivity through reducing metabolic degradation [56,57]. A good example of this is an amphiphilic lipopeptide comprising a palmitoyl (C₁₆) ester linked to Ala-Gly-Phe-Leu-Arg peptide motif, incorporating the anticancer drug Dalargin (Ala-Gly-Phe-Leu-Arg), which formed into nanofibres [58]. These fibres have high circulation times and also more importantly can cross the BBB. This allows the normally hydrophobic cancer drug to potentially be used to treat cancers in the brain, overcoming its low circulation and high hydrophobicity [58].

Biomaterial Templating

The various structures formed by PAs and lipopeptides have been exploited to template distinct inorganic materials (salts, metal oxides and metals), sometimes allowing for unique structures to be formed. PA nanostructures have the potential, for example, to be used to template the bone mineral hydroxyapatite, which is the body's primary storage depot for calcium and phosphorus in bones [59,60]. A further avenue of interest is

the incorporation of other metals, such as titanium, with mineralizing PAs to form hybrid bone implants. In one example, a titanium alloy foam was mixed with different PAs solutions and then further allowed to mineralise calcium phosphate. This was shown (Figure 6) to form new, immature bone adjacent and inside the hybrid, after 4 weeks [61]. This field of research is highly active and may, in the future, allow for treatment of bone diseases such as hydroxyapatite deposition disease or dystrophic calcification, among others.

Peptide Hormones

Peptide hormones are hormones made up of amino acid chains that primarily have an effect on the endocrine system. Based on the building units, hormones can be classified as either amino acid-based or steroid-based systems. The presence of amino acids in peptide hormones allows them to act on the surface of target cells via secondary messengers. This differs from steroid hormones that are lipid soluble, and so can move through the plasma membranes of target cells and act within the nuclei [62].

The endocrine system is composed of many different glands, and it can be divided into two categories: classical and non-classical. In the endocrine system, hormones are secreted into the circulatory system where they are distributed throughout the body, regulating bodily functions. The classical endocrine glands include the pituitary gland, pancreas, thyroid gland, adrenal cortex and medulla. The primary function of these glands is to manufacture specific hormones. Non-classical endocrine glands include the heart, hypothalamus, kidneys, liver and the gastrointestinal tract. Many of the classical hormones are controlled by the hypothalamus and pituitary, which can also be classified as being an extension of the nervous system [63].




Figure 6. Time-sequence photographs of a mineralising 3D PA matrix [61].

Gut–Brain Interactions

Gut–brain interactions are increasingly recognised as playing an important role in determining overall food intake [63]. Many peptides are synthesised and released from the gastrointestinal tract, and it has been shown that they physiologically influence eating behaviour via gut–brain signalling [64]. Ghrelin is an appetite-stimulating peptide produced in the stomach, which acts as a meal initiator. This differs from peptide YY, pancreatic polypeptide (PP), glucagon-like peptide-1 (GLP-1), oxyntomodulin (OXM) and cholecystokinin (CCK), which are all derived from the intestine and pancreas and have been shown to produce satiety signals. From this, it has been suggested that gut hormones can be manipulated to regulate energy balance, and as a result, gut hormone-based therapies could be a possible treatment for obesity (Figure 7) [65,66].




Figure 7. Interactions of gut and endocrine hormones with the brain and how they affect food intake [66].

Peptide Hormones Involved with the Gastro-intestinal Tract and Feeding

Leptin

Leptin is a hormone made by adipose cells that affects many biological mechanisms including reproduction, the immune and inflammatory response, haematopoiesis, angiogenesis, bone formation and wound healing. More interestingly, however, leptin helps to regulate energy balance by inhibiting hunger. This occurs via a feedback mechanism in which signals are sent to key regulatory centres in the brain to inhibit food intake [67].

After leptin is released by the adipose tissue into the bloodstream, it crosses the BBB and binds to the hypothalamic leptin receptors. This affects the activity of many hypothalamic neurones and the expression of various orexigenic (appetite stimulating) and anorexigenic neuropeptides. Orexigenic peptides include neuropeptide Y (NPY), and anorexigenic peptides include pro-opiomelanocortin. It has been suggested that the interaction with both types of these neuropeptides underpins the mechanism of action of leptin in the hypothalamus to inhibit hunger [67].

Ghrelin

Ghrelin is a 28-amino acid peptide with an octanoylated serine residue at position 3 [68] and is produced and secreted by cells within the oxyntic glands of the stomach [65]. Peripheral administration of ghrelin has been shown to stimulate food intake and decrease fat utilisation. This means it is involved in energy homeostasis, and it is the serine residue that appears to give ghrelin these effects [68]. What makes ghrelin unique is its function to increase food intake rather than decrease it, and as a result, it is a very important component of weight control. Evidence of this was shown when a study was carried out on mice that were lacking in ghrelin. The results showed that they were resistant to diet-induced obesity when fed a high-fat diet due to them eating less, and therefore utilising more stored fat as an energy source [66].

Cholecystokinin

Cholecystokinin is an endogenous gut hormone mainly found in the duodenum and jejunum, which exists in several molecular forms with differing numbers of amino acids. Examples include CCK-8 and CCK-54 (the number indicates the number of amino acid residues). CCK is known to act as a postprandial satiety signal, and it acts via two receptors: CCK₁ and CCK₂. The CCK₁ receptor is more important in appetite control [68]. The receptors are located on the peripheral vagal afferent terminals, which transmit signals to

the part of the brain stem that is associated with appetite, such as the nucleus of the solitary tract [66].

Oxyntomodulin

Oxyntomodulin is a 37-amino acid peptide expressed in the central nervous system and the L cells of the intestine and pancreas [69]. OXM seems to mediate its effects via the GLP-1 receptor as shown in experiments carried out on rat parietal cells [70]. This has been proven since its anorectic actions are blocked when the GLP-1 antagonist was administered [71]. Intravenous administration of OXM in humans inhibits gastric emptying and gastric acid secretion, which leads to a feeling of satiety [72]. This feeling of satiety can cause a reduction in both food intake and overall body weight, and this is brought about by the suppression of ghrelin.

Glucagon-like Peptide-1

Glucagon-like peptide-1 is a 30-amino acid gut-derived incretin peptide hormone [73] meaning that it stimulates insulin secretion in response to eating, and as a result, it suppresses glucagon secretion. In addition to this, GLP-1 inhibits gastric emptying and also reduces appetite and food intake [74]. GLP-1 is produced in the intestinal epithelial endocrine L cells in the distal small bowel and colon by differential processing of proglucagon [73,74]. Proglucagon is the gene that is expressed in the L cells and is regulated in the gut and brain [74]. Within minutes of food intake, the plasma levels of GLP-1 rise rapidly. GLP-1 exists in two circulating molecular forms: GLP-1(7–37) and GLP-1(7–36) amide, and it is GLP-1(7–36) amide that represents the majority of circulating active GLP-1 in human plasma. Both forms of GLP-1 are rapidly metabolised and inactivated by the enzyme dipeptidyl peptidase-4 (DPP-4) to GLP-1(9–37) or GLP-1(9–36) amide following the release from gut L cells [75]. This widely expressed enzyme cleaves both forms of GLP-1 at the position 2 alanine of the N-terminal to make them inactive. The expression of DPP-4 in the gut and vascular endothelium explains the short half-life of GLP-1 of just several minutes, because the majority of immunoreactive GLP-1 entering the portal venous circulation has already been inactivated by N-terminal cleavage [76].

Pancreatic Polypeptide

Pancreatic polypeptide is a 36-amino acid peptide that belongs to a family that includes NPY and peptide YY (PYY), and all of these peptides are members of the PP-fold peptide family. The PP-fold family binds to receptors Y_1 – Y_6 , but PP in particular has the highest affinity for the Y_4 and Y_5 receptors [68]. PP is similar to GLP-1 in that it is released into the circulation after the ingestion of food. However, it differs in that it is produced in the endocrine F cells, which are located in the periphery of the pancreatic islets [77,78]. PP is responsible for a number of regulatory actions, such as the inhibition of pancreatic exocrine secretion, and the modulation of gastric acid secretion, and gastric emptying [79,80]. The amount of PP released is affected by the digestive state, i.e. release is very low in the fasted state but is significantly increased throughout all phases of digestion. In addition to this, PP is affected by a decrease in blood glucose levels and insulin-induced hypoglycaemia, both being stimuli for PP secretion in the brain. As a result of this, it is thought that PP could potentially play a significant role in the regulation of feeding behaviour to control energy homeostasis [77].

Peptide YY

Peptide YY is a gut hormone that belongs to the PP family, along with PP and NPY, all of which are given the term PP-fold family. The PP-fold motif is found throughout this family and relates to the 3D structure. The PP-fold is formed through the incorporation of certain residues, which are predominately Pro2, Pro5, Pro8, Gly9, Tyr20 and Tyr27. The PP-fold has been found to protect the peptide against enzymatic attack as well as producing a hydrophobic pocket that is inherently overall energy-reducing. In addition to containing the PP-fold motif, PYY and its derivative PYY_{3–36} also have a high C-terminal α -helix content, which has also been suggested to be extremely important for the structural integrity of PYY. The sequence of the 36-amino acid peptide, PYY, is Tyr-Pro-Ala-Lys-Pro-Glu-Ala-Pro-Gly-Glu-Asp-Ala-Ser-Pro-Glu-Glu-Leu-Ser-Arg-Tyr-Tyr-Ala-Ser-Leu-Arg-His-Tyr-Leu-Asn-Leu-Val-Thr-Arg-Gln-Arg-Tyr-NH₂ [81].

Temperature-dependent circular dichroism (CD) studies have been carried out on human PYY to characterise its secondary structure. The results indicated an α -helical structure due to a maximum near 190 nm and two minima near 208 and 222 nm. A minimum near 200 nm was also observed, which corresponds to a random-coil structure (Figure 8) [82].

The PP-fold PP family all consists of a signal peptide, followed by a 36-amino acid active peptide and a carboxyl-terminal [83], and they mediate their effects through the NPY receptors Y_1 , Y_2 , Y_4 and Y_5 [84]. The Y-receptors belong to the G protein-coupled receptor family, and they mediate a wide variety of physiological effects such as regulation of blood pressure, anxiety, memory retention, hormone release and food intake.

Peptide YY is released by the L cells of the gastrointestinal tract following food intake, and there are two main endogenous forms: PYY_{1–36} and PYY_{3–36}. PYY_{1–36} is rapidly processed by the enzyme DPP-4 to the 34-amino acid peptide PYY_{3–36} [85]. DPP-4 hydrolyses PYY and removes the first two amino acids, tyrosine and proline, at the N-terminal, which changes the receptor selectivity. As a result of this, PYY_{3–36} has a high selectivity for the Y_2 -receptor, compared with PYY_{1–36}, which has selectivity for the Y_1 , Y_2 and Y_5 receptors. It is thought that the Y_1 receptor requires both the C-terminus and N-terminus for recognition, binding and then subsequent activation. The Y_2 receptor is thought to have a smaller receptor site and also only requires the C-terminus for recognition (Figure 9).

This could explain the reduced affinity for PYY_{3–36} on any Y receptor other than Y_2 [86]. Other studies replacing the amide bonds with ester bonds also confirm that the end section is important in binding and activation [87]. The Y_2 receptors are located in




Figure 8. Temperature dependant CD spectra of human PYY in 20 mM acetate buffer at pH 4.6 [82].




Figure 9. Binding sites for Y1 and Y2 with hPYY + hPYY_(3–36) [86].

the hippocampus, sympathetic and parasympathetic nerve fibres, intestines and certain blood vessels and have been implicated in regulating food intake and gastric emptying [88]. As a result of this, the Y₂ receptor is considered a target for the treatment of obesity and type 2 diabetes.

Neuropeptide Y

Neuropeptide Y is also a 36-amino acid peptide that has a very similar sequence homology to PYY and PP. NPY, however, differs from the other two peptides in the fact that it acts as a neurotransmitter rather than a hormone [89]. NPY is one of the most abundant peptides found in the brain [90], and it is synthesised and released by neurons, which in the peripheral nervous system are mostly sympathetic neurons [89]. NPY is associated with various biological responses, which include increased food intake, enhanced cognitive function associated with learning and memory and also reduction in anxiety [91]. In addition to this, in peripheral blood vessels, NPY has been shown to induce vasoconstriction [92]. Studies of NPY and its receptors suggest that it could be directly related to various pathological disorders such as obesity, depression and epilepsy [93].

Peptide Therapeutics

There are many peptide products on the market at present, and this are likely to increase because of their specificity, potency and low toxicity [94,95]. Table 1 shows some of the leading peptide therapeutics. Not all of these peptide therapeutics self-assemble or are formulated as solutions, but the table shows examples highlighting the importance of the use of peptides in drugs. The highest selling marketed diabetic drug Liraglutide incorporates a lipid chain to extent plasma circulation and ensures prolonged bioavailability [96,97]. Liraglutide is a GLP-1 agonist drug that self-assembles into an α -helical structure, and it requires once a day administration [98]. Lipid conjugation of a palmitoyl chain to a lysine residue at position 26 of liraglutide results in an extended half-life (around 13–14 h) in the blood. This is due to the palmitoyl chain allowing non-covalent binding to albumin, which delays

proteolytic attack by DPP-4 and also rapid renal clearance. Furthermore, the addition of the lipid chain could further prolong half-life by sterically hindering the DPP-4 enzyme from degradation [99].

Another peptide known to self-assemble is the octapeptide Lanreotide. This compound is a synthetic analogue of the peptide hormone somatostatin, and it is used to treat acromegaly [100] (a condition where the body produced too much growth hormone). In water, lanreotide self-assembles into monodisperse liquid crystalline nanotubes. The nanotubes are made up of dimers that self-assemble into a 2D crystal, which is held together by lateral chain interactions within antiparallel β -sheets [100,101].

Daptomycin is a lipopeptide therapeutic agent that is used as an antibiotic to treat infections caused by Gram-positive bacteria [102]. This drug is particularly effective against infections that are antimicrobial resistant such as MRSA. The antimicrobial activity of daptomycin is calcium-dependent. In the presence of a 1 : 1 ratio of calcium ions to daptomycin, self-assembly into micelles has been observed. The mechanism of action involves the disruption of the negatively charged bacterial cell membrane, in the presence of cations (for antibiotic activity, this is calcium ions) [53].

A further example of how self-assembly and bioactivity of peptide hormones are related involves self-assembling amyloid structures formed by peptide hormones and neuropeptides, which are crucial messenger molecules responsible for the function of different cells and organs. Peptide hormones and neuropeptides form aggregates that pack into dense-core vesicles (DCVs), which are used to temporarily store peptide messengers in secretory cells [103]. When DCVs are triggered, they release the stored contents into the blood or extracellular space [104], which results in amyloid disassembly, in order for action [103]. Therefore, for these types of peptides, reversibility of peptide aggregation is essential for their function.

As mentioned earlier, Table 1 shows some examples of peptide agents that are currently on the market.

Therapeutic Applications of PYY Peptides

A significant amount of research into the effects that PYY and PYY_(3–36) have on the body has been undertaken, predominately

in the food/biological science fields [129–131]. The application of PYY_(3–36) as an anti-obesity drug is an extremely interesting topic of research; however, it is limited because of its short half-life and lack of selectivity for the Y₂-receptor [85]. As a result, there is an unmet need for a treatment, and because peptide hormones already exist, *in vivo* peptide-based drugs are very favourable compared with synthetic molecules. This is due to them being less toxic, more selective and also more predictable in their *in vivo* behaviour [132]. In contrast, however, the use of peptides and proteins as therapeutic agents has its drawbacks because of their rapid degradation, excretion, and poor water solubility. In addition to this, they may cause allergic reactions due to immune responses. However, this applies more to proteins than peptides as peptides are generally less immunogenic. One of the main reasons for the short half-life of peptides is due to their rapid renal excretion, and this can be overcome by increasing the molecular weight (upper limited >2000 g mol^{−1}). A strategy to do this is to PEGylate the peptides. Covalently attaching PEG chains to the peptide improves enzymatic stability because of steric hindrance of proteolytic enzymes. This suppressed the immune response [132–134]. PEGylation has been shown to induce vacuolisation (formation of vacuoles within/adjacent to cells), but current research has focused on limiting this effect by using different carriers or shortening chain lengths [135–137]. In addition to extending half-life, PEGylation exhibits many properties that are favourable in pharmaceutical applications such as high water solubility, low toxicity and immunogenicity and ready clearance from the body. An example to show the effectiveness of PEGylation on half-life is the study by Lee *et al.*, where site-specific mono-PEGylation of GLP-1 led to a 16-fold increase in plasma half-life time in rats [138].

As previously mentioned, the ability of PYY and PYY_(3–36) to reduce food intake makes this peptide promising for use in anti-obesity or chronic eating disorder medications. A few studies have shown that even though PYY works as an appetite suppressant, it has some drawbacks that need to be overcome. A few studies that used high doses of PYY have reported taste aversion in animals and nausea in humans [139,140]. This was compounded by rapid dose administration. These side effects were combated by low doses at a steady and controlled infusion mainly through intravenous injections [141]. The two main problems with the administration of PYY seem to be both its concentration and how it is given, with the optimum dose being obtained by intravenous administration [142]. This is a detriment for the patient as they have to inject themselves, potentially daily, during the programme, and this can cause problems. Oral or nasal delivery gets over these issues, but with potential efficacy issues due to uptake [143,144].

A paper in 2002 commented on the potential for acute peripheral administration of PYY_(3–36) to inhibit food intake. PYY_(3–36) was injected into fasted (for 24 h) rats' hypothalamuses, and this led to a significant decrease in food intake, even at doses as low as 100 µg kg^{−1} [145]. This effect was initially controversial with very few other researchers being able to repeat the result [146]. A theory behind the controversy was proposed, this being that the mice used in the repeat tests were unaccustomed to the laboratory environment, which increased their stress levels. Stress inherently reduces baseline food intake so it makes any investigations in anorectic agents difficult to assess [146]. Further research into the use of PYY_(3–36) in human studies has found that obese subjects show normal levels of sensitivity to the appetite-reducing effects [147].

Summary and Conclusions

Peptides have many advantages as therapeutic agents. This is due to them being multifunctional and having a diverse range of applications; from antimicrobial agents to triggering a cellular release of hormones. Table 1 provides a list of examples of peptide drugs currently on the market.

Lipidation is a useful tool to reduce the degradation of a drug, or to extend half-life, with some of these effects probably being the result of self-assembly, although much more work is needed to investigate the possibility of many lipidated (and also non-lipidated) peptide drugs. Lipidation also enhances interactions of peptides with cell membranes.

As discussed earlier with appropriate examples, lipopeptides and PAs have a lot of interesting applications including use in antimicrobial and antifungal treatments, as skin care product ingredients, in drug delivery systems including those for hormone diseases as highlighted herein and in material templating in tissue engineering.

Peptide hormone therapeutics are of great interest as components of 'pre-built' drugs that the human body has already created and responds to. This significantly reduces the work needed to find a synthetic drug that responds to the active site. One drawback is that peptide hormones *in vivo* are stable for a specific time frame, before being removed. This may be a problem if the peptide hormone is of interest as a drug candidate, due to rapid clearance from the body. This is where lipidation may be beneficial as it can lead to a reduction in the drug clearance rate and proteolytic degradation. This offers great potential in the development of future therapeutics.

The peptide hormone PYY_(3–36) is released by the L cells of the gastrointestinal tract following food intake and has a high selectivity for the Y₂ receptor. The Y₂ receptor is associated with the regulation of food intake and gastric emptying. As a result of this, the use of PYY_(3–36) as a peptide hormone therapeutic to treat lifestyle conditions such as obesity and type 2 diabetes is a very promising area of research. As previously mentioned though, peptide hormones have short half-lives and are rapidly cleared from the body, which means that their activity as therapeutic agents will be short lived, and lipidation and PEGylation are strategies to overcome this.

In summary, this review has covered applications of peptide-based molecules, especially peptide amphiphiles and lipopeptides, with a particular focus on peptide hormones and their uses as drugs. We have highlighted the limited number of studies so far in which it has been suggested that self-assembly may influence bioactivity. Much more research is required into this fascinating subject as novel peptide-based molecules emerge as future therapeutic agents.

References

- 1 Hamley IW. Self-assembly of amphiphilic peptides. *Soft Matter* 2011; **7**: 4122.
- 2 Dehsorkhi A, Castelletto V, Hamley IW. Self-assembling amphiphilic peptides. *J. Pept. Sci.* 2014; **20**: 453–467. DOI: 10.1002/psc.2633.
- 3 Tu RS, Tirrell M. Bottom-up design of biomimetic assemblies. *Adv. Drug Deliv. Rev.* 2004; **56**: 1537–1563. DOI: 10.1016/j.addr.2003.10.047.
- 4 Giddi HS, Arunagirinathan MA, Bellare JR. Self-assembled surfactant nano-structures important in drug delivery: a review. *Indian J. Exp. Biol.* 2007; **45**: 133–159.
- 5 Torchilin VP. Micellar nanocarriers: pharmaceutical perspectives. *Pharm. Res.* 2007; **24**: 1–16. DOI: 10.1007/s11095-006-9132-0.

- 6 Kita-Tokarczyk K, Grumelard J, Haefele T, Meier W. Block copolymer vesicles – using concepts from polymer chemistry to mimic biomembranes. *Polymer* 2005; **46**: 3540–3563.
- 7 Hamley IW. Lipopeptides: from self-assembly to bioactivity. *Chem. Commun.* 2015; **51**: 8574–8583.
- 8 Zhou P, Deng L, Wang YT, Lu JR, Xu H. Different nanostructures caused by competition of intra and inter- β -sheet interactions in hierarchical self-assembly of short peptides. *J. Colloid Interface Sci.* 2016; **64**: 219–228.
- 9 Houston ME, Wallace A, Bianchi E, Pessi A, Hodges RS. Use of a conformationally restricted secondary structural element to display peptide libraries: a two-stranded α -helical coiled-coil stabilized by lactam bridges. *J. Mol. Biol.* 1996; **262**: 270–282.
- 10 Mandal D, Shirazi AN, Parang K. Self-assembly of peptides to nanostructures. *Org. Biomol. Chem.* 2014; **12**: 3544–3561. DOI: 10.1039/c4ob00447g.
- 11 Cui HG, Webber MJ, Stupp SI. Self-assembly of peptide amphiphiles: from molecules to nanostructures to biomaterials. *Biopolymers* 2010; **94**: 1–18. DOI: 10.1002/bip.21328.
- 12 Löwik DWPM, van Hest JCM. Peptide based amphiphiles. *Chem. Soc. Rev.* 2004; **33**: 234–245. DOI: 10.1039/b212638a.
- 13 Soukasene S, Toft DJ, Moyer TJ, Lu HM, Lee HK, Standley SM, Cryns VL, Stupp SI. Antitumor activity of peptide amphiphile nanofiber-encapsulated camptothecin. *ACS Nano* 2011; **5**: 9113–9121. DOI: 10.1021/nn203343z.
- 14 Boekhoven J, Stupp SI. Supramolecular materials for regenerative medicine. *Adv. Mater.* 2014; **26**: 1642–1659. DOI: 10.1002/adma.201304606.
- 15 Stupp SI. Bioactive nanostructures for regenerative medicine and cancer therapies. *Tissue Eng Pt A* 2008; **14**: 703–703.
- 16 Matson JB, Zha RH, Stupp SI. Peptide self-assembly for crafting functional biological materials. *Curr. Opin. Solid State Mater. Sci.* 2011; **15**: 225–235. DOI: 10.1016/j.cossms.2011.08.001.
- 17 Li A, Hokugo A, Yalom A, Berns EJ, Stephanopoulos N, McClendon MT, Segovia LA, Spigelman I, Stupp SI, Jarrahy R. A bioengineered peripheral nerve construct using aligned peptide amphiphile nanofibers. *Biomaterials* 2014; **35**: 8780–8790. DOI: 10.1016/j.biomaterials.2014.06.049.
- 18 Webber MJ, Berns EJ, Stupp SI. Supramolecular nanofibers of peptide amphiphiles for medicine. *Isr. J. Chem.* 2013; **53**: 530–554. DOI: 10.1002/ijch.201300046.
- 19 Shah RN, Shah NA, Del Rosario Lim MM, Hsieh C, Nuber G, Stupp SI. Supramolecular design of self-assembling nanofibers for cartilage regeneration. *Proc. Natl. Acad. Sci. U. S. A.* 2010; **107**: 3293–3298. DOI: 10.1073/pnas.0906501107.
- 20 Accardo A, Tesaro D, Mangiapià D, Pedone C, Morelli G. Nanostructures by self-assembled peptide amphiphile as potential selective drug carriers. *Pept. Sci.* 2007; **88**: 115–121.
- 21 Mnif I, Ghribi D. Review lipopeptides biosurfactants: mean classes and new insights for industrial, biomedical, and environmental applications. *Pept. Sci.* 2015; **104**: 129–147. DOI: 10.1002/bip.22630.
- 22 Singh P, Cameotra SS. Potential applications of microbial surfactants in biomedical sciences. *Trends Biotechnol.* 2004; **22**: 142–146. DOI: 10.1016/j.tibtech.2004.01.010.
- 23 Steller S, Vollenbroich D, Leenders F, Stein T, Conrad B, Hofemeister J, Jacques P, Thonart P, Vater J. Structural and functional organization of the fengycin synthetase multienzyme system from *Bacillus subtilis* b213 and A1/3. *Chem. Biol.* 1999; **6**: 31–41. DOI: 10.1016/S1074-5521(99)80018-0.
- 24 Hamley IW, Dehsorkhi A, Jauregi P, Seitsonen J, Ruokolainen J, Coutte F, Chataigné G, Jacques P. Self-assembly of three bacterially-derived bioactive lipopeptides. *Soft Matter* 2013; **9**: 9572–9578.
- 25 Hamley IW, Kirkham S, Dehsorkhi A, Castelletto V, Reza M, Ruokolainen J. Toll-like receptor agonist lipopeptides self-assemble into distinct nanostructures. *Chem. Commun.* 2014; **50**: 15948–15951.
- 26 Erridge C. Endogenous ligands of TLR2 and TLR4: agonists or assistants? *J. Leukoc. Biol.* 2010; **87**: 989–999. DOI: 10.1189/jlb.1209775.
- 27 Jin MS, Lee JO. Structures of the toll-like receptor family and its ligand complexes. *Immunity* 2008; **29**: 182–191. DOI: 10.1016/j.immuni.2008.07.007.
- 28 Omueti KO, Beyer JM, Johnson CM, Lyle EA, Tapping RI. Domain exchange between human Toll-like receptors 1 and 6 reveals a region required for lipopeptide discrimination. *J. Biol. Chem.* 2005; **280**: 36616–36625. DOI: 10.1074/jbc.M504320200.
- 29 Choe J, Kelker MS, Wilson IA. Crystal structure of human toll-like receptor 3 (TLR3) ectodomain. *Science* 2005; **309**: 581–585. DOI: 10.1126/science.1115253.
- 30 Mifsud EJ, Tan ACL, Jackson DC. TLR Agonists as modulators of the innate immune response and their potential as agents against infectious disease. *Front. Immunol.* 2014; **5**: 79. DOI: 10.3389/fimmu.2014.00079.article
- 31 Mauldin IS, Wang E, Deacon DH, Olson WC, Bao YD, Slingluff CL. TLR2/6 agonists and interferon- γ induce human melanoma cells to produce CXCL10. *Int. J. Cancer* 2015; **137**: 1386–1396. DOI: 10.1002/ijc.29515.
- 32 Pimentel-Nunes P, Roncon-Albuquerque R, Goncalves N, Fernandes-Cerqueira C, Cardoso H, Bastos RP, Marques M, Marques C, Sarmiento JA, Costa-Santos C, Macedo G, Pestana M, Dinis-Ribeiro M, Leite-Moreira AF. Attenuation of toll-like receptor 2-mediated innate immune response in patients with alcoholic chronic liver disease. *Liver Int.* 2010; **30**: 1003–1011. DOI: 10.1111/j.1478-3231.2010.02251.x.
- 33 Zeyer F, Mothes B, Will C, Carevic M, Rottenberger J, Nurnberg B, Hartl D, Handgretinger R, Beer-Hammer S, Kormann MSD. mRNA-mediated gene supplementation of toll-like receptors as treatment strategy for asthma *in vivo*. *PLoS One* 2016; **11**. DOI: 10.1371/journal.pone.0154001.
- 34 Wan WB, Cao L, Khanabdali R, Kalionis B, Tai XT, Xia SJ. The emerging role of HMGB1 in neuropathic pain: a potential therapeutic target for neuroinflammation. *J. Immunol. Res.* 2016. DOI: 10.1155/2016/6430423.
- 35 Shao L, Ma AQ, Li JW, Zhang P. TLR3 and TLR4 as potential clinical biomarkers for in-stent restenosis in drug-eluting stents patients. *Immunol. Res.* 2016; **64**: 424–430.
- 36 Castelletto V, Hamley IW, Perez J, Abezgauz L, Danino D. Fibrillar superstructure from extended nanotapes formed by a collagen-stimulating peptide. *Chem. Commun.* 2010; **46**: 9185–9187. DOI: 10.1039/c0cc03793a.
- 37 Jones RR, Castelletto V, Connon CJ, Hamley IW. Collagen stimulating effect of peptide amphiphile C₁₆-KTTKS on human fibroblasts. *Mol. Pharm.* 2013; **10**: 1063–1069. DOI: 10.1021/mp300549d.
- 38 Abu Samah NH, Heard CM. Topically applied KTTKS: a review. *Int. J. Cosmet. Sci.* 2011; **33**: 483–490.
- 39 Castelletto V, Hamley IW, Whitehouse C, Matts PJ, Osborne R, Baker ES. Self-assembly of palmitoyl lipopeptides used in skin care products. *Langmuir* 2013; **29**: 9149–9155. DOI: 10.1021/la401771j.
- 40 Pakalns T, Haverstick KL, Fields GB, McCarthy JB, Mooradian DL, Tirrell M. Cellular recognition of synthetic peptide amphiphiles in self-assembled monolayer films. *Biomaterials* 1999; **20**: 2265–2279.
- 41 Tysseling-Mattiace VM, Sahni V, Niece KL, Birch D, Czeisler C, Fehlings MG, Stupp SI, Kessler JA. Self-assembling nanofibers inhibit glial scar formation and promote axon elongation after spinal cord injury. *J. Neurosci.* 2008; **28**: 3814–3823.
- 42 Silva GA, Czeisler C, Niece KL, Beniash E, Harrington DA, Kessler JA, Stupp SI. Selective differentiation of neural progenitor cells by high-epitope density nanofibers. *Science* 2004; **303**: 1352–1355. DOI: 10.1126/science.1093783.
- 43 Freeman R, Boekhoven J, Dickerson MB, Naik RR, Stupp SI. Biopolymers and supramolecular polymers as biomaterials for biomedical applications. *MRS Bull.* 2015; **40**: 1089–1101. DOI: 10.1557/mrs.2015.270.
- 44 Harrington DA, Cheng EY, Guler MO, Lee LK, Donovan JL, Claussen RC, Stupp SI. Branched peptide-amphiphiles as self-assembling coatings for tissue engineering scaffolds. *J. Biomed. Mater. Res. A* 2006; **78a**: 157–167.
- 45 Perez CMR, Stephanopoulos N, Sur S, Lee SS, Newcomb C, Stupp SI. The powerful functions of peptide-based bioactive matrices for regenerative medicine. *Ann. Biomed. Eng.* 2015; **43**: 501–514.
- 46 Morgan CE, Dombrowski AW, Perez CMR, Bahnson ESM, Tsihlis ND, Jiang WL, Jiang Q, Vercammen JM, Prakash VS, Pitts TA, Stupp SI, Kibbe MR. Tissue-factor targeted peptide amphiphile nanofibers as an injectable therapy to control hemorrhage. *ACS Nano* 2016; **10**: 899–909. DOI: 10.1021/acs.nano.5b06025.
- 47 Webber MJ, Kessler JA, Stupp SI. Emerging peptide nanomedicine to regenerate tissues and organs. *J. Intern. Med.* 2010; **267**: 71–88. DOI: 10.1111/j.1365-2796.2009.02184.x.
- 48 Azmi F, Elliott AG, Marasini N, Ramu S, Ziora Z, Kavanagh AM, Blaskovich MAT, Cooper MA, Skwarczynski M, Toth I. Short cationic lipopeptides as effective antibacterial agents: design, physicochemical properties and biological evaluation. *Biorg. Med. Chem.* 2016; **24**: 2235–2241.
- 49 Zasloff M. Antimicrobial peptides of multicellular organisms. *Nature* 2002; **415**: 389–395. DOI: 10.1038/415389a.

- 50 Hancock REW, Sahl HG. Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. *Nat. Biotechnol.* 2006; **24**: 1551–1557.
- 51 Makovitzki A, Baram J, Shai Y. Antimicrobial lipopeptides composed of palmitoyl di- and tricationic peptides: *in vitro* and *in vivo* activities, self-assembly to nanostructures, and a plausible mode of action. *Biochemistry* 2008; **47**: 10630–10636. DOI: 10.1021/bi8011675.
- 52 Liu LH, Xu KJ, Wang HY, Tan PKJ, Fan WM, Venkatraman SS, Li LJ, Yang YY. Self-assembled cationic peptide nanoparticles as an efficient antimicrobial agent. *Nat. Nanotechnol.* 2009; **4**: 457–463. DOI: 10.1038/nnano.2009.153.
- 53 Kirkham S, Castelletto V, Hamley IW, Inoue K, Rambo R, Reza M, Ruokolainen J. Self-assembly of the cyclic lipopeptide daptomycin: spherical micelle formation does not depend on the presence of calcium chloride. *ChemPhysChem* 2016; **17**: 2118–2122. DOI: 10.1002/cphc.201600308.
- 54 Pogliano J, Pogliano N, Silverman JA. Daptomycin-mediated reorganization of membrane architecture causes mislocalization of essential cell division proteins. *J. Bacteriol.* 2012; **194**: 4494–4504. DOI: 10.1128/JB.00011-12.
- 55 Zhang P, Cheetham AG, Lin YA, Cui H. Self-assembled TAT nanofibers as effective drug carrier and transporter. *ACS Nano* 2013; **7**: 5965–5977. DOI: 10.1021/nn401667z.
- 56 Kostelnik KB, Els-Heindl S, Kloting N, Baumann S, von Bergen M, Beck-Sickinger AG. High metabolic *in vivo* stability and bioavailability of a palmitoylated ghrelin receptor ligand assessed by mass spectrometry. *Biorg. Med. Chem.* 2015; **23**: 3925–3932.
- 57 Schönauer R, Els-Heindl S, Fischer JP, Köbberling J, Riedl B, Beck-Sickinger AG. Adrenomedullin 2.0: adjusting key levers for metabolic stability. *J. Med. Chem.* 2016; **59**: 5695–5705. DOI: 10.1021/acs.jmedchem.6b00126.
- 58 Mazza M, Notman R, Anwar J, Rodger A, Hicks M, Parkinson G, McCarthy D, Daviter T, Moger J, Garrett N, Mead T, Briggs M, Schatzlein AG, Uchegbu IF. Nanofiber-based delivery of therapeutic peptides to the brain. *ACS Nano* 2013; **7**: 1016–1026. DOI: 10.1021/nn305193d.
- 59 Hartgerink JD, Beniash E, Stupp SI. Self-assembly and mineralization of peptide-amphiphile nanofibers. *Science* 2001; **294**: 1684–1688. DOI: 10.1126/science.1063187.
- 60 Spoerke ED, Anthony SG, Stupp SI. Enzyme directed templating of artificial bone mineral. *Adv. Mater.* 2009; **21**: 425–430. DOI: 10.1002/adma.200802242.
- 61 Sargeant TD, Guler MO, Oppenheimer SM, Mata A, Satcher RL, Dunand DC, Stupp SI. Hybrid bone implants: self-assembly of peptide amphiphile nanofibers within porous titanium. *Biomaterials* 2008; **29**: 161–171. DOI: 10.1016/j.biomaterials.2007.09.012.
- 62 Hutton JC, Siddle K. *Peptide Hormone Secretion/Peptide Hormone Action: A Practical Approach*. Oxford University Press: Oxford, 1991.
- 63 Neal JM. *How the Endocrine System Works*, (2nd edn). Wiley-Blackwell: Chichester, 2016.
- 64 Berthoud H-R, Morrison C. The brain, appetite, and obesity. *Ann. Rev. Psychology* 2008; **59**: 55–92.
- 65 Wren AM, Bloom SR. Gut hormones and appetite control. *Gastroenterology* 2007; **132**: 2116–2130. DOI: 10.1053/j.gastro.2007.03.048.
- 66 Coll AP, Farooqi IS, O'Rahilly S. The hormonal control of food intake. *Cell* 2007; **129**: 251–262. DOI: 10.1016/j.cell.2007.04.001.
- 67 Klok MD, Jakobsdottir S, Drent ML. The role of leptin and ghrelin in the regulation of food intake and body weight in humans: a review. *Obes. Rev.* 2007; **8**: 21–34. DOI: 10.1111/j.1467-789X.2006.00270.x.
- 68 Huda MSB, Wilding JP, Pinkney JH. Gut peptides and the regulation of appetite. *Obes. Rev.* 2006; **7**: 163–182. DOI: 10.1111/j.1467-789X.2006.00245.x.
- 69 Anini YJC, Chariot J, Nagain C. Oxyntomodulin inhibits pancreatic secretion through the nervous system in rats. *Pancreas* 2000; **20**: 348–360.
- 70 Schepp WDK, Riedel T, Schmidler J, Schaffer K, Classen M. Oxyntomodulin: a cAMP-dependant stimulus of rat parietal cell function via the receptor for glucagon-like peptide-1 (7-36)NH₂. *Digestion* 1996; **57**: 398–405.
- 71 Dakin CL, Gunn I, Small CJ, Edwards CM, Hay DL, Smith DM, Ghatei MA, Bloom SR. Oxyntomodulin inhibits food intake in the rat. *Endocrinology* 2001; **142**: 4244–4250. DOI: 10.1210/endo.142.10.8430.
- 72 Schjoldager B, Mortensen PE, Myhre J, Christiansen J, Holst JJ. Oxyntomodulin from distal gut. Role in regulation of gastric and pancreatic functions. *Dig. Dis. Sci.* 1989; **34**: 1411–1419.
- 73 Holst JJ. The physiology of glucagon-like peptide 1. *Physiol. Rev.* 2007; **87**: 1409–1439. DOI: 10.1152/physrev.00034.2006.
- 74 Drucker DJ, Nauck MA. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 2006; **368**: 1696–1705. DOI: 10.1016/S0140-6736(06)69705-5.
- 75 Drucker DJ. The biology of incretin hormones. *Cell Metab.* 2006; **3**: 153–165. DOI: 10.1016/j.cmet.2006.01.004.
- 76 Drucker DJ. Minireview: the glucagon-like peptides. *Endocrinology* 2001; **142**: 521–527. DOI: 10.1210/endo.142.2.7983.
- 77 Katsuura G, Asakawa A, Inui A. Roles of pancreatic polypeptide in regulation of food intake. *Peptides* 2002; **23**: 323–329.
- 78 Adrian TE, Besterman HS, Cooke TJ, Bloom SR, Bames AJ, Russell RC. Mechanism of pancreatic polypeptide release in man. *Lancet* 1997; **1**: 161–163.
- 79 Ri H. The pancreatic polypeptide (PP-fold) family: gastrointestinal, vascular, and feeding behavioural implication. *Exp. Biol. Med.* 1993; **203**: 44–63.
- 80 McTigue DM, Rogers RC. Pancreatic polypeptide stimulates gastric motility through a vagal-dependant mechanism in rats. *Neurosci. Lett.* 1995; **188**: 93–96.
- 81 Tatemoto K. Isolation and characterization of peptide YY (PYY), a candidate gut hormone that inhibits pancreatic exocrine secretion. *Proc. Natl. Acad. Sci.* 1982; **79**: 2514–2518.
- 82 Hegefelfd WA, Kuczerka K, Jas GS. Structural dynamics of neuropeptide hPYY. *Biopolymers* 2011; **95**: 487–502. DOI: 10.1002/bip.21608.
- 83 Blomqvist AG, Herzog H. Y-receptor subtypes – how many more? *Trends Neurosci.* 1997; **20**: 294–298.
- 84 Batterham RL, Bloom SR. The gut hormone peptide YY regulates appetite. *Ann. N. Y. Acad. Sci.* 2003; **994**: 162–168.
- 85 Ehrlich GK, Michel H, Truitt T, Riboulet W, Pop-Damkov P, Goelzer P, Hainzl D, Qureshi F, Lueckel B, Danho W, Conde-Knape K, Konkari A. Preparation and characterization of albumin conjugates of a truncated peptide YY analogue for half-life extension. *Bioconjug. Chem.* 2013; **24**: 2015–2024. DOI: 10.1021/bc400340z.
- 86 Nygaard R, Nielbo S, Schwartz TW, Poulsen FM. The PP-fold solution structure of human polypeptide YY and human PYY3-36 as determined by NMR. *Biochemistry* 2006; **45**: 8350–8357. DOI: 10.1021/bi060359l.
- 87 Albertsen L, Andersen JJ, Paulsson JF, Thomsen JK, Norrild JC, Stromgaard K. Design and synthesis of peptide YY analogues with c-terminal backbone amide-to-ester modifications. *ACS Med. Chem. Lett.* 2013; **4**: 1228–1232. DOI: 10.1021/ml400335g.
- 88 Keire DA, Bowers CW, Solomon TE, Reeve Jr JR. Structure and receptor binding of PYY analogs. *Peptides* 2002; **23**: 305–321.
- 89 Michel MC, Beck-Sickinger A, Cox H, Doods HN, Herzog H, Larhammar D, Quirion R, Schwartz T, Westfall T. XVI. International Union of Pharmacology recommendations for the nomenclature of neuropeptide Y, peptide YY, and pancreatic polypeptide receptors. *Pharmacol. Rev.* 1998; **50**: 143–150.
- 90 Chronwall BM, DiMaggio DA, Massari VJ, Pickel VM, Ruggiero DA, O'Donohue TL. The anatomy of neuropeptide-Y-containing neurons in rat brain. *Neuroscience* 1985; **15**: 1159–1181.
- 91 Dumont Y, Fournier A, Quirion R. Expression and characterization of the neuropeptide Y Y5 receptor subtype in the rat brain. *Neuroscience* 1998; **18**: 5565–5574.
- 92 Larhammar D. Structural diversity of receptors for neuropeptide Y, peptide YY and pancreatic polypeptide. *Regul. Pept.* 1996; **65**: 165–174.
- 93 Wahlestedt C, Reis DJ. Neuropeptide Y-related peptides and their receptors are the receptors potential therapeutic drug targets? *Annu. Rev. Pharmacol. Toxicol.* 1993; **32**: 309–352.
- 94 Craik DJ, Fairlie DP, Liras S, Price D. The future of peptide-based drugs. *Chem.Biol. Drug Design* 2013; **81**: 136–147.
- 95 Grant M, Leone-Bay A. Peptide therapeutics: it's all in the delivery. *Ther. Deliv.* 2012; **3**: 981–996.
- 96 Li Y, Shao MX, Zheng XM, Kong WL, Zhang JN, Gong M. Self-assembling peptides improve the stability of glucagon-like peptide-1 by forming a stable and sustained complex. *Mol. Pharm.* 2013; **10**: 3356–3365. DOI: 10.1021/mp4001734.
- 97 Gao Z, Bai G, Chen J, Zhang Q, Pan P, Bai F, Geng P. Development, characterization, and evaluation of a fusion protein of a novel glucagon-like peptide-1 (GLP-1) analog and human serum albumin in *Pichia pastoris*. *Biosci. Biotechnol. Biochem.* 2009; **73**: 688–694. DOI: 10.1271/bbb.80742.

- 98 Wang Y, Lomakin A, Kanai S, Alex R, Benedek GB. Transformation of oligomers of lipidated peptide induced by change in pH. *Mol. Pharm.* 2015; **12**: 411–419. DOI: 10.1021/mp500519s.
- 99 Frederiksen TM, Sonderby P, Ryberg LA, Harris P, Bukrinski JT, Scharff-Poulsen AM, Elf-Lind MN, Peters GH. Oligomerization of a glucagon-like peptide 1 analog: bridging experiment and simulations. *Biophys. J.* 2015; **109**: 1202–1213. DOI: 10.1016/j.bpj.2015.07.051.
- 100 Valery C, Artzner F, Robert B, Gulick T, Keller G, Grabielle-Madellmont C, Torres ML, Cherif-Cheik R, Paternostre M. Self-association process of a peptide in solution: from beta-sheet filaments to large embedded nanotubes. *Biophys. J.* 2004; **86**: 2484–2501. DOI: 10.1016/S0006-3495(04)74304-0.
- 101 Gobeaux F, Fay N, Tarabout C, Meneau F, Meriadec C, Delvaux C, Cintrat JC, Valery C, Artzner F, Paternostre M. Experimental observation of double-walled peptide nanotubes and monodispersity modeling of the number of walls. *Langmuir* 2013; **29**: 2739–2745. DOI: 10.1021/la304862f.
- 102 Baltz RH, Miao V, Wrigley SK. Natural products to drugs: daptomycin and related lipopeptide antibiotics. *Nat. Prod. Rep.* 2005; **22**: 717–741. DOI: 10.1039/b416648p.
- 103 Nespovita N, Gath J, Barylyuk K, Seuring C, Meier BH, Riek R. Dynamic assembly and disassembly of functional β -endorphin amyloid fibrils. *J. Am. Chem. Soc.* 2016; **138**: 846–856. DOI: 10.1021/jacs.5b08694.
- 104 Glombik MM, Gerdes H-H. Signal-mediated sorting of neuropeptides and prohormones: secretory granule biogenesis revisited. *Biochimie* 2000; **82**: 315–326.
- 105 Kolodny S, Ford C, Grinspan A, US Open-Label Glatiramer Acetate. Twenty years of continuous treatment of multiple sclerosis with glatiramer acetate 20 mg/ml daily: long-term disability outcomes in the US Open-Label Extension Study. *Mult Scler J* 2016; **22**: 417–417.
- 106 Mohammadi F, Rahimian R, Fakhraei N, Rezayat SM, Javadi-Paydar M, Dehpour AR, Afshari K, Mehr SE. Effect of glatiramer acetate on short-term memory impairment induced by lipopolysaccharide in male mice. *Fundam. Clin. Pharmacol.* 2016; **30**: 347–356. DOI: 10.1111/fcp.12202.
- 107 Guzman-Soto I, Salinas E, Quintanar JL. Leuprolide acetate inhibits spinal cord inflammatory response in experimental autoimmune encephalomyelitis by suppressing NF- κ B activation. *Neuroimmunomodulation* 2016; **23**: 33–40. DOI: 10.1159/000438927.
- 108 Rahimi M, Mobedi H, Behnamghader A. Aqueous stability of leuprolide acetate: effect of temperature, dissolved oxygen, pH and complexation with beta-cyclodextrin. *Pharm. Dev. Technol.* 2016; **21**: 108–115. DOI: 10.3109/10837450.2014.971377.
- 109 Candeias EM, Sebastiao IC, Cardoso SM, Correia SC, Carvalho CI, Placido AI, Santos MS, Oliveira CR, Moreira PI, Duarte AI. Gut-brain connection: the neuroprotective effects of the anti-diabetic drug Liraglutide. *World J. Diabetes* 2015; **6**: 807–827. DOI: 10.4239/wjd.v6.i6.807.
- 110 Rai Y, Iwata H, Masuda N, Anan K, Takeuchi T, Kohno N, Takei H, Yanagita Y, Noguchi S. Monthly versus three-monthly goserelin treatment in premenopausal patients with oestrogen receptor-positive early breast cancer. *EJC Suppl.* 2010; **8**: 193–193.
- 111 Sagara Y, Masuda N, Kinoshita T, Iwata H, Nakamura S, Yanagita Y, Nishimura R, Iwase H, Kamigaki S, Takei H, Noguchi S. The STAGE Study: A phase III comparison of anastrozole plus goserelin with tamoxifen plus goserelin as pre-operative treatments in premenopausal breast cancer patients. *Cancer Res.* 2010; **70**.
- 112 Gadelha MR, Bronstein MD, Brue T, Coculescu M, Fleseriu M, Guzelman M, Pronin V, Raverot G, Shimon I, Lieve KK, Fleck J, Aout M, Pedroncelli AM, Colao A, Grp PCS. Pasireotide versus continued treatment with octreotide or lanreotide in patients with inadequately controlled acromegaly (PAOLA): a randomised, phase 3 trial. *Lancet Diabetes Endo* 2014; **2**: 875–884.
- 113 Giustina A, Karamouzis I, Patelli I, Mazziotti G. Octreotide for acromegaly treatment: a reappraisal. *Expert. Opin. Pharmacother.* 2013; **14**: 2433–2447. DOI: 10.1517/14656566.2013.847090.
- 114 Chambers EE, Lame ME, Bardsley J, Hannam S, Legido-Quigley C, Smith N, Fountain KJ, Collins E, Thomas E. High sensitivity LC-MS/MS method for direct quantification of human parathyroid 1-34 (teriparatide) in human plasma. *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences* 2013; **938**: 96–104.
- 115 Gatti D, Rossini M, Viapiana O, Povino MR, Liuzza S, Fracassi E, Idolazzi L, Adami S. Teriparatide treatment in adult patients with osteogenesis imperfecta type I. *Calcif. Tissue Int.* 2013; **93**: 448–452. DOI: 10.1007/s00223-013-9770-2.
- 116 Fan H, Pan QR, Xu Y, Yang XC. Exenatide improves type 2 diabetes concomitant with non-alcoholic fatty liver disease. *Arq. Bras. Endocrinol. Metabol.* 2013; **57**: 702–708.
- 117 Qi F, Wu J, Fan QZ, He F, Tian GF, Yang TY, Ma GH, Su ZG. Preparation of uniform-sized exenatide-loaded PLGA microspheres as long-effective release system with high encapsulation efficiency and bio-stability. *Colloids Surf. B.* 2013; **112**: 492–498.
- 118 Chang YH, Chen WC, Hsieh PH, Chen DW, Lee MS, Shih HN, Ueng SWN. *In vitro* activities of daptomycin-, vancomycin-, and teicoplanin-loaded polymethylmethacrylate against methicillin-susceptible, methicillin-resistant, and vancomycin-intermediate strains of *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* 2011; **55**: 5480–5484. DOI: 10.1128/AAC.05312-11.
- 119 Larioza J, Girard A, Brown RB. Clinical experience with daptomycin for outpatient parenteral antibiotic therapy. *Am. J. Med. Sci.* 2011; **342**: 486–488. DOI: 10.1097/MAJ.0b013e31821e1e6b.
- 120 Kireev KA, Krasnopeev AV, Kireeva TS. Clinical case history of eptifibatide use during coronary intervention in patient with coronary failure. *Ration Pharmacother* 2015; **11**: 159–164.
- 121 Pancioli AM, Adeoye O, Schmit PA, Khoury J, Levine SR, Tomsick TA, Sucharew H, Brooks CE, Crocco TJ, Gutmann L, Hemmen TM, Kasner SE, Kleindorfer D, Knight WA, Martini S, McKinney JS, Meurer WJ, Meyer BC, Schneider A, Scott PA, Starkman S, Warach S, Broderick JP; CLEAR-ER investigators. Combined approach to lysis utilizing eptifibatide and recombinant tissue plasminogen activator in Acute Ischemic Stroke-Enhanced Regimen Stroke Trial. *Stroke* 2013; **44**: 2381–2387. DOI: 10.1161/STROKEAHA.113.001059.
- 122 Scatena R. Bivalirudin: a new generation antithrombotic drug. *Expert Opin. Investig. Drugs* 2000; **9**: 1119–1127.
- 123 Yamada T, Kurihara K, Ohnishi Y, Tamada T, Tomoyori K, Masumi K, Tanaka I, Kuroki R, Niimura N. Neutron and X-ray crystallographic analysis of the human alpha-thrombin-bivalirudin complex at pD 5.0: protonation states and hydration structure of the enzyme-product complex. *BBA-Proteins Proteom* 2013; **1834**: 1532–1538.
- 124 Cope SM, Sizemore SM, Roy A, Ghirlanda G, Vaiana SM. Structure and internal dynamics of calcitonin family peptides: implications for amyloid formation. *Biophys. J.* 2014; **106**: 687a–687a.
- 125 Deftos LJ, Nolan JJ, Seely BL, Clopton PC, Cote GJ, Whitham CL, Florek LJ, Christensen TA, Hill MR. Intrapulmonary drug delivery of bone-active peptides: bioactivity of inhaled calcitonin approximates injected calcitonin. *J. Bone Miner. Res.* 1996; **11**: 4–4.
- 126 Karsdal MA, Henriksen K, Arnold M, Christiansen C. Calcitonin – a drug of the past or for the future? Physiologic inhibition of bone resorption while sustaining osteoclast numbers improves bone quality. *BioDrugs* 2008; **22**: 137–144.
- 127 Caplin ME, Pavel M, Cwikla JB, Phan AT, Raderer M, Sedlackova E, Cadiot G, Wolin EM, Capdevila J, Wall L, Rindi G, Langley A, Martinez S, Gomez-Panzani E, Ruzniewski P, CLARINET Investigators. Anti-tumour effects of lanreotide for pancreatic and intestinal neuroendocrine tumours: the CLARINET open-label extension study. *Endocr. Relat. Cancer* 2016; **23**: 191–199. DOI: 10.1530/ERC-15-0490.
- 128 Sun LC, Yu CY, Mackey LV, Coy DH. Lanreotide and its potential applications in polycystic kidney and liver diseases. *Curr. Top. Med. Chem.* 2016; **16**: 133–140.
- 129 Batterham RL, Heffron H, Kapoor S, Chivers JE, Chandarana K, Herzog H, Le Roux CW, Thomas EL, Bell JD, Withers DJ. Critical role for peptide YY in protein-mediated satiation and body-weight regulation. *Cell Metab.* 2006; **4**: 223–233. DOI: 10.1016/j.cmet.2006.08.001.
- 130 Kaga T, Fujimiyama M, Inui A. Emerging functions of neuropeptide Y Y₂ receptors in the brain. *Peptides* 2001; **22**: 501–506.
- 131 Cabrele C, Beck-Sickinger AG. Molecular characterization of the ligand-receptor interaction of the neuropeptide Y family. *J. Pept. Sci.* 2000; **6**: 97–122. DOI: 10.1002/(SICI)1099-1387(200003)6:3<97::AID-PSC236>3.0.CO;2-E.
- 132 Bellmann-Sickert K, Elling CE, Madsen AN, Little PB, Lundgren K, Gerlach LO, Bergmann R, Holst B, Schwartz TW, Beck-Sickinger AG. Long-acting lipidated analogue of human pancreatic polypeptide is slowly released into circulation. *J. Med. Chem.* 2011; **54**: 2658–2667. DOI: 10.1021/jm101357e.
- 133 Toft DJ, Moyer TJ, Standley SM, Ruff Y, Ugolkov A, Stupp SI, Cryns VL. Coassembled cytotoxic and pegylated peptide amphiphiles form filamentous nanostructures with potent antitumor activity in models of breast cancer. *ACS Nano* 2012; **6**: 7956–7965. DOI: 10.1021/nn302503s.

- 134 Zhao T, Yang Y, Zong AZ, Tan HN, Song XL, Meng SO, Song CX, Pang GL, Wang FS. N-terminal PEGylation of human serum albumin and investigation of its pharmacokinetics and pulmonary microvascular retention. *Biosci. Trends* 2012; **6**: 81–88.
- 135 Prabhu S, Mutalik S, Rai S, Udupa N, Rao BSS. PEGylation of superparamagnetic iron oxide nanoparticle for drug delivery applications with decreased toxicity: an *in vivo* study. *J. Nanopart. Res.* 2015; **17**.
- 136 Casettari L, Vllasaliu D, Mantovani G, Howdle SM, Stolnik S, Illum L. Effect of PEGylation on the toxicity and permeability enhancement of chitosan. *Biomacromolecules* 2010; **11**: 2854–2865. DOI: 10.1021/bm100522c.
- 137 Zerbst C, Ring J, Frohlich M, Schliemann C, Mesters RM, Berdel WE, Schwoppe C. Site-directed and random PEGylation of retargeted tissue factor can improve the activity/toxicity profile of the molecule. *Cancer Res.* 2015; **75**.
- 138 Werle M, Bernkop-Schnürch A. Strategies to improve plasma half life time of peptide and protein drugs. *Amino Acids* 2006; **30**: 351–367. DOI: 10.1007/s00726-005-0289-3.
- 139 Chelikani PK, Haver AC, Reidelberger RD. Dose-dependent effects of peptide YY(3–36) on conditioned taste aversion in rats. *Peptides* 2006; **27**: 3193–3201. DOI: 10.1016/j.peptides.2006.08.001.
- 140 Degen L, Oesch S, Casanova M, Graf S, Ketterer S, Drewe J, Beglinger C. Effect of peptide YY_{3–36} on food intake in humans. *Gastroenterology* 2005; **129**: 1430–1436. DOI: 10.1053/j.gastro.2005.09.001.
- 141 Hurtado MD, Sergeyev VG, Acosta A, Spegele M, La Sala M, Waler NJ, Chiriboga-Hurtado J, Currlin SW, Herzog H, Dotson CD, Gorbatyuk OS, Zolotukhin S. Salivary peptide tyrosine-tyrosine 3–36 modulates ingestive behavior without inducing taste aversion. *J. Neurosci.* 2013; **33**: 18368–18380. DOI: 10.1523/JNEUROSCI.1064-13.2013.
- 142 Scott V, Kimura N, Stark JA, Luckman SM. Intravenous peptide YY_{3–36} and Y₂ receptor antagonism in the rat: effects on feeding behaviour. *J. Neuroendocrinol.* 2005; **17**: 452–457. DOI: 10.1111/j.1365-2826.2005.01330.x.
- 143 Olsen KM, Devlin JW. Comparison of the enteral and intravenous lansoprazole pharmacodynamic responses in critically ill patients. *Aliment. Pharmacol. Ther.* 2008; **28**: 326–333.
- 144 Laursen T, Grandjean B, Jorgensen JOL, Christiansen JS. Bioavailability and bioactivity of three different doses of nasal growth hormone (GH) administered to GH-deficient patients: comparison with intravenous and subcutaneous administration. *Eur. J. Endocrinol.* 1996; **135**: 309–315.
- 145 Batterham RL, Cowley MA, Small CJ, Herzog H, Cohen MA, Dakin CL, Wren AM, Brynes AE, Low MJ, Ghatei MA, Cone RD, Bloom SR. Gut hormone PYY_{3–36} physiologically inhibits food intake. *Nature* 2002; **418**: 650–654. DOI: 10.1038/nature02666.
- 146 Tschöp M, Castaneda TR, Joost HG, Thone-Reineke C, Ortmann S, Klaus S, Hagan MM, Chandler PC, Oswald KD, Benoit SC, Seeley RJ, Kinzig KP, Moran TH, Beck-sickinger AG, Koglin N, Rodgers RJ, Blundell JE, Ishii Y, Beattie AH, Holch P, Allison DB, Raun K, Madsen K, Wulff BS, Stidsen CE, Birringer M, Kreuzer OJ, Schindler M, Arndt K, Rudolf K, Mark M, Deng XY, Whitcomb DC, Halem H, Taylor J, Dong J, Datta R, Culler M, Craney S, Flora D, Smiley D, Heiman ML. Physiology: does gut hormone PYY_{3–36} decrease food intake in rodents? *Nature* 2004; **430**: 165.1 p following 165; discussion 2 p following
- 147 Batterham RL, Cohen MA, Ellis SM, Le Roux CW, Withers DJ, Frost GS, Ghatei MA, Bloom SR. Inhibition of food intake in obese subjects by peptide YY_{3–36}. *N. Engl. J. Med.* 2003; **349**: 941–948. DOI: 10.1056/NEJMoa030204.