University of Reading

A review of the financial impact of production diseases in poultry production systems

Article

Accepted Version

Jones, P. J. ORCID: https://orcid.org/0000-0003-3464-5424, Niemi, J., Christensen, J.-P., Tranter, R. B. ORCID: https://orcid.org/0000-0003-0702-6505 and Bennett, R. M. ORCID: https://orcid.org/0000-0003-3226-8370 (2019) A review of the financial impact of production diseases in poultry production systems. Animal Production Science, 59 (9). pp. 1585-1597. ISSN 1836-0939 doi: 10.1071/AN18281 Available at https://centaur.reading.ac.uk/78999/

It is advisable to refer to the publisher's version if you intend to cite from the work. See Guidance on citing.

To link to this article DOI: http://dx.doi.org/10.1071/AN18281

Publisher: CSIRO Publishing

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

University of Reading

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A review of the financial impact of production diseases in poultry production systems

Jones[1#], P.J., Niemi[2], J., Christensen[3], J-P., Tranter[1], R.B. and Bennett[1], R.M.

(#) Corresponding author: Tel: +44 (0)118 3788186; Email: p.j.jones@reading.ac.uk

(1) School of Agriculture, Policy and Development, University of Reading, PO Box 237, Whiteknight, Reading, RG6 6AR, England.

(2) Natural Resources Institute Finland (Luke), Bioeconomy and Environment Unit, Kampusranta 9, FI-60320 Seinäjoki, FINLAND.

(3) Department of Veterinary and Animal Sciences, Faculty of Health & Medical Sciences, University of Copenhagen, Stigbøjlen 4, 1870 Frederiksberg C, Denmark.

The financial impact of production diseases in poultry production systems

Abstract

Whilst the academic literature widely asserts that production diseases have a significant financial impact on poultry propduction, these claims are rarely supported by empirical evidence. There is a risk, therefore, that the information needs of poultry producers regarding the costs associated with particular diseases are not being adequately met.

A systematic literature review of poultry production diseases was undertaken, first to scope the availability of studies that estimate the financial impacts of production diseases on poultry systems and second, based on these studies, estimates were generated of the magnitude of these impacts. Nine production diseases, selected by a panel of stakeholders as being economically important in the EU, were examined.

The review found that the poultry disease literature has primarily an epidemiological focus, with very few publications provideing estimates of the financial impacts of diseases. However, some publications quantified the physical impacts of production diseases and control interventions, e.g. using measures such as output volumes, mortality rates, bacteria counts, etc. Using these data in standard financial models, partial financial analyses were possible for some poultry production diseases.

Coccidiosis and clostridiosis were found to be the most common production diseases in broiler flocks, with salpingoperitonitis the most common in layers. While the financial impact of untreated diseases varied, most uncontrolled diseases were estimated to make flocks loss-making. However, in all cases, interventions were available that signficantly reduced these losses. The review reinforces the concern that the available academic

literature is not providing sufficient information for poultry producers to decide on financially-optimal disease prevention and treatment measures.

Keywords: Poultry diseases; financial impacts; systematic literature review.

1. Introduction

There have been major changes in food consumption patterns in Western countries in the last 20 years, driven by increasing disposable incomes, changing food tastes and evolving health concerns (Traill, et al., 2014; European Commission, 2015). While egg consumption has remained fairly static (FAO, 2016a), there has been substantial growth in demand for poultry meat. Poultry meat is now the largest single source of meat-based protein in the diets of some countries, for example constituting 31% of all meat consumption in the UK and 43% in the USA in 2011 (FAO, 2016b). The chicken meat sector has responded to this increased demand by intensification of broiler production systems, involving more vertical integration, increases in production scale, use of new technologies and higher rates of input use, including higher stocking rates (FAO, 2016c).

A negative side of increasing production intensity has been a rise in the prevalence of so-called 'production diseases' in poultry systems. These usually originate from a complex interaction of pathogens, animal genetics and environment, including deficiencies in housing, nutrition and management. Production diseases constitute various infections, but also physical conditions, such as ascites, caused by genetic developments designed to increase physical performance, and physical damage caused by objects, or chemical irritants, in the rearing environment. What these diseases have in common is that, while they may be endemic, even in the wild, they can become increasingly problematic with the intensity of the production system and failures in management (Liverani, et al, 2013).

Production diseases compromise animal health and welfare and generate production inefficiencies, which can reduce profitability, and increase both environmental footprint and levels of antibiotic use. Bennett (2012) has provided a conceptual understanding of the way in which production diseases impact the economics of poultry production systems, i.e. through:

1. Economic impacts internal to the farm:

    - a loss of capital (i.e. animal mortality);
    - reduction in the level of marketable outputs;
    - reduction in (perceived or actual) output quality; and
    - waste of, or higher level of use of, inputs.

2. Economic impacts both internal, and external, to the farm:

    - resource costs associated with disease detection, diagnosis, prevention and control;
    - negative animal welfare impacts (i.e. animal suffering) associated with disease;
    - international trade restrictions due to disease and its control; and
    - human health costs associated with diseases or disease control.

3. Economic impacts external to the farm such as effects on rural economies and tourism.

With producer margins being squeezed by increasing costs and limited opportunity to transmit extra costs to consumers due to lack of market power, plus fierce competition from international suppliers, the response of the poultry industry has been to drive down those production costs that can be controlled, including disease costs (Narrod et al., 2008). To allow the industry to prioritise the most financially beneficial disease prevention and control measures, robust empirical data are required on: the risks posed by various production diseases; the financial impacts of different diseases; and the efficacy of, and financial benefits from, different disease control measures.

It might be be assumed that data to permit financially rational disease management decisions are available in the scientific literature. Much literature on poultry diseases exists, but data for individual diseases is seldom extensive and often lacking a financial dimension. Ubiquitous claims in research papers that particular poultry diseases lead to 'significant' financial impacts are seldom supported by empirical evidence. Consequently, data on the scale of financial losses associated with particular production diseases and the financial case for using control measures, are often lacking. Therefore, unless more informative industry data is available, there may be many poultry producers who are not implementing financially optimal disease prevention and treatment practices through lack of appropriate information. For example, in Denmark, vaccines are widely used to control salpingoperitonitis infections in layers without robust evidence of their efficacy (Christensen, 2016).

The study reported here undertook a systematic literature review to: determine the availability of data on the financial impacts of poulty production diseases; and a synthesis of this data to estimate the financial impacts of a number of production diseases and, where possible, the financial benefits of selected measures to control them. The study also had three sub-objectives. First, to show the relative risks presented by different production diseases, from data on their incidence. Second, to map the nature and distribution of disease costs, by showing where, in the production process, losses are occurring. Finally, to identify gaps in the literature on the financial impacts of poultry production diseases, to help guide future research.

2. Method

2.1 The choice of production diseases

To reduce the scope of the study, the most important production diseases were selected for analysis by a panel of 29 European animal scientists collaborating on the EU-funded PROHEALTH project.These came predominantly from veterinary medicine

123 or animal science backgrounds. Nine production diseases were identified as the most
124 important by virtue of rates of incidence, revenue losses, or control problems, i.e.
125 respiratory diseases (Ascites; Infectious bronchitis), enteric (Coccidiosis; Clostridiosis),
126 locomotory (Tibial dischrondoplasia; Foot pad dermatitis; Keel bone damage),
127 reproductive (Salpingoperitonitis) and other disorders (Injurious feather pecking).
128
129 2.2 The systematic literature review -
130 2.2.1 Introduction
131 A systematic review was undertaken to identify studies reporting financial or
132 productivity impacts of these nine production diseases/conditions. As a first step, a
133 Web of Science search was undertaken using a tailored search term with keywords to
134 capture:
135 (i) economic (or financial) studies;
136 (ii) poultry as study subjects;
137 (iii) specific production diseases;
138 (iv) exclusion of topics appearing in searches but not relevant to the review;
139 (v) exclusions to remove studies based on non-intensive production systems;
140 and
141 (vi) exclusions by text language, research domain, document type and
142 publication prior to 1995.
143
144 Abstracts found through the search were examined to exclude: duplications, those with
145 no physical performance measures or financial data, or were based on modelling
146 studies or reviews. This yielded 64 original studies. To supplement this list, additional
147 publications were found by: reviewing the reference lists of publications already
148 identified; a secondary web search using Google Scholar; website searches of
149 organisations with an interest in poultry health, such as the FAO; and reference lists
150 from recent poultry health research projects. This secondary search yielded a further

65 studies, making 129 in total. These publications encompassed peer-reviewed journals and conference proceedings, as well as 'grey' literature. Few publicatons assessed financial impacts, with most falling into the three categories shown in Sections 2.2.2 through 2.2.4.

2.2.2 Surveys of disease invidence and severity

A few studies surveyed the incidence of production diseases. Incidence, which is the number of (new) disease incidents (or outbreaks) over a specified period of time, can be viewed as an indicator of risk. Incidence might be reported for a particular flock, or as an average across flocks (e.g. average annual incidence). In the studies reviewed, flocks were generally only deemed to have experienced a disease outbreak when symptoms met a given severity criterion i.e. they either exhibited clinical symptoms, or where subclinical disease resulted in fiancial impacts. As we were only interested in disease episodes that cause financial losses, the analysis of incidence here was limited to those surveys where this criterion was explicitly used.

2.2.3 Studies exploring the impact of uncontrolled diseases on production

In this type of experimental study, birds could be deliberately exposed to a disease in either a controlled, or uncontrolled way. In the latter case, ambient levels of disease prevailed and therefore disease prevalence or severity was sometimes not elevated at all. Some of these studies employed a protected (or disease free) control group, while others did not.

2.2.4 Studies exploring the efficacy of measures to control production diseases

Intervention studies were the most common type of study in the reviewed literature. These involved trials of wide-ranging scale, from a few dozen birds to tens of thousands of birds across many poultry businesses. These studies had a variety of formats, depending on the:

- presence of a control group;
- presence of replicates;
- the number of interventions tested; and
- the level of control of environmental (rearing) conditions.

Studies with no control groups were excluded from the assessment. Where there were replicates of trials, averages over the replicates were calculated. When multiple, similar, interventions were used, for example several types of vaccine, an average over these interventions was taken. When multiple interventions were very different, for example contrasting a vaccine against a dietary nutrient, they were treated as separate interventions. When studies manipulated environmental conditions, in addition to target interventions, such as wetness of litter, then an average for the intervention over the multiple environmental conditions was estimated.

2.3 The standard financial models

Because financial data were rarely provided, the costs of diseases were estimated from data on changes to productive parameters (i.e. FCR, mortality and output volumes), using spreadsheet-based standard financial models for poultry enterprises. These were based on published data for market returns and production costs for EU 'average' conventional broiler and layer enterprises for 2013 (Appendix A).

2.5 Weighting of data

Recognizing that greater confidence can be placed on trials conducted on larger populations of birds, a weighting system was used in estimating averages across replicate trials. As studies often didn't state the exact number of birds in a trial, the value of the weights increases with size ranges using a geometric progression with a common ratio of two. By this means, data from experiments with up to 1,000 birds were

205 given a weight of one, 1,001-10,000 birds had a weight of two, 10,001-25,000 birds a
206 weight of four, and more than 25,000 birds a weight of eight.
207

208 ## 3. Results

209 3.1 The number of relevant studies identified from the systematic literature review
210 Table 1 lists the number of relevant studies identified for the nine study production
211 diseases, classified by the type of intervention used. Studies reporting no interventions
212 in Table 1 either examined the impacts of the uncontrolled disease, or were surveys of
213 disease incidence.
214

215 Table 1. The number of publications found reporting the impacts of poultry production
216 diseases and/or impacts of interventions to control them.

	Type of prevention/control intervention					
	None	Anti-microbials[1]	Vaccination	Housing	Other[3]	Total studies[2]
Respiratory diseases						
Pulmonary hypertension syndrome (ascites)	1	-	-	-	9	10
Infectious bronchitis (IB)	14	-	5	-	-	19
Enteric diseases						
Coccidiosis	1	7	8	-	-	16
Clostridiosis (*C. perfringens, C. septicum*)	1	10	3	-	1	15
Locomotory diseases						
Tibial dischrondoplasia	3	-	-	4	7	14
Foot pad dermatitis	3	-	-	-	12	15
Keel bone damage	10	-	-	3	5	18
Reproductive disorder						
Salpingoperitonitis syndrome, (colibacillosis)	9	1	-	-	-	10
Other disorders						

Injurious feather pecking	7	-	-	4	1	12
Total	45	18	16	11	35	129

[1] For either prophylactic or curative treatment.

[2] Some studies had multiple interventions, so the total number of studies may not equal the number of interventions.

[3] 'Other' usually involves changing parameters in the rearing environment, such as temperature, or humidity.

3.2 Disease incidence

This data came from studies ranging from large-scale surveys to small-scale laboratory trials. Because of the dominance of small-scale studies in the literature, the estimates in Table 2 should be treated with caution. Coccidiosis and clostridiosis would seem to be present in 90 - 100% of poultry flocks (Williams, 1998; Miller et al., 2010). There is a far greater incidence of the subclinical forms of these diseases, but these are only included in the incidence estimates where they cause productivity losses. The lowest reported disease incidence (at 5%) was reported for ascites, but most production diseases appear to have a reported incidence of over 30% of flocks.

Table 2. The incidence of production diseases and sources of this data

		Incidence (% of flocks)	Sources of data
Ascites	Broiler flocks	5	Hassanzadeh et al. (2005); Hassanzadeh et al. (2008); Maxwell and Robertson (1998)
Coccidiosis		90-100	Williams (1998, 1999)[1]
Clostridiosis		90-100	Miller et al. (2010)[1]
Footpad dermatitis		41.1	Allain et al. (2009); de Jong et al. (2014); Pagazaurtundua and Warriss (2006)
Tibial dyschondroplasia		35.6	Edwards (1990); Edwards and Sorensen (1987); Leeson et al. (1995); Lilburn and Lauterio (1989); Lui et al. (1992); Petek et al. (2005); Trablante et al. (2003); Yalcin et al. (2007)
Salpingoperitonitis	Laying flocks	49.5	Fossum et al. (2009)[2]

Injurious feather pecking	35	Lambton et al. (2013)

[1] Exact estimates of incidence for coccidiosis and clostridiosis are unavailable but sources indicate these infections are close to ubiquitous.

[2] Estimate of incidence of coliobaccilosis i.e. e-coli infections.

3.3 Mortality rates

Financial impacts resulting from elevated bird mortality comes from: loss of sales; expenditure on housing, feed and health care for birds that subsequently die; and the cost of disposal of carcasses. Once a disease is present in a flock, mortality rate is determined both by the severity of the disease challenge, and other factors such as the type of bird, breed, age at end of productive life-cycle and housing and production system, e.g. free-range. In an average commercial setting, with 'standard' disease management practice, cumulative mortality in layers, from all causes, ranges from 6 - 11%, with an average of 7.7% (van Horne, 2014; Weber et al., 2003; Merle et al., undated; Vitse et al., 2005; and Bell, 2012). Cumulative mortality in broilers is somewhat lower, ranging between 4 - 6% with an average of 4.7% (Havenstein et al., 2003; ACP, 2006; Gocsik et al., 2014; and van Horne and Bont, 2014). Table 3 shows the change in rate of mortality resulting from uncontrolled production diseases that are classified in studies as severe, i.e. where they have measureable financial impact. Also shown are the ranage of mortality values (in parentheses) found in the literature, where more than one usable estimate is available.

Table 3. Impact of severe uncontrolled production disease on flock mortality rates

	Mortality change (%) (range %)	**Sources of data**
Broilers		

Tibial dischondroplasia	+1	Morris (1993)
Acites	+36.3 (15.2 – 68)	Acar et al. (1995); Arce-Menocal et al. (2009); Camacho-Fernandez et al. (2002); Izadinia et al. (2010)
Clostridiosis	+336 (45.4 – 1500)	Lovland and Kaldhusdal (2001); Miller et al. (2010);Tactacan et al. (2013); Zhang et al. (2010)
Footpad dermatitis	+12.7 (-1 – 87.5)	Bilgili et al. (2009); Cengiz et al. (2011); de Jong et al. (2014); Ekstrand et al. (1997); Martland (1985); Mayne et al. (2007); Taira et al. (2013); Wang et al. (2010)
Laying flocks		
Keel bone damage	+71.5 (65.1 - 77.8)	Nasr et al. (2013); Petrik et al. (2015);
Salpingoperitonitis	+57	Jordan et al. (2005); Medina (2008), Thøfner et al. (2015)

Note: Change in mortality is the change to the base, or 'normal', mortality rate resulting from uncontrolled disease.

Note: Coccidiosis, Salpingoperitonitis and Injurious pecking are omited from the table due to lack of data.

Mortality impacts vary considerably between, and within, production diseases, and disease-driven mortality rates much higher than those in Table 3 have been observed in commercial practice. However, the headline observation is the paucity of studies on the mortality impacts of specific diseases in the literature. This problem is compounded by methodological weakness that affect the available data, i.e. some studies either have no experimental control, or they have a disease-challenged control, rather than a true (disease free) control. The lack of robustness in the available data is exemplified by the mortality impacts estimated for keel bone damage, which are considerably higher in the studies cited than have been observed by the authors in commercial farming practice.

3.4 Loss of physical outputs

Production diseases can lead to financial losses through reductions in the physical output from flocks (see Table 4). In broilers this can take the form of reduced terminal

277 weight (or rather, a longer growing period to reach the desired weight, requiring more
278 feeding and less efficient utilization of resources). In layers this would mean reduced
279 egg numbers, but also impairment of output quality. Loss of quality in broilers means
280 broken bones, damaged or discoloured muscle, or skin burns, leading to carcass
281 downgrades, or trimmings. In layers, this is experienced as smaller or mishapen eggs,
282 thin shells and colour change, resulting in downgrades or rejections.
283
284 Reviewed studies report reductions in terminal body weight in broilers range from zero
285 for ascites (although Swayne, 2013, suggests some weight loss is possible), to a high
286 of 17.7% for coccidiosis. There is a relatively high reported loss of body weight from
287 tibial dyschondroplasia. This effect is likely due to the fact that the condition can cause
288 considerable pain, and birds in pain move less and consume less food.
289
290 Table 4. Impact of severe and uncontrolled production disease on physical outputs

Broilers	**Live-weight (% change) (range)**	**Carcass downgrades (% change) (range)**	**Sources of data**
Tibial dischondroplasia	-10	<1	Burton et al. (1981); Edwards and Sorensen (1987); Morris (1993)
Acites	0	N.A.	Acar et al. (1995); Arce et al. (1992); Arce-Menocal et al. (2009); Camacho-Fernandez et al. (2002); Izadinia et al. (2010); Kalmar et al. (2013); Khajali et al. (2007); Maxwell and Morris (1992); Rincon (2000); Robertson (1998)
Clostridiosis	-1.24	N.A.	Lovland and Kaldhusdal (2001)
Coccidiosis	-17.7 (-17.3 - -18.1)	N.A.	Abdelrahman et al. (2014); Li et al. (2005)
Footpad dermatitis	-7.3 (0.8 – -14.6)	<1	Cengiz et al. (2011); de Jong et al. (2014); Martland (1985)
Laying Flocks	**Egg numbers**	**Egg weight**	**Egg quality**

Keel bone damage	-3.5 (-1.2 - -5.7)	-3.2		Nasr et al. (2012); Nasr et al (2013)
Infectious bronchitis	-32.9 (-3 - -50)	-8.7 (-7.3 - -11.36)	N.A.	Bisgaard M. (1976); Muneer et al. (1986); Muneer et al. (1987); Ignjatovic and Sapats (2000); Muneer et al. (2000)
Salpingoperitonitis	N.A.	N.A.	<1	Bisgaard and Dam (1981)
Injurious feather pecking	-5.1 (-2.6 - -7.5)	0	0	Glatz (2001); Hagger et al. (1989); Peguri and Coon (1993); Leeson and Morrison (1978)

Note: N.A. means that suitable data are not available.

Disease impacts on laying flocks (number of eggs) rangese between 3.5% and 32.9%, although greater losses may be observed in commercial practice. The impact of keel bone damage and injurious pecking on egg production should be low, unless birds contract secondary infections. In the case of feather pecking, feather loss means elevated loss of body heat, so that birds must eat more food to regulate body temperature and continue normal egg laying. While the impacts of infectious bronchitis can be severe, these effects last for only a small part of the productive life of a hen, typically 1-8 weeks. If a disease does not kill a hen, it will recover, and so, typically, will the laying percentage, although productivity may not always recover to pre-disease levels (Ignjatovic and Sapats, 2000; and Bisgaard, 1976). Based on available data, infectious bronchitis has the most significant impact on egg downgrades.

3.5 Impaired feed conversion ratio

All production diseases, if severe enough, impair birds' FCR i.e. they lower feed conversion efficiency. Where the bird cannot compensate by eating more, this can lead to loss of physical output. Where additional food is available and the bird has the capacity to consume it, physical outputs need not be reduced, but financial losses will still be experienced due to elevated feed consumption. Reductions in FCR ranged from zero for ascites to reductions of 25.9% for severe feather pecking (Table 5).

312

313 Table 5. Impact of severe, uncontrolled, production disease on the feed conversion

314 ratio (FCR)

	Reduction in FCR (%) (range)	Sources of data
Broilers		
Acites	0	Acar et al. (1995); Arce et al. (1992); Arce-Menocal et al. (2009); Camacho-Fernandez et al. (2002); Izadinia et al. (2010); Kalmar et al. (2013); Khajali et al. (2007); Maxwell and Robertson (1998); Morris (1992); Rincon, (2000)
Clostridiosis	16.4 (-3.7 – 70.5)	Lovland and Kaldhusdal (2001); Miller et al. (2010); Tactacan et al. (2013); Zhang et al. (2010)
Coccidiosis	17.7	Abdelrahman et al. (2014); Li et al. (2005)
Footpad dermatitis	3.3 (1.06 – 4.35)	Cengiz et al. (2011); de Jong et al. (2014)
Laying flocks		
Injurious feather pecking	25.9 (-5.1 - -49.7)	Glatz (2001); Leeson and Morrison (1978); Peguri & Coon (1993)

315 Note: Suitable data are not available for Tibial Dischondroplasia, Keel bone damage, Infectious bronchitis

316 and salpingoperitonitis.

317

318

319 3.6 Financial impacts of uncontrolled production diseases

320 The financial impacts of these diseases were estimated by applying percentage

321 changes in physical outputs to the standard broiler and layer financial models

322 (Appendix A). On the few occasions where data were available from the studies on

323 changes to input costs resulting from the diseases, or interventions, these were also

324 used in the financial models. For six of the diseases there were sufficient data to

325 undertake financial analyses, while for three there were not. In Figures 1 and 2, the

326 darker shaded bars represent the financial losses per bird, averaged over the flock,

327 arising from the uncontrolled diseases and the lighter bars show the losses that would

328 be incurred after applying the best available interventions to control them. Not

329 surprisingly, average losses for layers are higher than broilers because layers have a
330 longer productive life (around 56 weeks (RSPCA, 2016) and thus generate more
331 revenue. Broilers are usually slaughtered around 6-7 weeks in the EU and USA (EFSA,
332 2010; National Chicken Council, 2016), depending on growth rates and desired
333 slaughter weights.
334
335 Uncontrolled clostridiosis caused the greatest reported losses, at around €0.32 per bird
336 averaged over the flock, while losses from uncontrolled coccidiosis amounted to €0.21
337 per bird. Based on the financial model used here, confirmed by anecdotal industry
338 evidence, the net (profit) margin for a typical commercial broiler enterprise in the EU in
339 2013 was low, at around 10 Euro Cents per bird. With margins as tight as this, all of the
340 production diseases costed here would, when unconstrained, make affected flocks
341 loss-making.
342
343 Based on the standard financial model, laying hens typically generated a margin of
344 around €6 per bird in 2013. Figure 2 shows that, among the studied diseases, keel
345 bone damage causes the largest financial losses in laying hens, at around €3.5 per bird
346 averaged over the flock. However, this result should be treated with some caution in
347 view of the doubts raised above over the scale of mortality losses reported for this
348 disease.
349
350 A number of possible disease costs have not been accounted for, due to lack of data.
351 Typically, there are no data available from most disease impactstudies on labour, vet
352 and medicine costs, additional carcass disposal costs, or costs associated with the
353 disruption of normal husbandry practices resulting from diseases, such as delays to
354 thinning and depopulation to allow extra time for broilers to reach target weight. Also
355 not reported are losses from increased heterogeneity of broiler weights in a cohort,
356 meaning that a greater proportion of birds would fail to meet buyer requirements for

357 permissible weight range and would have to be sold at lower prices, often through
358 alternative marketing channels.

359

360 Figure 1. Financial losses due to four production diseases (controlled and uncontrolled) in
361 broiler flocks.

362



363

364

365 Figure 2. Financial losses due to two production diseases (controlled and uncontrolled) in
366 laying flocks.



367

368

369 3.7 The efficacy of interventions

370 Interventions to control production diseases in poultry are of two types, both adding to
371 production costs: treatment and prevention measures. Once a disease outbreak has
372 occurred, producers react with one or more courses of treatments, often with veterinary
373 support. Because many diseases are endemic, and difficult or expensive to control

once established, producers sometimes deploy preventive measures to try to reduce the risk of outbreaks and/or their severity. The cost of therapeutic treatments can be reduced if treatment begins early in a disease outbreak and so, producers may also increase expenditure on health monitoring to identify early signs of disease.

As Table 6 shows, many types of intervention have been evaluated in the literature, although there are few studies for any particular intervention. There is some heterogeneity within type of intervention studied for each disease. For example, in the anti-microbial category, treatments might be dietary supplements, probiotics, bacteriophage therapy, or antibiotics, with variation within these categories based on compounds or brands used, and concentrations of active ingredients. The data presented in Figures 1 and 2 represent the single most efficacious intervention reported in the literature for each disease. These estimates provide a sense of the higher end of the achievable levels of control that might be expected in a commercial setting.

Figures 1 and 2 show that there are considerable differences between these diseases in terms of both the financial losses caused when uncontrolled, and the extent to which interventions can reduce these losses. Tibial dyschondroplasia, for example, causes relatively small financial losses, but these are relatively difficult to eliminate. Conversely, diseases such as clostridiosis and infectious bronchitis, while resulting in very high financial costs when uncontrolled, can be reduced effectively through interventions. The diseases that would seem most problematic are those, such as keel bone damage, which lead to high financial costs when unconstrained and which resist attempts to control them. Based on this analysis, coccidiosis appears to fall into this class, with lower efficacy of interventions than for other diseases. However, producers report that both vaccines and anti-microbials offer significant means of disease control in a commercial setting.

Table 6. Types of intervention to control production diseases from the literature review.

	Class of measure	Types of intervention and data sources
Broilers		
Tibial dischondroplasia	Prevention	Manipulation of nutrients (Edwards, 1990) Manipulation of feed consumption (Edwards and Sorensen, 1987; Onbasilar et al., 2007) Manipulation of egg incubation temp. (Yalcin et al., 2007)
Acites	Prevention	Feed restriction – full rearing period (Arce et al. 1992; Camacho-Fernandez et al., 2002; Rincon 2000) Feed restriction – early weeks (Acar et al., 1995; Arce et al., 1992; Khajali et al., 2007)
Clostridiosis	Treatment	Antibiotics (Tactacan et al., 2013; Zhang et al., 2010) Bacteriophage therapy (Miller et al., 2010) Other antimicrobials (Tactacan et al., 2013)
Coccidiosis	Prevention	Vaccines (Lee et al., 2009; Li et al., 2005; Miguel et al., 2008; Shirley et al., 1995; Sou et al., 2006; Vermeulen et al., 2001; Williams et al., 1999; Williams and Gobbi, 2002) Probiotics (Abdelrahman et al., 2014) Herbal treatments (Miguel et al., 2008) Anticoccidials (Abdelrahman et al., 2014; Lee et al., 2009; Li et al., 2005; Miguel et al., 2008; Sou et al., 2006; Williams et al., 1999; Williams and Gobbi, 2002)
Footpad dermatitis	Prevention	Manipulation of litter moisture (Cengiz et al., 2011; de Jong et al., 2014; Ekstrand et al., 1997; Martland, 1985; Mayne et al., 2007; Taira et al., 2013; Wang et al., 2010) Variation of litter materials (Bilgili et al., 2009)
Laying flocks		
Keel bone damage	Prevention	Switch from unenriched to enriched cages (Petrik et al., 2015; Sherwin et al., 2010; Wilkins et al., 2011)
Infectious bronchitis	Prevention	Vaccines (Cook et al., 1999; Faramarzi et al., 2014; Jones et al., 2005; Tarpey et al., 2006; Tawfik et al., 2013)
Salpingoperitonitis	Prevention	Probiotics (Shini et al., 2013) Inoculation (Reid and Bocking, 2003) Vaccination (Gregersen, et al., 2010)
	Treatment	Antimicrobials (Balevi et al., 2001; Nahashon et al., 1996; Willis and Read, 2008)
Injurious feather pecking	Prevention	Housing (Fossum et al., 2009) Beak trimming (Craig and Lee,1990) Enriched environment (El-Lethey et al., 2000; Lambton et al., 2013) Reduced stocking rates (Nicol et al., 1999) Feed modification (Ambrosen and Petersen,1997)

There are two possible explanations for the discrepancy between the results of the scientific trials and real-world experience. First, that the few studies available are simply generating unrepresentative results and, second, and perhaps more likely, that the reviewed studies are capturing sub-clinical disease impacts. Observation of

410 commercial practice suggests that coccidiostats, such as ionophore antibiotics, while
411 effective at controlling clinical disease, are seemingly less effective at controlling
412 subclinical impacts, leading to losses through reduced feed intake and feed conversion
413 efficiency (Christensen, 2016).
414
415 **4. Discussion**
416 Our study found that there is an almost complete absence of published studies
417 generating data on the financial impacts of these nine poultry production diseases.
418 Generalising from this, it might be supposed that the entire poultry disease literature
419 has very much an epidemiological, rather than financial, focus.
420
421 To estimate the financial impacts of the nine poultry production diseases, and control
422 interventions, it was necessary to apply data on changes to productive parameters to
423 standard financial models (for broilers and layers) in order to monetise them. However,
424 there are significant gaps, even in the data on the impacts of diseases on productive
425 parameters, a case in point being salpingoperitonitis, where there are insufficient data
426 to permit any estimation of financial impact. This is perhaps explicable in view of the
427 tendency for salpingoperitonitis to occur in conjunction with other E. Coli-induced
428 conditions, such as airsaculitis, and secondary infections such as septicemia.
429
430 There is great heterogeneity of research objectives and methodology in the reviewed
431 studies, with some focusing on disease incidence, others on disease severity, others
432 seeking to capture the physical impacts of the disease itself, while others are
433 concerned only with the efficacy of control interventions. As a consequence of this,
434 together with the few studies, there is little or no replication in the literature and,
435 sometimes, essential data are only available from a single study. This limitation affects
436 the level of confidence that can be placed in the available data when generalising to
437 the whole sector.

438

439 The lack of focus on finanial impacts in studies means that, even if data on changes to
440 productive parameters are available and can be monetised, impacts on some cost
441 categories, such as vet and medicine costs, still cannot be captured. With very little
442 data on the impact of production diseases on the quality of outputs, the full financial
443 impact of downgrades to carcasses or eggs cannot be accounted for, and so disease
444 impacts may be underestimated. The lack of data on the cost of interventions means
445 that the estimates of the financial savings resulting from using them may be over-
446 estimated in our study.

447

448 Different studies often show a wide range of severity of impacts for the same disease.
449 More extreme impacts than estimated here might occur in commercial practice for a
450 number of reasons, including variations in: rearing environment; breed; management
451 quality; and the pathogenicity of infections. An additional cause of variation is the
452 occurrence of secondary infections. Most studies do not report data where secondary
453 infections are known to have occurred, on the grounds that such data would bias
454 impact estimates for the individual production diseases themselves. However, it must
455 be acknowledged that part of the set of negative consequences arising from the
456 occurrence of production diseases is an elevated risk of secondary infections from
457 other diseases.

458 For the reasons identified above, it is concluded that there are deficiencies in the
459 literature (and in the underlying reported research) resulting in data which are difficult
460 to use. Thus, the financial impacts estimated for the production diseases examined
461 here should be treated with some caution. Despite this, the claims made by many
462 authors in the poultry disease literature that production diseases can have significant
463 financial impacts would appear correct, even though these authors seldom supply any
464 empirical financial evidence supporting these claims.

While poultry farms with elevated levels of production diseases can make substantially less profit than farms with low disease levels, these losses can be significantly reduced by a range of prevention measures, such as vaccinations, or improved litter management, nutrition and hygiene, as well as curative treatments. The financial benefits of interventions to control production diseases vary greatly according to disease and the intervention chosen. The losses associated with diseases such as clostridiosis, for example, can be significantly reduced through use of antimicrobials, but others, such as keel bone damage, present a greater challenge.

The reliance of the poultry industry on the use of antimicrobials to control infectious diseases highlights the risks to the financial sustainability of the sector from the continuing growth in farm bacterial reservoirs with resistance to antimicrobial treatments (Aminov and Mackie, 2007; Sykes, 2010, EFSA and ECDPC, 2016).

These risks occur on three fronts. First, some antibiotics commonly used for the treatment of diseases may lose their efficacy. Second, government action plans, such as the EU Action Plan Against the Rising Threats from Antimicrobial Resistance (EU, 2011), which are designed to drive more responsible use of antibiotics, may make some antibiotics less readily available. Third, although there have been few official bans on the use of selected antibiotics so far, such as the US ban on Fluoroquinolones (FDA, 2005), governments may adopt the 'precautionary principle', and issue complete bans on the use of some antibiotics.

5. Conclusions

In light of this growing threat, there is a pressing need for the poultry research community to help identify cost-effective alternatives to antibiotics which offer similar levels of disease control. These could include: novel substances to strengthen the

493 poultry immune response to bacterial infection; naturally occurring bacteriophages;
494 novel vaccinations; and enhanced biosecurity measures on farm. Although some
495 rigorous individual studies of alternative approaches have been undertaken, there is
496 insufficient data across the literature to evaluate them. Failure to develop these
497 alternatives could significantly, and negatively, impact the future financial sustainability
498 of the global poultry industry.
499
500 There are strong hints in the literature that some interventions, particularly in relation to
501 biosecurity measures, reduce disease incidence, prevalence and severity, for multiple
502 production diseases simultaneously. The use of single interventions to control multiple
503 diseases would be very advantageous for an industry faced with small profit margins,
504 volatile markets, and the possibility of further regulation. The industry would, therefore,
505 benefit from a more holistic effort from the research community to identify the most
506 useful and cost-effective multi-functional interventions to reduce disease-related
507 financial losses.
508
509 The analysis above has revealed a disconnect between the requirements of the poultry
510 industry for data on the financial impacts of diseases and control measures and the
511 goals of researchers in the non-commercial poultry disease research community. As a
512 consequence, the value of such research, even if it targets relevant production
513 diseases and interventions, is of less value than it could be. In view of this, the question
514 might reasonably be asked, where are commercial producers and their advisors getting
515 the data on which to plan their disease management programmes?
516
517 In order to meet the future informational needs of the poultry industry, the focus of
518 academic poultry disease research needs to be changed. Studies need to generate
519 data not only on the first-order physical impacts of production diseases, but also
520 secondary and financial impacts, as is currently already being achieved commonly in

research on pig and dairy cow diseases. This means collecting data from abbatoirs on the impact of diseases on product quality, as well as data from farm trials and lab-based experiments on changes to the levels of input use resulting from diseases and the interventions to control them. This would require a more inter-disciplinary approach to research, involving not just veterinarians or animal scientists, but also agricultural economists.

Acknowledgement

The work on which this paper is based was made possible by funding from the EU's DG Research Framework 7 Programme PROHEALTH project http://www.fp7-prohealth.eu/

Conflict of interest

The authors know of no conflict of interest in relation to the production or publication of this article.

References

Abdelrahman, W., Mohnl, M., Teichmann, K., Doupovec, B., Schatzmayr, G., Lumpkins, B., Mathis, G., 2014. Comparative evaluation of probiotic and salinomycin effects on performance and coccidiosis control in broiler chickens. Poultry Sci. 93, 3002-3008.

Acar, N., Sizemore, F.G., Leach, G.R., Wideman, R.F., Owen, R.L., Barbatio, G.F.,1995. Growth of broiler chickens in response to feed restriction regimens to reduce ascites. Poultry Sci. 74, 833-843.

Agro Business Consultants Ltd., 2012. ABC Budgeting & Costings Book, 83rd ed. November 2016. Melton Mowbray, UK.

548

549 Allain, V., Mirabito, L., Arnould, C., Colas, M., Le Bouquin, S., Lupo, C., Michel, V.,
550 2009. Skin lesions in broiler chickens measured at the slaughterhouse: relationships
551 between lesions and between their prevalence and rearing factors. Brit. Poultry Sci. 50,
552 407-417.

553

554 Ambrosen, T., Petersen, V.E., 1997. The influence of protein level in the diet on
555 cannibalism and quality of plumage of layers. Poultry Sci, 76, 559–563.

556

557 Aminov, R.I., Mackie, R.I., 2007. Evolution and ecology of antibiotic resistance genes.
558 FEMS Microbiol. Lett. 271, 147-61.

559

560 Arce, J., Berger, M., Lopez Coello, C., 1992. Control of ascites syndrome by feed
561 restriction techniques. J. Appl. Poultry Res. 1, 1-5.

562

563 Arce-Menocal, J. Avila-Gonzalez, E., lopez-Cello, C., Garibay-Torres, G., Martinez-
564 Lemus, L.A. 2009. Body weight, feed-particle size and ascites incidence revisited. J.
565 Appl. Poultry Res. 18, 465-471.

566

567 Assured chicken production (ACP) standards, 2006. Key health and welfare indicators
568 for broiler production. http://www.journals.cambridge.org/ (accessed May 2016)

569

570 Balevi, T., Ucan, U.S., Coskun, B., Kurtoglu, V., Cetingul, I.S., 2001. Effect of dietary
571 probiotic on performance and humoral immune response in layer hens. Brit. Poultry
572 Sci. 42, 456-461.

573

574 Bell, D., 2012. U.S. experiences with Lohmann Selected Leghorn (LSL-Lite) layers,
575 Part 3: Livability. Lohmann Information, 47, 22-33.

576

577 Bennett, R. (2012) Economic rationale for interventions to control livestock disease.
578 Eurochoices, 11 (2), 5-10.
579 Bilgili, S.F., Hess, J.B., Blake, J.P., Macklin, K.S., Saenmahayak, B., Sibley, J.L., 2009.
580 Influence of bedding material on footpad dermatitis in broiler chickens. J. Appl. Poultry
581 Res. 18, 583-589.
582
583 Bisgaard, M., 1976. The influence of infectious bronchitis virus on egg production,
584 fertility, hatchability and mortality rate in chickens. Nor. Vet. Med. 28, 368-376.
585
586 Bisgaard, M., Dam, A., 1981. Salpingitis in poultry II: prevalence, bacteriology and
587 possible pathogenesis in egg laying chickens. Nord. Vet. Med. 33, 81-89.
588
589 BPC. 2016. The BPC antibiotic stewardship scheme: leading the way in the
590 responsible use of antibiotics. British Poultry Council. April 2016.
591 http://www.britishpoultry.org.uk/wp-
592 content/uploads/2016/04/The_BPC_Antibiotic_Stewardship_Scheme_April2016.pdf
593 (accessed May 2016)
594
595 Burton, R.W., Sheridan, A.K., Howlett, C.R., 1981. The incidence and importance of
596 tibial dyschondroplasia to the commercial broiler industry in Australia. Brit. Poultry Sci.
597 22, 153-60.
598
599 Camacho-Fernandez, D., Lopez, C., Avilla, E., Arce, J., 2002. Evaluation of different
600 dietary treatments to reduce ascites syndrome and their effects on corporal
601 characteristics in broiler chickens. J. Appl. Poultry Res. 11, 164-174.
602
603 Christensen, J-.P. (2016) Personal Communication.

Cengiz, O., Hess, J.B., Bilgili, S.F., 2011. Effect of bedding type and transient wetness on footpad dermatitis in broiler chickens. J. Appl. Poultry Res. 20, 554-560.

Cook, J.K.A., Orbell, S.J., Woods, M.A., Huggins, M.B., 1999. Breadth of protection of the respiratory tract provided by different live-attenuated infectious bronchitis vaccines against challenge with infectious bronchitis viruses of heterologous serotypes. Avian Path. 28, 477-485.

Craig, J.V., Lee, H.Y., 1990. Beak trimming and genetic stock effects on behavior and mortality from cannibalism in white leghorn-type pullets. Appl. Anim. Behav. Sci. 25, 107-123.

de Jong, I.C., Gunnink, H., van Harn, J., 2014. Wet litter not only induces footpad dermatitis but also reduces overall welfare, technical performance and carcass yield in broiler chickens. J. Appl. Poultry Res. 23, 51-58.

European Commission (2015) World food consumption patterns – trends and drivers. EU Agricultural Markets Briefs, No. 6, June 2015. https://ec.europa.eu/agriculture/markets-and-prices/market-briefs_en

Edwards, H.M., 1990. Efficacy of several vitamin D compounds in the prevention of tibial dyschondroplasia in broiler chickens. J. Nutr. 120, 1054-1061.

Edwards, H.M., Sorenssen, P., 1987. Effect of short fasts on the development of tibial dyschondroplasia in chickens. J. Nutr. 117, 194-200.

632 EFSA, 2010. Scientific opinion on the influence of genetic parameters on the welfare
633 and the resistance to stress of commercial broilers. EFSA J. 8, 1666.
634
635 EFSA (European Food Safety Authority) and ECDC (European Centre for Disease
636 Prevention and Control), 2016. The European Union summary report on antimicrobial
637 resistance in zoonotic and indicator bacteria from humans, animals and food in 2014.
638 EFSA J. 14, 4380.
639
640 Ekstrand, C, Algers, B., Svedberg, J., 1997. Rearing conditions and foot-pad dermatitis
641 in Swedish broiler chickens. Prev. Vet. Med. 31, 167-174.
642
643 El-Lethey, H., Aerni, V., Jungi, T.W., Wechsler, B., 2000. Stress and feather pecking in
644 laying hens in relation to housing conditions. Brit. Poultry Sci. 41, 22-28.
645 EUROSTAT, 2016. http://ec.europa.eu/eurostat/statistics-
646 explained/index.php/Agricultural_accounts_and_prices (accessed: August 2016).
647
648 EU, 2011. EU Action plan against the rising threats from antimicrobial resistance,
649 COM (2011) 748.
650 http://ec.europa.eu/dgs/health_consumer/docs/communication_amr_2011_748_en.pdf
651 (accessed May 2016).
652
653 FDA, 2005.
654 http://www.fda.gov/AnimalVeterinary/SafetyHealth/RecallsWithdrawals/ucm042004.htm
655 (accessed May 2016).
656
657 Follet, G., 2000. Antibiotic resistance in the EU - science, politics, and policy.
658 AgBioForum. 3, 148-155.
659

660 FAO, 2016a. Livestock in the balance, change in the livestock sector.
661 http://www.fao.org/docrep/005/y4252e/y4252e07.htm (accessed May 2016).
662
663 FAO, 2016b. http://faostat.fao.org/site/610/DesktopDefault.aspx?PageID=610#ancor
664 (accessed 25.05.16).
665
666 FAO, 2016c. http://www.fao.org/docrep/005/y4252e/y4252e07.htm (accessed
667 25.05.16).
668
669 Faramarzi, S., Bijanzad, P., Javaherzadeh, V., Moomivand, H., Stabraghi, E., Dehnavi,
670 E.N., Ghaedi, A., Zarghami, A., 2014. Evaluation of two different infectious bronchitis
671 vaccination programmes in broiler breeder chickens. Int. J. Biosci. 5, 210-216.
672
673 Fossum, O., Hansson, D.S., Engelsen Etterlin, P. and Vagsholm, I., 2009. Causes of
674 mortality in laying hens in different housing systems in 2001 to 2004. Acta Vet. Scand.
675 51, 3.
676
677 Glatz, P.C., 2001. Effect of poor feather cover on feed intake and production of aged
678 laying hens. Asian-Australian J. Anim. Sci. 14, 553-558.
679
680 Gocsik, E., Kortes, H.E., Oude Lansink, A.G.J.M., Saatkamp, H.W., 2014. Effects of
681 different broiler production systems on health care costs in the Netherlands. Poultry
682 Sci. 93, 1301-1317.
683
684 Gregersen, R.H., Christensen, H., Ewers, C., Bisgaard, M., 2010. Impact of
685 Escherichia coli vaccine on parent stock mortality, first week mortality of broilers and
686 population diversity of E. Coli in vaccinated flocks. Avian Pathol. 39, 287-295.
687

688 Hagger, C, Marguerat, C., Steiger-Stafl, D., Stranzinger, G., 1989. Plumage condition,
689 feed consumption and egg production relationships in laying hens. Poultry Sci. 68, 221-
690 225.
691
692 Havenstein, G.B., Ferket, P.R., Qureshi, M.A., 2003. Growth, livability, and feed
693 conversion of 1957 versus 2001 broilers when fed representative 1957 and 2001
694 broiler diets. Poultry Sci. 82, 1500-8.
695
696 Havenstein, G.B., Ferket, S.E., Scheideler, S.E., Larson, B.T., 1994. Growth, livability
697 and feed conversion of 1957 vs 1991 broilers when fed 'typical' 1957 and 1991 broiler
698 diets. Poultry Sci. 73, 1875-1794.
699
700 Hassanzadeh, M., Gilanpour, H., Charkkar, S., Buyse, J., Decuypere, E., 2005.
701 Anatomical parameters of cardiopulmonary system in three different lines of chickens:
702 further evidence for involvement in ascites syndrome. Avian Pathol. 34, 1.6.
703
704 Hassanzadeh, M., Buyse, J., Decuypere, E., 2008. Further evidence for the
705 involvement of anatomical parmeters of cardiopulmonary system in the development of
706 ascites syndrome in broiler chickens. Acta Vet.Hung. 71, 71-80.
707
708 Ignjatovic, J., Sapats, S., 2000. Avian infectious bronchitis virus. Rev. Sci. Tech. OIE.
709 19, 493-508.
710
711 Izadinia, M., Nobakht, M., Khajali, F., Faraji, M., Zamani, F., Qujeq, D., Karimi, I., 2010
712 Pulmonary hypertension and ascites as affected by dietary protein source in broiler
713 chickens reared in cool temperature at high altitudes. Anim. Feed Sci.Tech. 155, 194-
714 200.
715

716 Jordan F.T.W., Williams, N.J., Jones, W.T., 2005. Observations on salpingitis,
717 peritonitis and salpingoperitonitis in layer breeder flock. Vet. Rec. 157, 573-577.
718
719 Kalmar, I.D., Vanrompay, D., Janssens, G.P.J., 2013. Broiler ascites syndrome:
720 collateral damage from efficient feed to meat conversion. Vet. J. 197, 169-174.
721
722
723 Khajali, F., Zamani-Moghaddam, A., Asadi-Khoshoei, E., 2007. Application of an early
724 skip-a-day feed restriction on physiological parameters, carcass traits and development
725 of ascites in male broilers reared under regular or cold temperatures at high altitude.
726 Anim. Sci. 78, 159-163.
727
728 Lambton, S.L., Nicol, C.J., Friel, M., Main, D.C.J., McKinstry, C.M, Sherwin, J., Weeks,
729 C.A., 2013. A bespoke management package can reduce levels of injurious pecking in
730 loose-housed laying hen flocks. Vet. Rec. 172, 423.
731
732 Lee, J.T., Broussard, C., Fitz-Coy, S., Burke, P., Eckert, N.H., Stevens, S.M.,
733 Anderson, P.N., Caldwell, D.J., 2009. Evaluation of live oocyst vaccination or
734 salinomycin for control of field-strain Eimeria challenge in broilers on two different
735 feeding programs. J. Appl. Poultry Res. 18, 458-464.
736
737 Leeson, S., Gonzalo, J., Summers, J.D., 1995. Poultry metabolic disorders and
738 mycotoxins. University Books, Guelph, Ontario, Canada.
739
740 Leeson, S., Morrison, W.D., 1978. Effect of feather cover on feed efficiency in laying
741 birds. Poultry Sci. 57, 1094-1096.
742

743 Lilburn, M.S., Lauterio, T.J., 1989. Relationships among mineral balance in the diet,
744 early growth manipulation and incidence of tibial dyschondroplasia in different strains of
745 meat type chickens. Poultry Sci. 68, 1263-1273.
746
747 Liverani, M., Waage, J., Barnett, T., Pfeiffer, D.U., Rushton, J., Rudge, J.W.,
748 Loevinsohn, M.E., Scoones, E., Smith, R.D., Cooper, B.S., White, L.J. Goh, S., Horby,
749 P., Wren, B., Gundogdu, O., Woods, A., and Coker, R.J (2013) Understanding and
750 Managing Zoonotic Risk in the New Livestock Industries. Understanding and Managing
751 Zoonotic Risk in the New Livestock Industries. Environmental Health Perspectives,
752 121, 873–877.
753
754 Lovland, A., Kaldhusdal, M., 2001. Severely impaired production performance in broiler
755 flocks with high incidence of Clostridium perfringens-associated hepatitis. Avian Pathol.
756 30, 73-81.
757
758 Lui, L., Tong, J., Huang, J., 1992. Effect of dietary chloride and magnesium on on the
759 incidence of tibial dyschondroplasia in chickens fed on Chinese practical diets. Brit.
760 Poultry Sci. 3: 603-611.
761
762 Martland, M.F., 1985. Ulcerative dermatitis in broiler chickens: the effects of wet litter.
763 Avian Pathol. 14, 353-364.
764
765 Maxwell, M.H., Robertson, G.W., 1998. UK survey of broiler ascites and sudden death
766 syndromes in 1993. Brit. Poultry Sci. 39, 203-215
767
768 Mayne, R.K., Else, R.W., Hocking, P.M., 2007. High litter moisture alone is sufficient to
769 cause footpad dermatitis in growing turkeys. Brit. Poultry Sci. 48, 538-545.
770

771 Medina, H.A. 2008. Factors that could indice peritonitis in commercial egg layers.
772 Zootechnica, 2 October 2008. http://www.zootecnicainternational.com/article-
773 archive/veterinary/234-factors-that-could-induce-peritonitis-in-commercial-egg-layers-
774 .html (accessed May 2016)
775
776 Miguel, J.A., Asenjo, B., Ciria, J., del Cacho, E., Calco, J.L., 2008. Comparison of
777 control methods for coccidiosis in native Spanish 'Castellana Negra' chickens. Spanish
778 J. Agr. Res. 6, 531-536.
779
780 Miller, R.W., Skinner, E.J., Sulakvelidze, A., Mathis, G.F., Hofacre, C.L., 2010.
781 Bacteriophage therapy for control of necrotic enteritis of broiler chickens experimentally
782 infected with clostridium perfringens. Avian Dis. 54, 33-40.
783
784 Morris, M.P.,1992. Ascites in broilers. Poultry Int. October, 26-32.
785
786 Morris, M.P., 1993. National survey of leg problems. Broiler Industry 93 (May): 20-24.
787
788 Muneer, M.A., Halvorson, D.A., Sivanandan, V., Newman, J.A., and Coon, C.N.,1986.
789 Effects of infectious bronchitis virus (Arkansas strain) on laying chickens. Avian Dis. 30,
790 644-7.
791
792 Muneer, M.A., Newman, J.A., Halvorson, D.A., Sivanandan, V. and Coon, C.A., 1987.
793 Effects of Avian Infectious Bronchitis Virus (Arkansas Strain) on Vaccinated Laying
794 Chickens. Avian Dis. 31(4), 820-828.
795
796 Muneer, M.A., Chaudhry, K.M. and Khawaja, K.M., 2000. Losses due to infectious
797 bronchitis virus infection in laying and breeding hens. Pakistan Vet. J. 20, 64-70.
798

799 Nahashon, S.N., Nakaue, H.S. and Mirosh, L.W., 1996. Performance of single comb
800 white Leghorn fed a diet supplemented with a live microbial during the growth and egg
801 laying phases. Anim. Feed Sci. Tech. 57, 25-38.
802
803 Narrod, C., Tiongco, M. and Costales, A., 2008. Global poultry sector trends and
804 external drivers of structural change.
805 http://www.fao.org/AG/againfo/home/events/bangkok2007/docs/part1/1_1.pdf
806 (accessed 25 May, 2016).
807
808 National Chicken Council, (2016) http://www.nationalchickencouncil.org/about-the-
809 industry/statistics/u-s-broiler-performance/ (accessed19.05.16).
810
811 Nasr, M.A.F., Murrell, J., Wilkins, L.J., Nicol, C.J., 2012. The effect of keel fractures on
812 egg-production parameters, mobility and behaviour in individual laying hens. Anim.
813 Welfare 21, 127-135.
814
815 Nasra, M.A.F., Murrella, J., Nicol, C.J., 2013. The effect of keel fractures on egg
816 production, feed and water consumption in individual laying hens. Brit. Poultry Sci. 54,
817 165-170.
818
819 Nicol, C.J., Gregory, N.G., Knowles, T.G., Parkman, I.D.,Wilkins, L.J., 1999. Differential
820 effects of increased stocking density, mesiated by increased flock size, on feather
821 pecking and aggression in laying hens. Appl. Anim. Behav. Sci. 65, 137-152.
822
823 Onbasilar, E.E., Erol, H. Cantekin, Z., Kaya, U.,2007. Influence of intermittent lighting
824 on broiler performance, incidence of tibial dyschondroplasia, tonic immobility, some
825 blood parameters and antibody production. Asian-Australian J. Anim. Sci. 20, 550-555.
826

827 Pagazaurtundua, A., Warriss, P.D., 2006. Levels of foot pad dermatitis in broiler
828 chickens reared in 5 different systems. Brit. Poultry Sci. 47, 529-535.
829
830 Peguri, A., Coon, C., 1993. Effect of feather coverage and temperature on layer
831 performance. Poultry Sci. 72, 1318-1329.
832
833 Petek, M. Sonmez, G., Yildiz, H., Baspinar, H., 2005. Effects of different management
834 factors on broiler performance and incidence of tibial dyschondroplasia. Brit. Poultry
835 Sci. 46, 16-21.
836
837 Petrik, M.T., Guerin, M.T., Widowski, T.M., 2015. On-farm comparison of keel fracture
838 prevalence and other welfare indicators in conventional cage and floor-housed laying
839 hens in Ontario, Canada. Poultry Sci. 94, 579–585.
840
841 Reid, G. and Bocking, A., 2003. The potential for probiotics to prevent bacterial
842 vaginosis and preterm labor. Am. J. Obstet. Gynecol. 189, 1202-8.
843
844 Rincon, M.U., 2000. Mild feed restriction and compensatory growth in the broiler
845 chicken. MSc thesis, University of Guelph, Guelph, Canada.
846
847 RSPCA, 2015. Laying hens - farming (egg production).
848 http://www.rspca.org.uk/adviceandwelfare/farm/layinghens/farming (accessed
849 19.05.16).
850
851 RSPCA (2016) http://www.rspca.org.uk/adviceandwelfare/farm/layinghens/farming
852 (accessed 19.05.16).
853

Rural Business Research (RBR), 2014. Poultry production in England: Farm Business Survey 2012/13. Crane, R., Davenport, R., Laney, S., Vaughan, R. RBR, February 2014.

Sherwin, C.M., Richards, G.J., Nicol, C.J., 2010. Comparison of the welfare of layer hens in 4 housing systems in the UK. Brit. Poultry Sci. 51, 488-499.

Shini S., Shini, A., Blackall, P J., 2013. The potential for probiotics to prevent reproductive tract lesions in free-range laying hens. Anim. Prod. Sci. 53, 1298-1308.

Shirley, M.W., Bushell, A.C., Bushell, J.E., McDonald, V., Roberts, B., 1995. A live attenuated vaccine for the control of avian coccidiosis: trials in broiler breeders and replacement layer flocks in the UK. Vet. Rec. 137, 453-457.

Sou, X., Zhang, J.X., Li, Z.G., Yang, C.T., Min, Q.R., Xu, L.T., Lui, Q. And Zhu, X.Q., 2006. The efficacy and economic benefits of Supercox®, a live anticoccidial vaccine in a commercial trial in broiler chickens in China. Vet. Parasitol. 142, 63-70.

Swayne, D.E., 2013. Diseases of poultry, 13th ed, in: Swayne, D.E. (Ed.), Glisson, J.R., McDougald, L.R., Nolan, L.K., Suarez, D.L., Nair, V. (Associate Eds.), Wiley-Blackwell, New York, USA.

Sykes, R., 2010. The 2009 Garrod lecture: the evolution of antimicrobial resistance: a Darwinian perspective. J. Antimicrob. Chemoth. 65, 1842-52.

Tactacan, G.B., Schmidt, J.K., Miille, M.J., Jimenez, D.R., 2013. A Bacillus subtilis (QST 713) spore-based probiotic for necrotic enteritis control in broiler chickens. J. Appl. Poultry Res. 22, 825-831.

Taira, K, Nagai, T., Obi, T., Takase, K., 2013. Effect of litter moisture on the development of footpad dermatitis in broiler chickens. J. Vet. Med. Sci. 76, 583-586.

Tarpey, I., Orbell, S.J., Britton, P., Casais, R., Hodgson, T., Lin, F., Hogan, E., Cavanagh, D., 2006. Safety and efficacy of infectious bronchitis virus used for chicken embryo vaccination. Vaccine 24, 6830-6838.

Tawfik, H.I., Salama, E., Hassan, O.M., Ahmed, A.,2013. Preparation and evaluation of live bivalent bronchitis vaccine in chicken. Researcher 5, 31-35.

Thøfner, I., Ladefoged Poulsen, L., Bisgaard, M., Christensen, H., Heidemann Olsen, R., Christensen, J.P., 2015. Longitudinal study of mortality observed in four broiler breeder flocks. Paper presented at 19th WVPA Congress, Cape Town, 7-11.11.15.

Trablante, N.L. Estevez, I., Russek-Cohen, E., 2003. Effect of perches and stocking density on tibial dyschondroplasia and bone mineralization as measured by bone ash in broiler chickens. J. Appl. Poultry Res. 12, 53-59.

Traill,W. B., Mazzocchi, M., Shankar, B., Hallam, D. (2014) Importance of government policies and other influences in transforming global diets. Nutrition Reviews, 72(9), 591–604.

Vanhemelrijck, J., 1999. Risk management: from antibiotic discovery to market. Paper presented at the OIE Scientific Conference, Paris, France. March, 1999.

908 Van Horne P.M.L., 2014. Competitiveness of the EU egg sector: international
909 comparison base year 2013. Report LEI 2014-041. LEI Wageningen UR, Wageningen,
910 December 2014.
911
912 Van Horne, P.L.M, Bont, N., 2014. Competitiveness of the EU poultry meat sector:
913 International comparison base year 2013. LEI Report 2013-068, LEI Wageningen UR,
914 The Hague, December, 2013.
915
916 Vits, A., Weitzenburger, D., Hamann, H., Distl, O., 2005. Production, egg quality, bone
917 strength, claw Length, and keel bone deformities of laying hens housed in furnished
918 cages with different group sizes. Poultry Sci. 84, 1511–1519.
919
920 Wang, G. Ekstrand, C., Svedberg, J., 2010. Wet litter and perches as risk factors for
921 the development of footpad dermatitis in floor-housed hens. Brit. Poultry Sci. 39, 191-
922 197.
923
924 Vermeulen, A.N., Schaap, D.C., Schetters, T.P., 2001. Control of coccidiosis in
925 chickens by vaccination. Vet. Parasitol. 100, 13-20.
926
927 Wilkins, L.J., McKinstry, J.L., Avery, N.C., Knowles, T.G., Brown, S.N., Tarlton, J.,
928 Nicol, C.J., 2011. Influence of housing system and design on bone strength and keel
929 bone fractures in laying hens. Vet. Rec. 169, 414.
930
931 Williams, R.B., 1998. Anticoccidial vaccines for broiler chickens: pathways to success.
932 Avian Pathol. 31, 317-353.
933
934 Williams, R.B., 1999. A compartmentalised model for the estimation of the cost of
935 coccidiosis to the world's chicken production industry. Int. J. Parasitol. 29, 1209-1229.

Williams, R.B., Carlyle. W.W., Bond, D.R., Brown, I.A., 1999. The efficacy and economic benefits of Paracox, a live attenuated anticoccidial vaccine, in commercial trials with standard broiler chickens in the UK. Int. J. Parasitol. 29, 341-355

Williams, R.B., Gobbi, L., 2002. Comparison of an attenuated anticoccidial vaccine and an anticoccidial drug programme in commercial broiler chickens in Italy. Avian Pathol. 31, 253-265.

Willis, W.L., Read, L., 2008. Investigating the effects of dietary probiotic feeding regimens on broiler chicken production and Campylobacter jejuni presence. Poultry Science 87: 606-611.

Yalcin, S., Molayoglu, H.B., Baka, M., Genin, O., Pines, M., 2007. Effect of temperature during the incubation period on tibial growth plate chondrocyte differentiation and the incidence of tibial dyschondroplasia. Poultry Sci. 86, 1772-1783.

Zhang, G., Mathis, G.F., Hofacre, C.L., Yaghmaee, P., Holley, R.A., Duranc, T.D., 2010. Effect of a radiant energy-treated lysozyme antimicrobial blend on the control of clostridial necrotic enteritis in broiler chickens. Avian Dis. 54, 1298-1300.

Appendix A. Standard financial models for broiler and layer enterprises, 2013

	Broilers	Layers
Sales:	**Revenues (€/ 100 kg live weight)** [6]	**Revenues (€ / hen)** [6]
Broilers, (2.276 g of meat per bird at €107.7/100 kg liveweight); Layers, 340 eggs at €7.6/100 eggs [7]	107.7	25.84
Spent hens	-	0.36
Expenditure:	**Production costs (€/ 100 kg live weight)** [10]	**Production costs (€ / hen)** [8, 9]
Day old chicks / pullets (17 weeks)	15.20	3.30
Mortality[1]	2.02	0.87
Feed	67.00	10.29
Medication [5]	1.40	0.09 [4]
Heating and electricity	2.20	1.41 [4]
Water	0.60	
Litter (incl. cleanout & disposal)	3.70	
Labour	3.40	1.10
Housing [2]	6.40	2.75
General [3]	1.00	0.41
Total costs	102.92	20.22
Net margin	**4.74**	**5.98**

[1] Mortality costs assumed to be 50% of total rearing costs per dead bird. Mortality rate for layers assumed to be 9%.

[2] Housing costs includes: poultry house and inventory.

[3] General costs include: insurance, office, consultancy, telephone, transport.

[4] Medication, heating and electricity, water and litter costs are equated with the 'Other variable costs' category of Van Horne (2014), which includes: heating, electricity, litter, animal health and catching.

[5] Medication costs for broilers taken from Cocsik et al. (2014); layers from RBR (2014).

[6] 2013 broiler meat and egg prices; Eurostat Median of EU28 prices (authors' own calculations)

http://ec.europa.eu/eurostat/statistics-explained/index.php/Agricultural_accounts_and_prices

971 [7] Number of eggs produced per housed bird = 340 (source: van Horne, 2014), based
972 on enriched cage system).
973 [8] Sources: Van Horne (2014); Agro-Business Consultants Ltd (2012); RBR (2014).
974 [9] 2013 prices (based on 2010 (Van Horne, 2014) prices adjusted for inflation using
975 annual GDP deflators for the EU. Source: World Bank National Accounts Data & OECD
976 National Accounts files 2010-2015)
977 [10] Sources: Van Horne (2014); Agro-Business Consultants Ltd (2012).
978