

Ecological intensification: bridging the gap between science and practice

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Kleijn, D., Bommarco, R., Fijen, T. P. M., Garibaldi, L. A., Potts, S. G. ORCID: <https://orcid.org/0000-0002-2045-980X> and van der Putten, W. H. (2019) Ecological intensification: bridging the gap between science and practice. Trends in Ecology & Evolution, 34 (2). pp. 154-166. ISSN 0169-5347 doi: 10.1016/j.tree.2018.11.002 Available at <https://centaur.reading.ac.uk/82030/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.tree.2018.11.002>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Ecological intensification: bridging the gap between science and practice

David Kleijn^{1*}, Riccardo Bommarco², Thijs P.M. Fijen¹, Lucas A. Garibaldi³, Simon G. Potts⁴ & Wim H. van der Putten^{5,6}

¹ *Wageningen University, Plant Ecology and Nature Conservation Groups, Wageningen, The Netherlands*

² *Swedish University of Agricultural Sciences, Department of Ecology, SE-75007 Uppsala, Sweden*

³ *Instituto de Investigaciones en Recursos Naturales, Agroecología y Desarrollo Rural (IRNAD), Sede Andina, Universidad Nacional de Río Negro (UNRN) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Mitre 630, CP 8400, San Carlos de Bariloche, Río Negro, Argentina.*

⁴ *Centre for Agri-Environment Research, School of Agriculture, Policy and Development, University of Reading, UK*

⁵ *Department of Terrestrial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, The Netherlands*

⁶ *Laboratory of Nematology, Wageningen University and Research Centre, Wageningen, The Netherlands*

* Corresponding author: Kleijn, D. (David.Kleijn@wur.nl)

Keywords: agricultural production, biodiversity, ecosystem services, farmer behaviour

Abstract

There is worldwide concern about the environmental costs of conventional intensification of agriculture. Growing evidence suggests that ecological intensification of mainstream farming can safeguard food production with accompanying environmental benefits, however, the approach is rarely adopted by farmers. Our review of the evidence for replacing external inputs with ecosystem services shows that scientists tend to focus on processes (e.g. pollination) rather than outcomes (e.g. earnings), and express benefits at spatio-temporal scales that are not always relevant to farmers. This results in mismatches in perceived benefits of ecological intensification between scientists and farmers which hinders its uptake. We provide recommendations for overcoming these mismatches and highlight important additional factors driving uptake of nature-based management practices such as social acceptability of farming.

Ecological intensification shows potential to sustainably safeguard food security...

Meeting the demands for agricultural products from a growing and more affluent world population through conventional intensification of agriculture is impossible without causing significant damage to the environment [1-3]. Ecological intensification has been proposed as a nature-based alternative that complements or (partially) replaces external inputs such as agro-chemicals with production-supporting ecological processes to sustain agricultural production while minimizing adverse effects on the environment [4, 5]. Ecological intensification is based on the assumption that delivery of ecosystem services is suboptimal in high-input agricultural systems (e.g. [6-10]), and that management of specific components of biodiversity can be used to either complement artificial inputs and increase agricultural productivity (Ecological Enhancement; Fig. 1) or replace artificial inputs (Ecological Replacement; Fig. 1) which results in reduced environmental costs without negatively impacting crop productivity [5].

The last few years, the evidence base underlying ecological intensification has steadily strengthened with studies demonstrating that management can enhance the delivery of a range of regulating and supporting ecosystem services [11-14] or even produce win-win situations for agricultural production and the environment [15-18]. Scientists are therefore increasingly highlighting the benefits of ecologically intensifying agriculture through a greater reliance on biodiversity and ecosystem services. Policy makers likewise are starting to embrace ecological

intensification as an environmentally friendly way towards food security [19, 20] by supporting the implementation of biodiversity and ecosystem service enhancing practices. In some regions, notably Europe and North America, this has been through considerable public expenditure (e.g. agri-environment schemes) to (partially) offset farmer's opportunity costs associated with implementation [21].

...but sees little uptake by the agricultural sector

Knowledge of how farmers perceive the costs and benefits of ecological intensification practices is limited [22] but European farmers generally seem to have little interest in the topic. A recent survey on farmer attitudes towards biodiversity and ecosystem service enhancing practices in seven European countries [23] showed that farmers generally favour practices that interfere little with normal farming operations. For example, farmers appreciate relatively simple management changes targeting landscape features such as hedgerows, ditch banks or trees (Fig. 2). However, on-field management practices, such as cover crops, conservation headlands or beetle banks, were consistently among the least preferred practices (Fig. 2). Strikingly, the establishment of wildflower strips, the practice with the strongest evidence base for agronomic and/or economic benefits [12, 16, 24], and often eligible to subsidy support, is amongst the most disliked practices by farmers. Understanding why these practices are poorly adopted may explain why ecological intensification has seen little uptake to date by farmers, farmer organizations as well as agricultural corporations [19, 25, 26].

Here we explore why the perceptions of the costs and benefits of ecological intensification differ between scientists and farmers. We first synthesise the scientific evidence for nature-based contributions to agricultural production that underlie ecological intensification, and reflect on its relevance for farming enterprises. We consider both aboveground and belowground ecosystem services as both are relevant to farming, and ecological intensification has a greater potential of delivering benefits when targeting the full range of production-enhancing ecosystem services. We then highlight key knowledge gaps and suggest ways to overcome these. Finally we discuss the role of scientific evidence in shaping farm management and which additional factors are important drivers of farmer behaviour. Our focus is on ecosystem service enhancing practices rather than on farming systems (e.g. organic farming) and is mainly on high-input farming

systems since this is where biodiversity and ecosystem services are most degraded and where enhancing such services can potentially have the most pronounced effects [27].

Evidence for benefits of aboveground ecosystem services contributing to agricultural production

The species providing the two key aboveground ecosystem services relevant to agriculture, pollination and pest regulation, are mostly mobile organisms such as bees, hover flies, parasitoid wasps, spiders and carabid beetles. Although agricultural fields offer them important forage and shelter resources, these often come in short-lived fluxes, and beneficial species are generally highly dependent on semi-natural habitats in the surrounding landscape [28, 29]. Delivery of ecosystem services is therefore often inferred from the spatial configuration of landscape elements [30-32] with increasing landscape complexity (e.g. cover of semi-natural habitats, percentage non-arable land, distance to nearest semi-natural habitat, presence of wildflower strips) leading to higher pollination or pest regulation services. A wealth of studies have examined the relationship between the diversity and abundance of service providing species and landscape complexity and, on average, find positive relationships (Fig. 3) [29, 33-36]. However, there are notable exceptions, for example because pollinators do not always relate positively to landscape complexity [37, 38]. Also, natural enemies of crop pests are a taxonomically varied group of organisms that not necessarily all depend on semi-natural habitats and that may even be negatively related to cover of semi-natural habitats [39] (Fig. 3). Moreover, landscape complexity can also be related to delivery of dis-services, in the form of pests, but this relationship is highly variable and unresolved [36].

The relationship between landscape complexity and the diversity of service providing arthropods has led many scientists to conclude that delivery of ecosystem services can be influenced by maintaining or enhancing landscape complexity [40-42]. However, the relationship between landscape complexity and the actual delivery of the pollination and pest regulation services is less pronounced and more variable than that between the service providing taxa and landscape complexity [14, 33, 43-46]. Furthermore, the relationship between landscape complexity and crop yield, the main variable the agricultural sector is interested in, is even weaker and often absent [13, 41, 47-51]. The difference in focus on the main response variable may well contribute to the

122 difference in perceptions by scientists and farmers of the ecosystem service benefits that can be
123 obtained by manipulating landscape complexity (Fig. 3).

124 To date, only a few studies have convincingly demonstrated that management enhancing
125 pollination and pest regulation produces net agronomic or economic benefits. These studies have
126 in common that they examine the effects of establishing vegetation or wildflower strips on or
127 next to arable fields. Such measures invariably boost densities of pollinators and natural enemies
128 locally [52, 53] and can enhance crop pollination and pest regulation [54, 55] as well as a number
129 of other ecosystem services (e.g. reduce water runoff, increase soil and phosphorus retention [56].
130 However, only two of these studies suggest that yield increases were sufficient to compensate for
131 the opportunity costs (i.e. loss of cropped area) of establishing these new landscape elements [12,
132 24]. Only one study shows that yield increases were larger than both establishment and
133 opportunity costs so that farmers benefit economically from enhancing flower-rich habitats for
134 pollinators [16]. Further studies, across a range of crops and localities, are desperately needed.
135 With increasing demands for agricultural products and tight economic margins, farmers may
136 require more than just a proof of concept provide before they risk adopting ecological
137 intensification as a viable alternative or complementary approach to external input-based
138 practices.

140 **Evidence for benefits of belowground ecosystem services contributing agricultural** 141 **production**

142 The belowground communities of agricultural fields provide important ecosystem services such
143 as enhancing nutrient availability, prevention of pests and diseases, carbon storage and
144 improvement of soil structure and water holding capacity [57]. Soils contain a wealth of
145 biodiversity of microbes, invertebrates and some vertebrates, which can add up to thousands of
146 species per square metre of soil surface [58]. Recent studies suggest that soil biodiversity can be
147 engineered to specifically enhance the beneficial soil biota providing multiple ecosystem services
148 [59, 60]. In addition to the engineering approaches that often focus on introducing specific
149 organisms, such as for nutrient provision or plant protection, a more holistic approach has shown
150 how the stability of soil food webs depends on its structure [61]. Whereas individual groups of
151 soil biota correlate with specific ecosystem services [62], the connectedness of the entire soil
152 community corresponds with, for example, increased efficiency of carbon uptake by soil [63].

Organic matter may promote belowground biodiversity and ecosystem processes, and can even influence aboveground-belowground interactions by for example enhancing aboveground abundance of natural enemies [64]. Worldwide agriculture is causing loss of organic matter, except in areas with intensive animal farming [65] and in certain no-till conditions [66]. The question is how ecological intensification can make use of these novel insights into the relationship between soil biodiversity and functioning to improve crop production.

Key on-field practices that can improve the delivery of agriculturally relevant belowground ecosystem services are conservation tillage, the use of cover crops, increasing the diversity of the number of crops in a rotation or mixed cropping [60]. Figure 4 synthesises the impact of these practices and suggests that on average, and across all examined services, they have considerable positive effects. However, Figure 4 also highlights that none of the practices consistently enhance all of the ecosystem services considered here. For example, conservation tillage invariably reduces soil erosion and saves farmers tilling costs but has less consistent positive effects on soil structure, water retention and biodiversity [8, 67, 68], and has overall negative effects on nutrient retention, greenhouse gas emissions and weed control [14, 68]. The use of cover crops consistently improves soil structure and reduces soil erosion, however, it has less consistent positive effects on weed control and biodiversity [10, 20, 69]. Cover crops may improve nutrient retention and greenhouse gas emissions depending on whether nitrogen fixing cover crops are being used [9, 17]. However, in arid systems competition for water with the main crop generally results in yield reductions. Moreover, cover cropping requires additional sowing and sometimes killing the crop before planting the main crop which may bring substantial costs and the use of herbicides. Mixed cropping, or having a more diverse crop rotation, on average positively affects ecosystem service delivery [15, 46, 70-73], but for mixed cropping key information on the costs is still missing. Whether this is considered convincing evidence to a farmer may depend on which services are enhanced and which are reduced and probably what that means to the farm economically. In tandem with above-ground services, a greater number of studies on below-ground services in different cropping systems control [14, 68]. The use of cover crops consistently improves soil structure and reduces soil erosion, however, it has less consistent positive effects on weed control and biodiversity [10, 20, 69]. Cover crops may improve nutrient retention and greenhouse gas emissions depending on whether nitrogen fixing cover crops are being used [9, 17]. However, in arid systems competition for water with the main crop generally

results in yield reductions. Moreover, cover cropping requires additional sowing and sometimes killing the crop before planting the main crop which may bring substantial costs and the use of herbicides. Mixed cropping, or having a more diverse crop rotation, on average positively affects ecosystem service delivery [15, 46, 70-73], but for mixed cropping key information on the costs is still missing. Whether this is considered convincing evidence to a farmer may depend on which services are enhanced and which are reduced and probably what that means to the farm economically. In tandem with above-ground services, a greater number of studies on below-ground services in different cropping systems and locations is needed to help build a robust evidence base to support changes in farmer practices.

Knowledge gaps in the evidence base of ecological intensification

To be more convincing to farmers, scientific studies on ecological intensification need to address the costs and benefits that are most relevant to farmers (see Outstanding Questions). In addition to measuring straight-forward yield variables, parameters such as quality, commercial grading and stability of yield should be quantified as they also determine production value in many crops. Potential costs of ecological intensification should be an integral component of research. These include direct costs (e.g. establishment and maintenance of wildflower strips [16] as well as opportunity costs (e.g. loss of crop production on land used to establish wildflower strips). Ideally this should be done under a range of scenario's to account for context-dependence of the costs and benefits. For example, land prices in the Netherlands are an order of magnitude higher than in the United States (in 2009 approximately €47,000 ha⁻¹ and €3,700 ha⁻¹ respectively, [74, 75] resulting in higher fixed (mortgage) costs, which necessitates greater financial benefits from ecosystem services to break even. The benefits of particular ecosystem services can also be variable over time, as is illustrated by the pest regulation services provided by bats to cotton production in the south-western U.S.A.. These benefits declined by 79% between 1990 and 2008 due to falling global cotton prices and the widespread adoption of genetically modified Bt cotton [76]. Furthermore, when multiple services are considered, the cost-benefit analysis is complicated because different ecosystem services are usually expressed in different units making it difficult to assess whether a decline in one service is compensated by an increase in another. Cost-benefit analyses should additionally distinguish between private benefits and public goods delivered by ecological intensification. Public goods, such as reduced

greenhouse gas emissions or wildlife conservation, can benefit society at large but represent little or no direct benefit to individual farmers. For example, non-nitrogen fixing cover crops clearly outperform nitrogen fixing cover crops in terms of reducing greenhouse gas emissions and nutrient leaching (Fig. 4). However, leguminous cover crops are preferred by many farmers because they can result in higher yields in follow crops [17].

A second set of knowledge gaps concern the limited spatio-temporal scope of the evidence for ecosystem service benefits that is currently available (see Outstanding Questions). To date, most studies examine service delivery in a single crop at the field level in one or two years only [11-13, 46, 47, 54, 55, 77]. Studies that consider the spatio-temporal dimensions most relevant to farmers are rare. The key issues that need to be addressed are, first, that the populations of service providing species often need to build up before measurable effects can be established resulting in a time lag between implementation of ecological intensification and manifestation of ecosystem service benefits. Such time lags [78] may range from two or more years for pollination [16] to one or several decades for soil services [79]. Especially in farming systems where economic margins are low, farmers may not be willing to invest in practices of which they don't know when they will reap the benefits. Second, there is little information on pollination and pest regulation benefits across the crop rotation in annual cropping systems. The benefits of ecological intensification generally improve with increased targeting of the specific species groups providing the bulk of the services to a particular crop [12]. However, annual farming systems often rotate crops on individual fields. Ecological intensification should produce benefits across all crops in the rotation to be attractive to farmers. Third, information is lacking on benefits from ecological intensification at the farm scale, arguably the most relevant scale from the perspective of a farmer. In many countries, farms do not consist of a contiguous block of land, and fields can be scattered throughout the landscape. Most of the species providing pollination or pest regulation services are mobile and can be influenced by semi-natural habitats or crops up to several km away from the target location [28, 37, 46, 55, 77]. Their foraging ranges therefore generally supersede the size of individual farms [80]. The net farm-level benefits of enhancing pollination or pest regulation are difficult to predict as they depend on the implementation of nature-based management on the focal farm, on neighbouring farms and on biodiversity supporting habitat on public land such as protected areas, roadside verges and railway embankments [81]. Finally, although ecological theory predicts that service delivery

becomes more stable with increasing biodiversity [82], this has only been empirically demonstrated in small scale studies using experimental plant communities [83]. Variability in the profitability of farms as a result of adverse effects of inclement weather conditions on crop growth and yield is of major concern to farmers. Evidence of improved yield stability could be a powerful argument to interest farmers in ecological intensification.

Can scientific evidence of the benefits of ecological intensification increase its uptake?

Studies of farmer behaviour consistently show that short-term economic benefits enhance the adoption of novel biodiversity enhancing practices [25, 84, 85]. However, proven benefits alone do not guarantee uptake of management practices [86]. For example, conservation tillage in wheat has met with large-scale adoption in south Asia due to a 15-16% cost saving, but has met with limited uptake in Mexico and Southern Africa despite evidence of higher and more stable yields both for maize and wheat [87]. Farmers may decide not to follow scientific evidence because they are unsure about the relevance of generic recommendations from scientific studies for their specific farms and conditions. For example, a farm may be located on a different soil type than the study or bad weather can change a crop's response to a management practice [88]. Apart from economic considerations, key decisions by farmers and land managers are based on previous experience, familiarity with technologies, interactions with peers and advisors, labour requirements and perceived risks [25, 89]. Currently, advice to farmers often comes from advisors or sales representatives from agro-chemical companies that may sell both seeds and pesticides, and have financial incentives to promote their products [90]. In contrast, advice on nature-based management coming from parties such as independent extension services, NGO's and scientists may not reach as many farmers as this is not always a well-resourced core part of their business. Furthermore, agro-chemical applications offer quick, highly visible, short-term solutions to perceived problems. Rate and method of application are readily available as label instructions or otherwise provided by the manufacturers and effects can be easily observed. Ecological intensification tends to offer longer-term solutions. However, it relies on complex networks of service providing communities and management has mostly indirect effects that are rarely clear-cut and easily observed. For example, the relationship between semi-natural habitat or wildflowers in the wider countryside and pest regulation or fruit set may not be obvious to a farmer. Even with clear evidence of the benefits, using ecosystem services requires more

knowledge and initiative from farmers than spraying pesticides or adding honey bees at recommended rates. For some farmers, this alone may be an argument not to adopt ecological intensification practices. Finally, there is a general lack of practical, on the ground information to help farmers adopt nature-based management practices. We still have very little information on where how much of what kind of measures should be implemented to achieve a certain effect. This is because the proof of concept for ecological intensification has only recently been established and the amount of research on the topic is still small compared to the long-term and wide-ranging research on conventional farming practices [91]. Even today, conventional farming still receives not only the majority of the governmental funding but also almost all of the research investments by the private sector [92].

Farmers may, however, also adopt functional biodiversity enhancing practices without clear evidence of economic benefits as human behaviour is not solely driven by economic or other rational considerations [93]. Public attitude, in particular, can have strong direct and indirect effects on uptake of nature-based management practices by farmers. Farmers with strong social motivations can be influenced directly as adoption of ecological intensification contribute to a desired more positive image of their own farm by society and their peers [94]. Indirectly, public attitude can influence management of a much wider range of farms. For example, concern of the general public, in many parts of Europe, about intensive farming practices such as the use of pesticides or genetically engineered crops [95, 96] contributed to the EU restriction on the use of neonicotinoid pesticides in 2013. Uptake of ecological intensification may also be influenced by conflicts of interests between farmer communities and agribusiness multinationals and governments [97]. Many agribusinesses aim to generate societal support for the implementation of industrial forms of agriculture in new territories by emphasizing aspects of efficiency, productivity, economies of size, trade liberalization, free markets, and the need to feed the world [98]. Especially in the southern hemisphere, social movements such as La Via Campesina counter this by emphasizing benefits of family-based diversified agro-ecological farming such as small-scale production of healthy, local food, good stewardship of the rural environment and cultural heritages and the peasant or family farm way of life [99]. Such agroecology movements are now also gaining interest in northern countries with more industrial farming systems [100].

Conclusions

Large scale adoption of ecological intensification requires a stronger evidence base than is currently available. To date most research has focused on the ecological mechanisms and processes underlying ecological intensification in specific cropping systems. More knowledge is needed particularly on the quantification of the costs and benefits of ecological intensification using variables that are relevant to farmers (e.g. crop yield and earnings at the farm level) and the effectiveness of different ecological intensification practices, on their own and in combination, over longer periods of time and in a range of crops, farming systems and locations. The results of studies that have been carried out so far suggest that in the majority of crops and under the current economic paradigm it will be difficult for ecological intensification to achieve higher revenues than under conventional intensification. However, this could change in the near future as the price of external inputs are expected to rise, and climate change will make production more variable.

We propose that there are three complementary pathways towards wide-scale adoption of ecological intensification: through market driven processes, regulatory instruments and through reputational concerns. Market driven adoption will occur if a greater reliance on ecosystem services produces direct and net economic benefits [16] in which case it may simply become part of farm business models. Large-scale adoption through regulatory instruments require political will to promote nature-based farm management, for example through compulsory practices to support functional biodiversity linked to payments or by taxing agro-chemical inputs to integrate the environmental costs associated with the use of pesticides and artificial fertilizers into their price. Making external inputs more expensive would make biodiversity-based alternatives more attractive economically. Reputational concerns will increase adoption, if a sufficiently large part of the general public is worried about adverse effects of industrial farming and intensive use of agro-chemicals. This may influence farmers directly to manage their farms in ways to promote functional biodiversity when they can do this without economic repercussions. Moreover, given the global nature of the food market, changes in consumption patterns towards more environmental-friendly products (e.g. organic food) can influence farming practices all over the world. Just as importantly, public concern can be a strong driver of the political will to promote ecological intensification directly or indirectly (i.e. the regulatory instruments pathway). Future research should therefore not only address ecological, agronomic, and economic aspects of ecological intensification but also the sociological aspects.

339
340 **Acknowledgements:** This paper has been written in the framework of the European Union (EU)
341 FP7 projects LIBERATION (grant 311781). The paper benefited from additional support from the
342 Netherlands Organization for Scientific Research to DK (NWO-ALW project 841.11.001) and
343 LAG (NWO travel grant 040.11.577).

344

References

1. Foley, J.A. et al. (2005) Global consequences of land use. *Science* 309 (5734), 570-574.
2. Tilman, D. et al. (2011) Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences of the United States of America* 108 (50), 20260-20264.
3. Lundgren, J.G. and Fergen, J.K. (2011) Enhancing predation of a subterranean insect pest: A conservation benefit of winter vegetation in agroecosystems. *Applied Soil Ecology* 51, 9-16.
4. Cassman, K.G. (1999) Ecological intensification of cereal production systems: Yield potential, soil quality, and precision agriculture. *Proceedings of the National Academy of Sciences of the United States of America* 96 (11), 5952-5959.
5. Bommarco, R. et al. (2013) Ecological intensification: harnessing ecosystem services for food security. *Trends in Ecology & Evolution* 28 (4), 230-238.
6. Kremen, C. et al. (2002) Crop pollination from native bees at risk from agricultural intensification. *Proceedings of the National Academy of Sciences of the United States of America* 99 (26), 16812-16816.
7. Jonsson, M. et al. (2012) Agricultural intensification drives landscape-context effects on host-parasitoid interactions in agroecosystems. *Journal of Applied Ecology* 49 (3), 706-714.
8. Ulen, B. et al. (2010) Soil tillage methods to control phosphorus loss and potential side-effects: a Scandinavian review. *Soil Use and Management* 26 (2), 94-107.
9. Poeplau, C. and Don, A. (2015) Carbon sequestration in agricultural soils via cultivation of cover crops - A meta-analysis. *Agriculture Ecosystems & Environment* 200, 33-41.
10. Venter, Z.S. et al. (2016) The impact of crop rotation on soil microbial diversity: A meta-analysis. *Pedobiologia* 59 (4), 215-223.
11. Tschumi, M. et al. (2016) Perennial, species-rich wildflower strips enhance pest control and crop yield. *Agriculture Ecosystems & Environment* 220, 97-103.
12. Tschumi, M. et al. (2015) High effectiveness of tailored flower strips in reducing pests and crop plant damage. *Proceedings of the Royal Society B-Biological Sciences* 282 (1814), 189-196.
13. Gras, P. et al. (2016) How ants, birds and bats affect crop yield along shade gradients in tropical cacao agroforestry. *Journal of Applied Ecology* 53 (3), 953-963.
14. Tamburini, G. et al. (2016) Conservation tillage mitigates the negative effect of landscape simplification on biological control. *Journal of Applied Ecology* 53 (1), 233-241.
15. Davis, A.S. et al. (2012) Increasing Cropping System Diversity Balances Productivity, Profitability and Environmental Health. *Plos One* 7 (10), 8.
16. Blaauw, B.R. and Isaacs, R. (2014) Flower plantings increase wild bee abundance and the pollination services provided to a pollination-dependent crop. *Journal of Applied Ecology* 51 (4), 890-898.
17. Valkama, E. et al. (2015) Meta-analysis of the effects of undersown catch crops on nitrogen leaching loss and grain yields in the Nordic countries. *Agriculture Ecosystems & Environment* 203, 93-101.
18. Quemada, M. et al. (2013) Meta-analysis of strategies to control nitrate leaching in irrigated agricultural systems and their effects on crop yield. *Agriculture Ecosystems & Environment* 174, 1-10.
19. IPBES (2016) The assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production.

- Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany.
20. Blanco-Canqui, H. et al. (2015) Cover Crops and Ecosystem Services: Insights from Studies in Temperate Soils. *Agronomy Journal* 107 (6), 2449-2474.
 21. Batary, P. et al. (2015) The role of agri-environment schemes in conservation and environmental management. *Conservation Biology* 29 (4), 1006-1016.
 22. Uyttenbroeck, R. et al. (2016) Pros and cons of flowers strips for farmers. A review. *Biotechnologie Agronomie Societe Et Environnement* 20, 225-235.
 23. Bailey, A.P., Degani, E., Leigh, S. & Zhang, M. (2015) Report on farmer's attitude towards on-site ecosystem services. Deliverable 5.1 of the Liberation project.
 24. Pywell, R.F. et al. (2015) Wildlife-friendly farming increases crop yield: evidence for ecological intensification. *Proceedings of the Royal Society B-Biological Sciences* 282 (1816), 8.
 25. Liebman, M. et al. (2016) Ecologically sustainable weed management: How do we get from proof-of-concept to adoption? *Ecological Applications* 26 (5), 1352-1369.
 26. Vanloqueren, G. and Baret, P.V. (2008) Why are ecological, low-input, multi-resistant wheat cultivars slow to develop commercially? A Belgian agricultural 'lock-in' case study. *Ecological Economics* 66 (2-3), 436-446.
 27. Kleijn, D. et al. (2011) Does conservation on farmland contribute to halting the biodiversity decline? *Trends in Ecology & Evolution* 26 (9), 474-481.
 28. Steffan-Dewenter, I. et al. (2002) Scale-dependent effects of landscape context on three pollinator guilds. *Ecology* 83 (5), 1421-1432.
 29. Shackelford, G. et al. (2013) Comparison of pollinators and natural enemies: a meta-analysis of landscape and local effects on abundance and richness in crops. *Biological Reviews* 88 (4), 1002-1021.
 30. Lonsdorf, E. et al. (2009) Modelling pollination services across agricultural landscapes. *Annals of Botany* 103 (9), 1589-1600.
 31. Jonsson, M. et al. (2014) Ecological production functions for biological control services in agricultural landscapes. *Methods in Ecology and Evolution* 5 (3), 243-252.
 32. Maes, J. et al. (2015) More green infrastructure is required to maintain ecosystem services under current trends in land-use change in Europe. *Landscape Ecology* 30 (3), 517-534.
 33. Garibaldi, L.A. et al. (2011) Stability of pollination services decreases with isolation from natural areas despite honey bee visits. *Ecology Letters* 14 (10), 1062-1072.
 34. Kennedy, C.M. et al. (2013) A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters* 16 (5), 584-599.
 35. Bianchi, F. et al. (2006) Sustainable pest regulation in agricultural landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B-Biological Sciences* 273 (1595), 1715-1727.
 36. Chaplin-Kramer, R. et al. (2011) A meta-analysis of crop pest and natural enemy response to landscape complexity. *Ecology Letters* 14 (9), 922-932.
 37. Holzschuh, A. et al. (2008) Agricultural landscapes with organic crops support higher pollinator diversity. *Oikos* 117 (3), 354-361.
 38. Kleijn, D. et al. (2015) Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications* 6.
 39. Jauker, F. et al. (2009) Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landscape Ecology* 24 (4), 547-555.

40. Holzschuh, A. et al. (2012) Landscapes with wild bee habitats enhance pollination, fruit set and yield of sweet cherry. *Biological Conservation* 153, 101-107.
41. Nayak, G.K. et al. (2015) Interactive effect of floral abundance and semi-natural habitats on pollinators in field beans (*Vicia faba*). *Agriculture Ecosystems & Environment* 199, 58-66.
42. Nicholson, C.C. et al. (2017) Farm and landscape factors interact to affect the supply of pollination services. *Agriculture Ecosystems & Environment* 250, 113-122.
43. Winfree, R. et al. (2007) Native bees provide insurance against ongoing honey bee losses. *Ecology Letters* 10 (11), 1105-1113.
44. Karp, D.S. et al. (2013) Forest bolsters bird abundance, pest control and coffee yield. *Ecology Letters* 16 (11), 1339-1347.
45. Tschamntke, T. et al. (2016) When natural habitat fails to enhance biological pest control - Five hypotheses. *Biological Conservation* 204, 449-458.
46. Rusch, A. et al. (2013) Flow and stability of natural pest control services depend on complexity and crop rotation at the landscape scale. *Journal of Applied Ecology* 50 (2), 345-354.
47. Bommarco, R. et al. (2012) Insect pollination enhances seed yield, quality, and market value in oilseed rape. *Oecologia* 169 (4), 1025-1032.
48. Mitchell, M.G.E. et al. (2014) Agricultural landscape structure affects arthropod diversity and arthropod-derived ecosystem services. *Agriculture Ecosystems & Environment* 192, 144-151.
49. Liere, H. et al. (2015) Trophic cascades in agricultural landscapes: indirect effects of landscape composition on crop yield. *Ecological Applications* 25 (3), 652-661.
50. Zou, Y. et al. (2017) Landscape effects on pollinator communities and pollination services in small-holder agroecosystems. *Agriculture Ecosystems & Environment* 246, 109-116.
51. Sutter, L. et al. Landscape greening and local creation of wildflower strips and hedgerows promote multiple ecosystem services. *Journal of Applied Ecology*, n/a-n/a.
52. Scheper, J. et al. (2013) Environmental factors driving the effectiveness of European agri-environmental measures in mitigating pollinator loss - a meta-analysis. *Ecology Letters* 16 (7), 912-920.
53. Ramsden, M.W. et al. (2015) Optimizing field margins for biocontrol services: The relative role of aphid abundance, annual floral resources, and overwinter habitat in enhancing aphid natural enemies. *Agriculture Ecosystems & Environment* 199, 94-104.
54. Feltham, H. et al. (2015) Experimental evidence that wildflower strips increase pollinator visits to crops. *Ecology and Evolution* 5 (16), 3523-3530.
55. Holland, J.M. et al. (2012) Agri-environment scheme enhancing ecosystem services: A demonstration of improved biological control in cereal crops. *Agriculture Ecosystems & Environment* 155, 147-152.
56. Schulte, L.A. et al. (2017) Prairie strips improve biodiversity and the delivery of multiple ecosystem services from corn-soybean croplands. *Proceedings of the National Academy of Sciences of the United States of America* 114 (42), 11247-11252.
57. Wall, D.H. et al. (2015) Soil biodiversity and human health. *Nature* 528 (7580), 69-76.
58. Bardgett, R.D. and van der Putten, W.H. (2014) Belowground biodiversity and ecosystem functioning. *Nature* 515 (7528), 505-511.
59. Wagg, C. et al. (2014) Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences of the United States of America* 111 (14), 5266-5270.
60. Bender, S.F. et al. (2016) An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. *Trends in Ecology & Evolution* 31 (6), 440-452.

61. Neutel, A.M. et al. (2007) Reconciling complexity with stability in naturally assembling food webs. *Nature* 449 (7162), 599-U11.
62. de Vries, F.T. et al. (2013) Soil food web properties explain ecosystem services across European land use systems. *Proceedings of the National Academy of Sciences of the United States of America* 110 (35), 14296-14301.
63. Morrien, E. et al. (2017) Soil networks become more connected and take up more carbon as nature restoration progresses. *Nature Communications* 8, 10.
64. Birkhofer, K. et al. (2008) Long-term organic farming fosters below and aboveground biota: Implications for soil quality, biological control and productivity. *Soil Biology & Biochemistry* 40 (9), 2297-2308.
65. Reijneveld, A. et al. (2009) Soil organic carbon contents of agricultural land in the Netherlands between 1984 and 2004. *Geoderma* 152 (3-4), 231-238.
66. Pittelkow, C.M. et al. (2015) Productivity limits and potentials of the principles of conservation agriculture. *Nature* 517 (7534), 365-U482.
67. Morris, N.L. et al. (2010) The adoption of non-inversion tillage systems in the United Kingdom and the agronomic impact on soil, crops and the environment-A review. *Soil & Tillage Research* 108 (1-2), 1-15.
68. Soane, B.D. et al. (2012) No-till in northern, western and south-western Europe: A review of problems and opportunities for crop production and the environment. *Soil & Tillage Research* 118, 66-87.
69. Tonitto, C. et al. (2006) Replacing bare fallows with cover crops in fertilizer-intensive cropping systems: A meta-analysis of crop yield and N dynamics. *Agriculture Ecosystems & Environment* 112 (1), 58-72.
70. Iverson, A.L. et al. (2014) Do polycultures promote win-wins or trade-offs in agricultural ecosystem services? A meta-analysis. *Journal of Applied Ecology* 51 (6), 1593-1602.
71. McDaniel, M.D. et al. (2014) Does agricultural crop diversity enhance soil microbial biomass and organic matter dynamics? A meta-analysis. *Ecological Applications* 24 (3), 560-570.
72. Yu, Y. et al. (2015) Temporal niche differentiation increases the land equivalent ratio of annual intercrops: A meta-analysis. *Field Crops Research* 184, 133-144.
73. Pappa, V.A. et al. (2011) Nitrous oxide emissions and nitrate leaching in an arable rotation resulting from the presence of an intercrop. *Agriculture Ecosystems & Environment* 141 (1-2), 153-161.
74. Eurostat (2015) Agricultural prices and price indices (EU and candidate countries). URL: <http://ec.europa.eu/eurostat/web/agriculture/data> (accessed: 11-11-2015).
75. USDA (2015) Land Values 2015 Summary. USDA, National Agricultural Statistics Service.
76. Lopez-Hoffman, L. et al. (2014) Market Forces and Technological Substitutes Cause Fluctuations in the Value of Bat Pest-Control Services for Cotton. *Plos One* 9 (2), 7.
77. Gardiner, M.M. et al. (2009) Landscape diversity enhances biological control of an introduced crop pest in the north-central USA. *Ecological Applications* 19 (1), 143-154.
78. Fremier, A.K. et al. (2013) Understanding Spatiotemporal Lags in Ecosystem Services to Improve Incentives. *Bioscience* 63 (6), 472-482.
79. Brady, M.V., Sahlin, U., Clough, Y., Bailey, A., Elek, Z., Hedlund, K., Koellner, T., Marini, L., Olsson, O., Poppenborg, P., Redlich, S., Świtek, S., Takacs, V., van Gils, S. & Smith, H.G. (2016) Report on ecological-economic models and evaluation of effects of ecological intensification on farm income. Deliverable 5.4 of the Liberation Project.

80. McKenzie, A.J. et al. (2013) Landscape-scale conservation: collaborative agri-environment schemes could benefit both biodiversity and ecosystem services, but will farmers be willing to participate? *Journal of Applied Ecology* 50 (5), 1274-1280.
81. Cong, R.G. et al. (2014) Managing ecosystem services for agriculture: Will landscape-scale management pay? *Ecological Economics* 99, 53-62.
82. Yachi, S. and Loreau, M. (1999) Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis. *Proceedings of the National Academy of Sciences of the United States of America* 96 (4), 1463-1468.
83. Isbell, F. et al. (2015) Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature* 526 (7574), 574-U263.
84. Willock, J. et al. (1999) Farmers' attitudes, objectives, behaviors, and personality traits: The Edinburgh Study of Decision Making on Farms. *Journal of Vocational Behavior* 54 (1), 5-36.
85. van der Horst, D. (2011) Adoption of payments for ecosystem services: An application of the Hagerstrand model. *Applied Geography* 31 (2), 668-676.
86. Garibaldi, L.A. et al. (2017) Farming Approaches for Greater Biodiversity, Livelihoods, and Food Security. *Trends in Ecology & Evolution* 32 (1), 68-80.
87. Erenstein, O. et al. (2012) Conservation Agriculture in Maize- and Wheat-Based Systems in the (Sub)tropics: Lessons from Adaptation Initiatives in South Asia, Mexico, and Southern Africa. *Journal of Sustainable Agriculture* 36 (1-2), 180-206.
88. Sheriff, G. (2005) Efficient waste? Why farmers over-apply nutrients and the implications for policy design. *Review of Agricultural Economics* 27 (4), 542-557.
89. Martinez-Garcia, C.G. et al. (2013) Factors influencing adoption of improved grassland management by small-scale dairy farmers in central Mexico and the implications for future research on smallholder adoption in developing countries. *Livestock Science* 152 (2-3), 228-238.
90. Wilson, C. and Tisdell, C. (2001) Why farmers continue to use pesticides despite environmental, health and sustainability costs. *Ecological Economics* 39 (3), 449-462.
91. Kremen, C. and Miles, A. (2012) Ecosystem Services in Biologically Diversified versus Conventional Farming Systems: Benefits, Externalities, and Trade-Offs. *Ecology and Society* 17 (4), 25.
92. Tittonell, P. (2014) Ecological intensification of agriculture - sustainable by nature. *Current Opinion in Environmental Sustainability* 8, 53-61.
93. Cerasoli, C.P. et al. (2014) Intrinsic Motivation and Extrinsic Incentives Jointly Predict Performance: A 40-Year Meta-Analysis. *Psychological Bulletin* 140 (4), 980-1008.
94. Greiner, R. et al. (2009) Motivations, risk perceptions and adoption of conservation practices by farmers. *Agricultural Systems* 99 (2-3), 86-104.
95. Madsen, K.H. and Sandoe, P. (2001) Herbicide resistant sugar beet - what is the problem? *Journal of Agricultural & Environmental Ethics* 14 (2), 161-168.
96. Wollaege, H.M. et al. (2015) Consumer Preferences for Traditional, Neonicotinoid-free, Bee-friendly, or Biological Control Pest Management Practices on Floriculture Crops. *Hortscience* 50 (5), 721-732.
97. Rosset, P.M. and Martinez-Torres, M.E. (2012) Rural Social Movements and Agroecology: Context, Theory, and Process. *Ecology and Society* 17 (3).
98. Rosset, P.M. and Martinez-Torres, M.E. (2017) La Via Campesina's Open Book: Celebrating 20 Years of Struggle and Hope. URL: <https://viacampesina.org/downloads/pdf/openbooks/EN-12.pdf> (accessed: 21-1-2017).

99. Borras, S.M. et al. (2008) Transnational agrarian movements: Origins and politics, campaigns and impact. *Journal of Agrarian Change* 8 (2-3), 169-204.
100. Tomlinson, I. (2013) Doubling food production to feed the 9 billion: A critical perspective on a key discourse of food security in the UK. *Journal of Rural Studies* 29, 81-90.
101. Holland, J.M. (2004) The environmental consequences of adopting conservation tillage in Europe: reviewing the evidence. *Agriculture Ecosystems & Environment* 103 (1), 1-25.
102. Dabney, S.M. et al. (2001) Using winter cover crops to improve soil and water quality. *Communications in Soil Science and Plant Analysis* 32 (7-8), 1221-1250.
103. Tiemann, L.K. et al. (2015) Crop rotational diversity enhances belowground communities and functions in an agroecosystem. *Ecology Letters* 18 (8), 761-771.
104. Rusch, A. et al. (2013) Effect of crop management and landscape context on insect pest populations and crop damage. *Agriculture Ecosystems & Environment* 166, 118-125.
105. Palmu, E. et al. (2014) Landscape-scale crop diversity interacts with local management to determine ground beetle diversity. *Basic and Applied Ecology* 15 (3), 241-249.
106. Jeuffroy, M.H. et al. (2013) Nitrous oxide emissions from crop rotations including wheat, oilseed rape and dry peas. *Biogeosciences* 10 (3), 1787-1797.
107. Karlen, D.L. et al. (2006) Crop rotation effects on soil quality at three northern corn/soybean belt locations. *Agronomy Journal* 98 (3), 484-495.
108. Dassou, A.G. and Tixier, P. (2016) Response of pest control by generalist predators to local-scale plant diversity: a meta-analysis. *Ecology and Evolution* 6 (4), 1143-1153.
109. Letourneau, D.K. et al. (2011) Does plant diversity benefit agroecosystems? A synthetic review. *Ecological Applications* 21 (1), 9-21.
110. Liebman, M. and Dyck, E. (1993) Crop-rotation and intercropping strategies for weed management. *Ecological Applications* 3 (1), 92-122.
111. Zhou, X.G. et al. (2011) Effects of intercropping cucumber with onion or garlic on soil enzyme activities, microbial communities and cucumber yield. *European Journal of Soil Biology* 47 (5), 279-287.
112. Lacombe, S. et al. (2009) Do tree-based intercropping systems increase the diversity and stability of soil microbial communities? *Agriculture Ecosystems & Environment* 131 (1-2), 25-31.

Glossary

Biodiversity: the variety of all forms of life, from genes to species and ecosystems. Within the context of ecological intensification.

Conservation tillage: the practice of reducing tillage intensity and retaining crop residues to conserve soil, water and energy.

Cover crop: crop grown between two cash crops to suppress weeds, improve soil fertility and reduce pest pressure and that is generally not harvested.

Crop rotation: the practice of growing different crops in succession on the same land to maintain soil productivity and control weeds, pests, and diseases.

Ecosystem service: benefits obtained by people from ecosystems.

External inputs: non-renewable or industrially made resources, such as fertilizers or pesticides used by growers to increase yield or avoid yield loss.

Functional biodiversity: the part of all biodiversity that makes a direct contribution to agricultural production.

Habitat quality: the extent to which a habitat offers all the resources required by species to successfully complete their life cycle.

High-input farming systems: Farming systems in which crop production is primarily based on external inputs such as fuel, fertilisers and pesticides.

Landscape complexity: the extent to which a landscape is covered by a variety of semi-natural, non-crop habitats.

Natural enemies: the naturally occurring predators and parasitoids of crop pests.

Mixed cropping: the practice of growing multiple crops simultaneously in the same field to enhance overall yield and reduce pressure of pests, weeds and diseases.

Pest regulation service: control of herbivore pests of (crop) plants by wild predators such as beetles, spiders, parasitoid wasps and birds.

Pollination service: pollination of crop and wild plants by wild pollinators such as bees, hoverflies and bats.

Outstanding questions

Response variables considered

- What are the effects of ecological intensification on parameters relevant to farmers?
- What are the (opportunity) costs of ecological intensification and are they balanced by the benefits?
- Are there synergies or trade-offs between delivery of multiple ecosystem services?
- Does ecological intensification have different effects on delivery of private benefits and public goods?

Spatio-temporal scales considered

- How long are time-lags between implementing management and delivery of benefits?
- What are the pollination and pest regulation benefits across the full rotation of annual crops?
- What are the farm-scale costs and benefits of ecological intensification?
- Does ecological intensification reduce yield variability?
- How can ecological intensification best be implemented practically (e.g. how much, where, when)?

Highlights

Ecological intensification aims to harness ecosystem services to sustain agricultural production while minimizing adverse effects on the environment.

Ecological intensification is championed by scientists as a nature-based alternative to high-input agriculture but meets with little interest from growers.

Scientific evidence underlying ecological intensification is often unconvincing to growers as it is based on small-scale studies of ecological processes.

Grower interest can be enhanced by evidence of the agronomic and economic benefits most relevant to farmers and measured at the scales of operation of farm enterprises.

In addition to concrete benefits, concerns of the general public about adverse effects of industrial farming can promote adoption of ecological intensification, both directly and indirectly by enhancing political will to use regulatory instruments.

Figure 1




Figure 1. An illustration of the relationships assumed under ecological intensification in high-input farming systems between functional biodiversity and (a) crop yield and (b) dependence on synthetic inputs (pesticides and artificial fertilizers) under ecological replacement and ecological enhancement.

Figure 2

1
2
3
4
5
6
7
8
9
10




Figure 2. Preferences of farmers from Germany, Hungary, Italy, Netherlands, Poland, Sweden and UK for management practices that may contribute to biodiversity enhancement and ecosystem service. Number of responses per management practice ranged between 55 and 84. Based on data from [23].

Figure 3




Figure 4




Figure 4. Radar plots graphically summarising the effects of the most frequently implemented management practices to increase sustainability of farming on multiple ecosystem services. Dark green/red - consistent positive/negative effects found in meta-analyses, reviews and individual studies; Intermediate green/red: positive/negative effects dominate but some studies show no effects; Light green/red: positive/negative effects dominate but many studies show no effect and some even negative effects. Effects based on refs [14, 67, 68, 101] for conservation tillage; [3, 9, 10, 17, 18, 20, 69, 102, 103] for cover crops; [10, 15, 71, 103-107] for crop rotation; [72, 73, 108-112] for mixed cropping.