

The impact of increased flooding occurrence on the mobility of potentially toxic elements in floodplain soil – a review

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Ponting, J., Kelly, T. J., Verhoef, A. ORCID: <https://orcid.org/0000-0002-9498-6696>, Watts, M. J. and Sizmur, T. ORCID: <https://orcid.org/0000-0001-9835-7195> (2021) The impact of increased flooding occurrence on the mobility of potentially toxic elements in floodplain soil – a review. Science of the Total Environment, 754. 142040. ISSN 0048-9697 doi: 10.1016/j.scitotenv.2020.142040 Available at <https://centaur.reading.ac.uk/92613/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.scitotenv.2020.142040>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1

2 The impact of increased flooding occurrence on the
3 mobility of potentially toxic elements in floodplain soil
4 – A review

5

6 Jessica Ponting ^{a,b}, Thomas J. Kelly ^{a,c}, Anne Verhoef ^a, Michael J. Watts ^b and
7 Tom Sizmur ^{a*}

8

9 ^a Department of Geography and Environmental Science, University of Reading, Reading, RG6
10 6DW, UK

11 ^b Inorganic Geochemistry, Centre for Environmental Geochemistry, British Geological
12 Survey, Keyworth, Nottingham, NG12 5GG, UK

13 ^c Department of Geography, Queen Mary University of London, Mile End Road, London, E1
14 4NS, UK

15

16 * Corresponding author

17 Department of Geography and Environmental Science
18 Russell Building
19 University of Reading
20 Reading
21 RG6 6DW, UK

22 t.sizmur@reading.ac.uk

23 +44(0)118 3788913

24

25

26

27

Abstract

The frequency and duration of flooding events are increasing due to land-use changes increasing runoff of precipitation, and climate change causing more intense rainfall events. Floodplain soils situated downstream of urban or industrial catchments, which were traditionally considered a sink of potentially toxic elements (PTEs) arriving from the river reach, may now become a source of legacy pollution to the surrounding environment if PTEs are mobilised by unprecedented flooding events.

When a soil floods, the mobility of PTEs can increase or decrease due to the net effect of five key processes; (i) the soil redox potential decreases which can directly alter the speciation, and hence mobility, of redox sensitive PTEs (e.g. Cr, As), (ii) pH increases which usually decreases the mobility of metal cations (e.g. Cd^{2+} , Cu^{2+} , Ni^{2+} , Pb^{2+} , Zn^{2+}), (iii) dissolved organic matter (DOM) increases, which chelates and mobilises PTEs, (iv) Fe and Mn hydroxides undergo reductive dissolution, releasing adsorbed and co-precipitated PTEs, and (v) sulphate is reduced and PTEs are immobilised due to precipitation of metal sulphides. These factors may be independent mechanisms, but they interact with one another to affect the mobility of PTEs, meaning the effect of flooding on PTE mobility is not easy to predict. Many of the processes involved in mobilising PTEs are microbially mediated, temperature dependent and the kinetics are poorly understood.

Soil mineralogy and texture are properties that change spatially and will affect how the mobility of PTEs in a specific soil may be impacted by flooding. As a result, knowledge based on one river catchment may not be particularly useful for predicting the impacts of flooding at another site. This review provides a critical discussion of the mechanisms controlling the mobility of PTEs in floodplain soils. It summarises current understanding, identifies limitations to existing knowledge, and highlights requirements for further research.

Key words; floodplain soil, flooding, climate change, potentially toxic elements, contamination, mobility

1. Introduction

Flooding is a major event that currently affects an estimated 20 to 300 million people per year, and accounts for around 40% of natural disasters occurring worldwide, threatening both social security and sustainable development (Euripidou and Murray, 2004; Hirabayashi and Kanae, 2009). Alterations to land use and land cover are having widespread implications for catchment characteristics; with soil sealing and impermeable surfaces increasing surface run-off, as well as a reduction of natural buffering environments such as forests and wetlands, meaning there is less capacity to accommodate flood waters in the same river reach (Dadson et al., 2017; Kundzewicz et al., 2014). There is growing evidence, from climate models, that short-term extreme weather events (e.g. high-frequency rainstorms, heat waves and wind storms) are likely to become increasingly frequent in many parts of the world, threatening the long-term functioning of the terrestrial system (Harvey et al., 2019; Kharin et al., 2007; Madsen et al., 2014; Pendergrass, 2018; Stagl et al., 2014). It is likely that populations will experience warmer and drier summers, and an increase in the intensity of heavy rainfall, contributing to more frequent pluvial, fluvial, groundwater or coastal flooding, and resulting in the occasional inundation of land that has rarely been flooded in the past (Barber et al., 2017; Kundzewicz et al., 2014; Schaller et al., 2016). The likelihood of flooding is also determined by antecedent soil moisture conditions. The proportion of soil pore space that is filled with water at any given time is largely dependent on local hydrological processes and stores including; infiltration, surface and sub-surface runoff (when rainfall intensity exceeds infiltration capacity), redistribution and drainage to/from groundwater, evaporation, and transpiration (Stagl et al., 2014).

Soil contamination is among the most serious threats to soil resources globally (Nriagu et al., 2007; Srivastava et al., 2017; Tóth et al., 2016b). Since many commercial, industrial, residential and agricultural developments have historically been situated adjacent to rivers; they contribute to the contamination of river sediments, and these sediments are often deposited onto the floodplain soils downstream by overbanking river water during a flooding event (Arnell et al., 2015; Nshimiyimana et

al., 2014; Zhao et al., 1999). Here we use the term PTEs, also referred to in the literature as ‘trace elements’ or ‘heavy metals’, to encompass all metals, metalloids, non-metals and other inorganic elements in the soil–plant–animal system, of which their mobility and potential toxicity to that system and/or humans is largely dependent upon their concentration, bioavailability and chemical form (Hooda, 2010; Rodgers et al., 2015). The term "mobility" is a concept that has been frequently used to estimate the risk of contamination from the soil to the surrounding environment by PTEs (Domergue and Vedy, 1992). Here we define mobile PTEs as those elements that are dissolved in soil porewater or associated with colloids and thus capable of leaching from the soil profile, or being taken up into plants or soil organisms. The mobility and subsequent fate of PTEs in periodically (occasionally) flooded soils (such as floodplain soils) are imperfectly understood. The legacy of historic contamination and continuing increases in emissions from urban activities pose a serious environmental threat globally (de Souza Machado et al., 2016; Srivastava et al., 2017). Human actions to mitigate and adapt to the impacts of climate change may influence the fate of contaminants, with climate change itself also potentially affecting the toxicity of the contaminants within the environment (Stahl et al., 2013).

The aim of this literature review is to provide an understanding of the factors involved in the mobility of PTEs in soil by pulling together interdisciplinary knowledge in this area. The review will first consider in more detail the expected changes to global rainfall patterns, the implications of these changes for flooding, and the role that floodplains play during inundation, as well as the changes they undergo. The review will then showcase how PTEs have entered the floodplain soil and how flooding influences soil biogeochemical processes which, in turn, influence PTEs mobility, using examples from the literature. Finally, this knowledge is used to identify gaps that will help to make recommendations for future research into the effects of flooding on the mobility and fate of PTEs.

1.1. Expected changes to global rainfall patterns and implications for flooding

Anthropogenic (human) activities including intensified land use; urbanisation, forestry, cultivation, and fossil energy use have increased atmospheric greenhouse gas concentrations which are driving changes in climate and leading to increases in rainfall intensity and surface run-off that are associated with increased flood risk (Bronstert, 2003; Chang and Franczyk, 2008; Kharin et al., 2007; Kundzewicz et al., 2014; Wheeler and Evans, 2009). Mean global temperatures have risen by 1.1 °C since the end of the 19th century; the “Paris Climate Agreement” seeks to contain global mean temperatures well below 2°C and, ambitiously, below 1.5°C (Alfieri et al., 2017; Bronstert, 2003; Huddart et al., 2020; Mullan et al., 2019). The Intergovernmental Panel on Climate Change (IPCC) has predicted that under the A1B (medium) emissions scenario, temperatures will increase between 1.1 and 6.4 °C by the year 2100, leading to an increase in atmospheric water holding capacity and therefore variations to seasonal rainfall (Arnell et al., 2015; Bell et al., 2012; Chan et al., 2014; Clemente et al., 2008; González-Alcaraz and van Gestel, 2015; Jenkins et al., 2009). It has been argued that we will experience an intensification of short-duration heavy rainfall events rather than a uniform increase in the daily average rainfall (Chan et al., 2014; Hirabayashi et al., 2008; Kharin et al., 2007; Kundzewicz et al., 2014).

An IPCC Special Report (SREX) on climate extremes (IPCC, 2012) assessed it is *likely* there have been statistically significant increases in the number of heavy precipitation events in more regions than significant decreases, with strong regional and sub-regional variation. The observed changes to precipitation extremes have been found to be far less spatially coherent or statistically significant compared with changes found in temperature extremes (Kundzewicz et al., 2014). Projected scenarios with 4°C warming showed more than 70% of the global population will face increased flood risk (Alfieri et al., 2017). Increases in flood frequency are expected in; Europe, America, Southeast Asia, eastern Africa, and Peninsular India. Populations in regions such as Bangladesh, Mumbai and Thailand are potentially at higher risk from flooding due to predicted increases in rainfall, coupled with changes in land use (e.g. irrigation schemes and construction of dams), and increasing population size requiring

rapid expansion of urban areas (Hirabayashi et al., 2013; Kundzewicz et al., 2014; Tockner et al., 2010). However, climate projections tend to have relatively low levels of model confidence, particularly for the prediction of fluvial floods because there is still relatively limited evidence and the causes of regional changes to flood occurrence are complex (Hirabayashi et al., 2013).

Flooding tends to be heterogeneous as it is affected not only by variability of the climatological and hydrological systems but also by land-use and the effect it has on the storage capacity of the receiving catchment (storage and drainage basin conditions). Changes to the characteristics of precipitation (the frequency, intensity and timing of rainfall) will have decisive implications for flood risk (Bronstert, 2003; Hirabayashi and Kanae, 2009; Kundzewicz et al., 2014). However, pre-existing high river levels and groundwater levels, as well as saturated soils are equally important to establish the capacity of the receiving catchment to cope with further rainfall (Maggioni and Massari, 2018; Wilby et al., 2008). The extent of flooding in a particular catchment will depend largely on the topography (variation in elevation), along with vegetation type, proportion of land used for cultivation and the extent of urbanised areas positioned upstream (Arnell et al., 2015; Bell et al., 2012; Bronstert, 2003; Chang and Franczyk, 2008; Kundzewicz et al., 2014; Qiao et al., 2019). Urbanisation is a global issue; with more than half the world's population now living in cities, the process of urbanisation is leading to greater human occupation of floodplains, often with inadequate drainage planning (Kundzewicz et al., 2014; Pathirana et al., 2014).

The probability of flooding occurring in a particular region is often related to regional processes like El Niño Southern Oscillation (ENSO) cycle and the North Atlantic Oscillation (NAO) that, in turn, cause global impacts. The intensity (frequency and amplitude) of both ENSO and NAO are influenced by other modes of variability, for example; Pacific Decadal Oscillation (PDO) and Interdecadal Pacific Oscillation (IPO) which cause opposite atmospheric and sea surface temperatures and can therefore determine the magnitude of floods (Grimm and Tedeschi, 2009; Johnson et al., 2020). ENSO is a rapid warming of the sea surface temperature (by 1–5 °C) of the equatorial Pacific over the duration of a

few weeks, resulting in extreme rainfall and increased cyclone activity in some regions, and risk of drought and forest fires in others (Berz et al., 2001; Grimm and Tedeschi, 2009; Karl and Trenberth, 2003; Kundzewicz et al., 2014; Tedeschi and Collins, 2016). Periods of extreme rainfall and subsequent flooding have been found to correlate with ENSO events in North and South America as well as in Africa (Berz et al., 2001; Brönnimann, 2007; Kundzewicz et al., 2014). NAO is an atmospheric pattern that affects the severity of winter temperatures and precipitation over Europe and eastern North America (Karl and Trenberth, 2003). Intense rainfall is a common cause of river basin flooding; however, in high latitude regions it is changes in temperatures altering the timing of seasonal snowmelt and causing glacier retreat that commonly causes flooding, for example in north-eastern Europe, Central and South America, and in polar regions such as the Russian Arctic (Blöschl et al., 2017; Hirabayashi et al., 2008; Kharin et al., 2007; Kundzewicz et al., 2014; Stagl et al., 2014). Rising global sea-level (11-16cm in the 20th century and a further 0.5m predicted this century) will certainly increase risk of flooding caused by tidal processes, with current estimates that 630 million people live on land below projected annual flood levels for 2100 (Kulp and Strauss, 2019). While there is uncertainty regarding the effect that future climate change will have on river levels (Prudhomme and Davies, 2009), changes made to land-use, and land cover, for example by urbanisation, will drive changes in the local climate (at the kilometre scale) influencing the hydrometeorological regime and resulting in more flooding (Foley et al., 2005; Hirabayashi and Kanae, 2009). Pathirana et al. (2014), using a 3D atmospheric model coupled with a land surface model (WRF-ARW) in southern India, found that in three out of four simulated cases there was a significant increase in local extreme rainfall when urbanisation in the area increased. This work was conducted in southern India, however the model could be applied and validated to other regions to establish whether this correlation is found globally.

1.2. The role of floodplains during floods

Floodplains are by definition dynamic environments subjected to fluctuations between flooding and drying (Vijver et al., 2007). They are distinctive landscape features, often on low-lying ground, and

characterised by a high spatio-temporal heterogeneity (Schulz-Zunkel et al., 2015; Stuart and Lapworth, 2011; Tockner et al., 2010; Tockner and Stanford, 2002). Periodic overbank inundation from the adjacent watercourse, overland flow, subsurface flow, and changes to the groundwater levels result in a constantly changing water balance and degree of floodplain saturation (Stuart and Lapworth, 2011; Tockner and Stanford, 2002). Floodplain topography and variations in elevation are usually slight but have an important effect on the degree of soil saturation across the floodplain, depending on the overall water balance from surface and sub-surface run-off (Arnell et al., 2015; Kundzewicz et al., 2014; Qiao et al., 2019).

There are various sources and pathways of water that can lead to the inundation of a floodplain, including lateral overflow of rivers or lakes, rising groundwater, upland sources, and direct precipitation. Several different factors and water sources normally contribute to a flooding event, thus making flooding a complex phenomenon to study (Junk et al., 1989; Tockner and Stanford, 2002). Fluvial flooding tends to occur when excessive rain falls over an extended period of time, leading to a river exceeding its capacity, or because of heavy snow that subsequently melts and, via surface run-off, rapidly fills the river channels when infiltration is low because of frozen soils below the snow layer (Blöschl et al., 2017).

River flow regimes are affected by the increased rainfall and this also has the potential to affect erosion and generate additional sediment loads and particulate organic matter (POM) for deposition within river channels, lakes and estuaries (Arnell et al., 2015; Le Gall et al., 2018; Rinklebe and Du Laing, 2011). Intense rainfall over a short timescale (usually less than six hours i.e. “flash floods”) can also cause rivers to overbank leading to an intense, high velocity torrent of water that moves through river beds, disturbing river sediments and potentially bringing more PTEs contamination with the flood water, greatly influencing the contaminated status of the floodplain (Blöschl et al., 2017; Maggioni and Massari, 2018). The water inundating the floodplain contains dissolved matter (i.e. free ions, inorganic and organic complexes and uncharged molecules) as well as particulate matter (i.e.

large organic and inorganic polymers, oxides, clay minerals and organic matter) (Kirk, 2004). The sediment loads travel at different rates due to their particle size, which reflects the texture of the river bed and bank (Malmon et al., 2004). Approximately 90% of PTEs load has been associated with sediment particles, with dissolved PTEs playing a comparatively minor role in pollutant transfer to floodplains (Ciszewski and Grygar, 2016). There have been many fluvial geomorphology studies showing how erosion and sedimentation have been influenced by climatic variability in the past (e.g. Lewin and Macklin, 2010; Macklin and Rumsby, 2007; Mullan et al., 2019), indicating that rivers are sensitive to climatic change (Arnell et al., 2015). Fluvial flooding is receiving increased scientific and political interest because of the potential impact that climate change may have on this type of flooding, with climate model projections showing an increased flood risk at a global scale (Pappenberger et al., 2012; Wilby et al., 2008).

In floodplains that are underlain by permeable deposits, increased rainfall causes groundwater to rise (leading to groundwater flooding), which can result from direct rainfall recharge, when the soil water storage potential is exceeded, as well as flow into the floodplain sediments from rivers with high water levels, and from areas inundated with fluvial flooding. However, good hydraulic connection between river and aquifer means that the aquifer can drain quickly as fluvial flood waters recess. Groundwater flooding in these settings is relatively short-lived compared with other groundwater flood settings, for example in chalk catchments (MacDonald et al., 2012).

With increased frequency of rainfall events predicted, it has become widely recognised that the storage of floodwater on floodplains can help to reduce the magnitude of a flood downstream. Thus, floodplains are useful for flood risk management (Acreman et al., 2003; Vink and Meeussen, 2007). As a result, floodplains may be deliberately managed to allow flooding to occur through engineered soakaways in order to protect an urban residential area (Lane, 2017; Wheeler and Evans, 2009). It is important to understand the potential implications of these types of management practices

on mobilisation of PTEs that may be associated with the sediments deposited on the floodplain during past flooding events.

1.3. Changes that floodplain soils undergo during and after inundation

Extreme rainfall events leading to flooding have generally been found to alter soil physical and chemical properties and influence biological processes (Harvey et al., 2019). The fluctuations between inundation and subsequent drying, associated with periodically flooded soils, are major drivers of spatial and temporal differences in soil properties that affect the biogeochemical processes taking place in floodplain soils (Schulz-Zunkel et al., 2015; Tockner et al., 2010). These changes include; a decrease in redox potential (E_H), which leads to, for example, reduction of iron (Fe) and manganese (Mn), which in turn can influence the soil pH (Rinklebe and Shaheen, 2017). Other processes affected include sulphur-cycling, changes to the presence of chelating agents such as dissolved organic carbon, mineralisation of POM and suppression of microbial activity (Poot et al., 2007; Puchalski, 2003; Schulz-Zunkel et al., 2015; Schulz-Zunkel and Krueger, 2009). Ibragimow, Walna, and Siepak (2013) showed, through analyses of fluvial samples before and after a flood, that the physicochemical properties (grain size, E_H , pH, POM, and calcium carbonate contents) as well as the total and available concentration of PTEs had changed. Harvey et al. (2019) found that after UK floods receded in the winter of 2013-14 there was a decrease in the soil bulk density, pH and available P. The flood was found to have had a negative effect on the overlying vegetation and caused a shift in the microbial community structure.

Inundation during a flooding event can carry PTEs dissolved in rising groundwater and potentially contaminated suspended sediment from upstream overbanking water, depositing this onto the floodplain during a flood (Acreman et al., 2003; Bednářová et al., 2015; Du Laing et al., 2009; Gröngröft et al., 2005; Junk et al., 1989; Rudiš et al., 2009; Tockner and Stanford, 2002; Weber et al., 2009). Subsequently, this deposition of suspended riverine sediments/POM by flood water results in the floodplain topsoil becoming a sink for PTEs (Du Laing et al., 2009; Frohne et al., 2011; Nshimiyimana et al., 2014; Overesch et al., 2007; Rinklebe et al., 2007; Visser et al., 2012; Zhao and

Marriott, 2013). As a result, floodplain topsoil (uppermost 15cm) can often initially contain elevated concentrations of PTEs such as the metalloid; arsenic (As), and metals; chromium (Cr), copper (Cu), lead (Pb), and zinc (Zn), but later due to post-depositional reactions with organic matter/other organic components the PTEs concentrations will vary (Adekanmbi et al., 2020; Ciszewski and Grygar, 2016; Hurley et al., 2017; Izquierdo et al., 2013; Jiao et al., 2014; Kelly et al., 2020). When laboratory experiments are undertaken on samples gathered from floodplain site, soils are collected as single or composite samples, air or oven dried and then homogenised, resulting in a loss of soil stratigraphy and therefore the potential differences in PTEs concentration with depth may be unaccounted for (Ciszewski and Grygar, 2016). Zhao and Marriott, (2013) looked at PTEs concentrations along a vertical profile and found that there were peak values at varying depths; affected by translocation and duration of inundation. The process of breaking up of soil samples for laboratory experiments will make interpretation of PTEs levels difficult. Kelly et al. (2020) took intact soil cores to overcome this and more closely reflect natural samples, they too found the duration of inundation influenced the fate of PTEs.

The biological health of floodplain soils is important as they act as an interface between terrestrial and aquatic environments, therefore playing an important role in maintaining the environmental quality of surface waters (Izquierdo et al., 2013; Stuart and Lapworth, 2011). Artificial or constructed wetlands have been used for flood and pollution control; storing and filtering excess water to protect rivers from various kinds of runoff e.g. high nutrient loads from farm land (Blackwell and Pilgrim, 2011; Ellis et al., 2003; Rizzo et al., 2018). An example of this technology is demonstrated at the Rothamsted Research North Wyke experimental farm (Pulley and Collins, 2019). Even if river and groundwater water quality improves due to the implementation of more stringent environmental policy, contaminated floodplains remain as a legacy of historic upstream pollution (Bradley and Cox, 1990; Förstner, 2004; Kowalik et al., 2004). With increased frequency and duration of flooding, there is the possibility that changes to soil properties and biogeochemical processes will ultimately lead to the mobilisation of PTEs from floodplain soils (Ciszewski and Grygar, 2016; González-Alcaraz and van

280 Gestel, 2015). Therefore, historically contaminated floodplains may become a source of legacy
281 pollution to the surrounding environment (Kelly et al., 2020; Pulchalski, 2003; Schulz-Zunkel and
282 Krueger, 2009), as shown in Figure 1.




Figure 1: How floodplains may switch from being a sink of pollution to becoming a source of legacy pollution: A) PTEs contaminated river sediment (red) due to industry in the catchment upstream, B) heavy rainfall influences the receiving catchment (increased river flow and groundwater level), resulting in flooding and the deposition of contaminated sediment onto the adjacent floodplain; dissolved contaminants may also reach the floodplain surface via rising groundwater, C) Later, the river is uncontaminated (brown) due to rising environmental quality standards, with legacy of PTEs contamination (red) in the river bank and floodplain soil, D) heavy rainfall results in flooding of the contaminated floodplain, mobilisation of the legacy PTEs by desorption and resuspended particulate matter into the surrounding environment and thus making them potentially available for uptake by

vegetation and soil organisms, as well as pollutant transfer leaching into the overlying flood water, the groundwater, and ultimately the river. Created with BioRender.com.

2.Impact of flooding on the mobility of potentially toxic elements in floodplain soil

2.1 PTEs in floodplain soil

Several PTEs are also essential nutrients that are required in low concentrations for healthy functioning and reproduction of microorganisms, plants, and animals, although may become toxic in high concentrations, these include; Cu, Cobalt (Co), Nickel (Ni), Vanadium (V), Zn, chlorine (Cl), Mn, Fe, boron (B), and molybdenum (Mo) (Adamo et al., 2014; Hooda, 2010; Wyszowska et al., 2013). Other PTEs are non-essential and can cause toxicity even when they are found at low concentrations, these include; As, Pb,) and mercury (Hg); (Adamo et al., 2014; Nriagu et al., 2007; Wuana et al., 2011; Wyszowska et al., 2013). Cadmium (Cd) is generally considered a non-essential element to soil organisms, but it has been found to be beneficial to some microalgae (Xu et al., 2008) Chromium can be considered a micronutrient but its toxicity depends on its valence state (i.e. Cr (VI) is the more mobile and toxic form compared with Cr (III)). Redox potential therefore not only affects the mobility of PTEs, but also their toxicity (Lee et al., 2005; Shahid et al., 2017). The consequences of PTEs contamination of soils are rarely observed with immediate effect, rather they tend to cause delayed adverse ecological changes, due to the fact that PTEs are persistent in the environment for long periods, non-biodegradable and can only be bio-transformed through complex physico-chemical and biological processes (Chrzan, 2016; Czech et al., 2014; Hooda, 2010). PTEs cause adverse ecological effects on plants and organisms such as impacting their activity, growth rate/yield, metabolism and reproduction, causing symptoms of physiological stress and potentially death. The extent of the adverse effect depends on the exposure route (ingestion, dermal absorption or uptake of pore water) and time, resistance (related to residence time of the PTEs in the environment) and detoxification mechanisms of the plant or animal (Alloway, 2013; Eggleton and Thomas, 2004; Ehlers and Loibner, 2006; Hooda, 2010; Pan et al., 2018; Shahid et al., 2017; Winger et al., 1998). Leaching of PTEs from

the floodplain soil into the groundwater or river will also cause adverse effects to aquatic organism in these environments (Zia et al., 2018).

PTEs are either present naturally in the floodplain soil from the underlying or upstream geology and subsequent geogenic processes (e.g. weathering of parent material, emissions from volcanoes, forest fires) or introduced by anthropogenic sources, including solid and dissolved inputs from; aerial deposition, transport emissions, industrial, municipal and diffuse runoff from agricultural practices landfills and sewage treatment facilities (Alloway, 1995; Álvarez-Ayuso et al., 2012). PTEs can be adsorbed to colloidal suspended particulate material, transported in the river water and accumulate in the floodplain soil during inundation (Du Laing et al., 2009; Frohne et al., 2011; Peijnenburg et al., 2007; Rinklebe et al., 2007). PTEs have been found to be primarily associated with fine-grained clay or silt minerals and can reside in the floodplain for longer when compared with river sediments, as they are less likely to be susceptible to erosion (Lučić et al., 2019; Malmon et al., 2002). Contamination of the floodplain soil may result from a point source such as a sewage treatment facility, or from diffuse sources that have no specific point of discharge (e.g., agricultural applications). Impacts of diffuse pollution are difficult to predict as they can be affected by weather systems, meaning soils far from the source may be affected (Gregory et al., 2015; Neal et al., 1996). The anticipated changes to intense rainfall may result in increased delivery of diffuse pollution to rivers and groundwater (Arnell et al., 2015; Foulds et al., 2014), particularly as contaminated floodplain soils may become a diffuse source of pollution themselves during a flooding event (Schulz-Zunkel and Krueger, 2009).

2. 2. Influence of flooding on PTEs mobility

During a flooding event, biogeochemical processes occur in the floodplain soil at the oxic-anoxic interface and in the anoxic layers. The kinetics of these processes are of great importance because the location of the oxic-anoxic interface is subject to change due to fluctuating water table levels (Du Laing et al., 2009; Puchalski, 2003). In their review of trace metal behaviour in floodplain

sediments, Du Laing et al. (2009) state that the spatial occurrence of processes affecting metal mobility and availability is largely determined by the topography of the floodplain. Remobilisation of PTEs from sediments into the overlying water column during a flooding event depends on the flood regime; the frequency of these intense floods which flush or remobilise contaminated material as well as the duration or alternation of flood with dry spells (Arnell et al., 2015; Foulds et al., 2014; González-Alcaraz and van Gestel, 2015). Whilst research has suggested that the longer the flood duration, the greater the metal mobility (Shaheen et al., 2014a, 2014b), Stafford et al. (2018) suggest that even short periods of soil saturation can have an influence the solubility of PTEs.

There are conflicting results in the literature regarding the effect of flooding on the mobility of PTEs, expressed by changes in PTEs concentration (increase or decrease), in floodplain soils (Table S1). This may largely be the result of different site-specific conditions (e.g. soil pH, texture, mineralogy) or different laboratory set-ups (e.g. submerging soils in deionised water, or the use of inert gas to simulate the anoxic conditions of a flood), illustrating the complexity of the processes involved in mediating PTEs mobility in floodplain soils (Abgottspon et al., 2015; Du Laing et al., 2007; Frohne et al., 2011; Schulz-Zunkel et al., 2015). Many of the considerations in the literature are founded on research of soils or sediments in microcosm experiments, which often involves homogenising the soil samples, resulting in loss of natural soil structure, loss of roots and biota, short-exposure time to flood conditions, and the control of variable factors such as temperature and soil water conditions (Frohne et al., 2011; Rinklebe et al., 2010). Redox conditions are often simulated and controlled through additions of O_2 , to increase E_H , and N_2 , to lower E_H (Frohne et al., 2014, 2011; Schulz-Zunkel et al., 2015; Shaheen et al., 2016; Shaheen and Rinklebe, 2017). These differences make extrapolation of these laboratory-based findings to field situations difficult (Hooda, 2010).

A key factor in determining the fate and transport of PTEs is their chemical form which, in combination with environmental factors, can influence their mobility in the soil. The chemical form of an element is often referred to as its “speciation”, “oxidation state”, or “valence” (Rodgers et al., 2015;

Wuana et al., 2011). There are important redox sensitive PTEs for which the oxidation state has a large influence on solubility and mobility. For example, Cr(VI) is more mobile than Cr(III), but As(V) is less mobile than As(III) (Frohne et al., 2015; Rinklebe et al., 2016; Schulz-Zunkel et al., 2015; Shaheen et al., 2014b; Yang et al., 2015). Speciation of PTEs within the environment has a distinct influence upon their behaviour; specifically, reactivity, toxicity, mobility and bioavailability within the floodplain (Du Laing et al., 2009; Gambrell, 1994; Hooda, 2010; Rodgers et al., 2015). This understanding is important for predicting the environmental impact of contaminated soils, although we are only beginning to converge on consensus on how bioavailability or speciation soil tests can help with risk-assessments, while this is slowly introduced into legislation (Cipullo et al., 2018; Naidu et al., 2015, 2008; Ng et al., 2015).

2. 3. Changes to soil physical and chemical properties that influences PTEs mobility

Potentially toxic elements present in soils are often adsorbed to or protected within aggregates that are stabilised by organic matter. During a flooding event, these particles may be leached through the soil profile, or suspended in flood waters where they may be redistributed across floodplain soils, or be carried downstream by the river, potentially contributing to river pollution of the contamination of downstream floodplains. The solubility and therefore mobility of PTEs from the soils to the surrounding environment depends largely on the intrinsic soil physical and chemical properties (texture, availability of soil particulate surfaces and dissolved organic matter, salinity and the presence of Fe/Mn oxides, carbonates, phosphates and sulphides) and a range of variables that are directly affected by periodic inundation of the floodplain, including; soil pH, redox potential (E_H), dissolved organic carbon (DOC) and the valance of individual PTEs (Adewuyi and Osobamiro, 2016; Dawson et al., 2010; Du Laing et al., 2009; Frohne et al., 2015; González-Alcaraz and van Gestel, 2015; Lee et al., 2005; Puchalski, 2003; Rinklebe and Du Laing, 2011; Schulz-Zunkel and Krueger, 2009; Shaheen et al., 2016; Shaheen and Rinklebe, 2014; Steinnes, 2013). A conceptual model (Figure 2) has been produced based on our literature review (Table S1) as a way of visualising the various factors and processes influencing the solubility of PTEs in a floodplain soil.




Figure 2: Conceptual model depicting the key processes influencing the solubility of Potentially Toxic Elements (PTEs) after a soil becomes flooded.

a) Oxygen is rapidly consumed by microbial and root respiration, decreasing the redox potential (E_H). b) Decreasing E_H can lead to redox sensitive elements (e.g. As and Cr) changing valence state, directly affecting solubility. c) Greater soil moisture brings dissolved organic matter (DOM) into solution. d) Reducing conditions (lower E_H) leads to the release of more DOM. e) Lower E_H results in the reduction of Fe and Mn, consuming protons (H^+) and increasing pH. f) an increase in pH often results in the release of more DOM. g) DOM acts as a chelating agent, forming soluble organo-metal complexes with PTEs desorbed from soil surfaces. h) as pH increases metal cations (e.g. Cu, Pb, Zn) are adsorbed on pH-dependent adsorption sites of particulate matter. i) as pH increases, anions and oxy-anions (e.g. As) are desorbed from pH-dependent adsorption sites. j) Dissolution of reducible Fe and Mn oxides is facilitated by increasing pH. k) Microbial reduction of Mn and Fe oxides increases their solubility and can cause reductive dissolution of co-precipitated PTEs. l) An increase in pH facilitates the precipitation of insoluble metal sulphides. m) Microbial reduction of sulphate results in the precipitation of metal sulphides. n) Release of adsorbed PTEs from soil surfaces increases PTEs solubility. o) Immobilisation of PTEs through adsorption processes reduces PTEs solubility. p) Reductive dissolution of PTEs associated with Fe and Mn oxides increases PTE solubility. q) Precipitation of PTEs as metal sulphides decreases PTEs solubility.

Soil physical, chemical and biological processes determine the mobility and redistribution of PTEs (Hooda, 2010). These processes include; sorption, desorption, dissolution and precipitation (Puchalski, 2003; Wijngaard et al., 2017). Subsequently, PTEs are redistributed into different geochemical fractions, associated with other soluble species, released from the soil matrix into the soil solution or porewater, and transferred through the ecosystem and food web to other terrestrial or riparian areas downstream from the floodplain; thus potentially becoming a risk to human and environmental health (Adamo et al., 2014; Adewuyi and Osobamiro, 2016; Baran and Tarnawski, 2015;

Dang et al., 2002; Du Laing et al., 2009; Rinklebe et al., 2016; Schulz-Zunkel et al., 2015; Shaheen et al., 2014a, 2014b; Sizmur et al., 2011). Sorption processes that control PTEs mobility and bioavailability in soil are affected by the soil pH, redox and their interactions with other ions and substances present in soil solution (Antoniadis et al., 2018; Frohne et al., 2011; Ostergren et al., 2000; Violante, 2013).

Sorption processes are influenced by the changing conditions that flooding brings, particularly with regards to soil moisture content, temperature and redox potential. The mobility of PTEs in flooded soils is closely related to changes in redox potential which, in turn, is altered by flooding. This can have direct impacts on the mobility of redox sensitive PTEs (e.g. As and Cr). Inundation of soils with floodwater may indirectly affect PTEs mobility and speciation because it also influences, the population, community composition, and behaviour of invertebrates inhabiting the floodplain which, in turn, influence the mobility of PTEs through their burrowing and bioturbation behaviour. For example, earthworms are known to increase the mobility of PTEs due to passage through the earthworm gut (Sizmur et al., 2011; Sizmur and Richardson, 2020) and their populations are suppressed by flooding events (Plum and Filser, 2008; Kiss et al., in review). Bioturbation/bioirrigation behaviour by chironomid larvae has been found to increase oxygen uptake at the soil/sediment-water interface, promoting POM decomposition that results in the release of dissolved organic matter and subsequent release of PTEs (He et al., 2019). Furthermore, the reduction of Mn and Fe can cause reductive dissolution of co-precipitated PTEs, and an increase in pH facilitates the precipitation of PTEs as insoluble sulphides. The mobility of PTEs can therefore increase or decrease due to the net effect of these processes (Figure 2). Which process dominates will depend primarily on the mineralogy of the soil.

The following sub-sections will explain how key soil physical and chemical properties are affected by flooding and how this influences PTEs mobility, followed by a discussion on the role of soil organisms and plants in mediating PTEs mobility in floodplain soils. Attention will be given to how each of these factors influence each other to distinguish direct and indirect impacts on PTEs mobility.

2.3.1 Soil texture and related properties

Soil texture is a stable property that refers to the physical composition of mineral fragments; sand, silt and clay and varies due to differences in underlying or upstream geology. The texture and related clay mineralogy reflect the particle/pore size distribution and overall soil surface area (Amacher et al., 1986) which, in turn, affects the soils' water holding capacity (WHC); the maximum quantity of water a soil can potentially contain, also known as the field capacity (Stürck et al., 2014). Therefore, soil physical properties play a role in flood duration because they determine the soils' ability to receive (via infiltration) and drain water during a rainfall event (Rinklebe et al., 2007). Clayey soils are likely to be saturated for longer than freely draining sandy soils (Sherene, 2010). Soil hydraulic (water retention and hydraulic conductivity curve) as well as thermal properties (thermal conductivity and heat capacity) affect the hydrothermal regime of the soil. Together these properties determine the ease in which water, and dissolved PTEs, moves through the soil pore continuum, how much water can be stored in the pore volume, and how soil temperature varies with depth. These properties are strongly dependent on soil texture, pore size distribution and mineralogy (Hillel, 1998; Tack et al., 2006; Thomas et al., 2016). Soil temperature affects the flow of water through the soil due to changes in viscosity and hence affects infiltration calculations (Gao and Shao, 2015; Prunty and Bell, 2005), so this is often corrected for when reporting hydraulic conductivity data (Thomas et al., 2016).

PTEs must be in the soluble phase or associated with colloids to be transported through the soil. The soil properties will play a part in the movement of PTEs into and out of the soil solution. Clay minerals and organic matter compounds have a large number of binding sites, so act as adsorption surfaces for PTEs in soils. The type of clay mineral present (kaolinite, illite, montmorillonite etc.) will also affect the specific surface area (Meegoda and Martin, 2019; Tack et al., 2006). As a result, soils with high clay and silt (fine fractions) tend to retain higher amounts of PTEs, compared to coarse textured sandy soils (Sherene, 2010; Zhao et al., 1999).

2.3.2 Organic matter

Soil POM, along with the surfaces of clay particles and Fe and Al oxides, acts as a binding phase for PTEs due to the attraction of positively charged cations to negatively charged surfaces (Evans, 1989). Thus, dissolved organic matter raises the cation exchange capacity (CEC) of a soil, and is thus considered to be an important factor controlling PTEs distribution and mobility in floodplain soils and sediments (Baran and Tarnawski, 2015; Bufflap and Allen, 1995; Du Laing et al., 2009; Ehlers and Loibner, 2006). The mechanisms that bind the PTEs with particulate and dissolved organic matter include adsorption, complexation and chelation (Alvim Ferraz and Lourenço, 2000; He et al., 2019; Selinus et al., 2005). Floodplains are subject to changing water table levels and occasional inundation that brings about associated changes in redox conditions. This can result microbially mediated soil POM degradation, either during prolonged periods of flooding or in the subsequent oxidising conditions when the flood recedes, which releases organically bound PTEs, such as As, Cu, Co, Cr, Ni, Pb, and Zn from the soil into the soil solution (Adewuyi and Osobamiro, 2016; Alvim Ferraz and Lourenço, 2000; Dang et al., 2002; Kalbitz and Wennrich, 1998; Koretsky et al., 2007; Rinklebe and Du Laing, 2011). Therefore, the extent to which flooding of soils results in the mobilisation of PTEs into solution is mediated by the proportion of the PTEs that are associated with soil POM, and the susceptibility of this organic matter to degradation (as a result of microbial activity (Fe(III) and Mn(IV)-reducing micro-organisms) under reducing conditions. The free ions that are then in solution are highly reactive with the solid phase and are thought to be a major determinant of bioavailability and causing the most significant biological effects (Bufflap and Allen, 1995; Dang et al., 2002; Dawson et al., 2010; Degryse et al., 2009; Lloyd, 2003).

2.3.3 Salinity

Salinity is proportional to the conductivity of a sample solution; which is a measure of its ability to conduct or carry electric current and depends on the presence of charged ion species (anions and cations) (Ander et al., 2016; de Souza Machado et al., 2018; De Vivo et al., 2008)). Increasing salinity in flood water is associated with an increase in major cations that compete with PTEs for sorption

502 sites. This competition promotes PTEs desorption from the floodplain soil in the absence of sulphides
503 and hence increases total PTEs concentrations in the soil porewater (Rinklebe and Du Laing, 2011).
504 The presence of Ca-salts releases more PTEs into the soil solution compared with Na-salts that are less
505 competitive for sorption (Du Laing et al., 2009; Hahne and Kroontje, 1973).

506 Changes in salinity may affect the soil physical properties and result in a destabilisation of the
507 soil structure (Gregory et al., 2015). The salinity of the water causes a neutralisation of negatively
508 charged clay particles, followed by flocculation (particles attaching together) which increases the
509 deposition of sediments (along with the PTEs adsorbed to them) onto the floodplain. This process
510 results in the floodplain becoming a sink for PTEs (Rinklebe and Du Laing, 2011). An extended flood
511 duration, particularly when accompanied by low flow-rates (including stagnant water), results in
512 sedimentation of fine grain sediment and organic matter that may have PTEs bound (Ciszewski and
513 Grygar, 2016; Du Laing et al., 2009; Shaheen and Rinklebe, 2014).

514 2.3.4 Redox potential (E_H)

515 Waterlogging of soils generally results in a reduction in oxygen availability due to rapid
516 consumption of oxygen by soil microbial activity and root respiration (Du Laing et al., 2007; Rinklebe
517 and Du Laing, 2011) and because the dissolution of oxygen through water is many times slower than
518 through air (Alloway, 1995; Du Laing et al., 2009; Frohne et al., 2015; Schulz-Zunkel et al., 2015). The
519 soil microbial community (e.g. bacterial species such as *Thiobacillus ferrooxidans*, *Thiobacillus*
520 *thiooxidans* and *Leptospirillum ferrooxidans*) then uses alternative electron acceptors (such as nitrate,
521 sulphate and Fe/Mn oxides), in anaerobic respiration, which results in a decrease in redox potential
522 (E_H) (Maluckov, 2017) as the floodplain soils change from oxic ($[O_2] > 30 \mu\text{mol L}^{-1}$) to anoxic
523 ($[O_2] < 14 \mu\text{mol L}^{-1}$) conditions (Bellanger et al., 2004). Associated alkalinity generation drives increases
524 in soil pH, a change which can be observed after a few days (Du Laing et al., 2007; Johnston et al.,
525 2014; Karimian et al., 2017). Soil temperature has been found to dictate the rate and type of redox
526 reactions; with soils at low temperatures (1 - 4 °C) requiring greater durations of saturation (20 days)

before the onset of reducing conditions were seen, whereas soils at higher temperatures (above 9 °C) only required 2 days of saturation (Vaughan et al., 2009).

Redox potential has important effects on the speciation of As, Cu and Cr, as well as N, S, Fe, Mn, because these elements can exist in soils in more than one oxidation state (Selinus et al., 2005) and solubility depends on oxidation state. Copper solubility decreases after reduction from Cu (II) to Cu (I) under anaerobic conditions and the presence of electron donors (Fe (II)) and bacteria. However, other PTEs such as Cd and Zn change valence state as a consequence of redox dependent pH changes, complexation with organic matter or precipitation with Fe and Mn (hydr)oxides or sulphides (Du Laing et al., 2009; Frohne et al., 2011). Shaheen et al. (2014a) demonstrated that sufficient time is needed for transformations between valence states to take place. For example, the oxidation of Cr from Cr (III) to the highly mobile Cr (IV) form was found to be a slow process. This means that with shorter flooding duration and quicker cycling between oxic and anoxic conditions, Cr mobility may be difficult to predict.

The presence of variable charge minerals, such as Fe and Mn oxides, phosphates, carbonates and sulphides provide a reaction surface for sorption processes, allowing PTEs to bind and become immobilised (Antoniadis et al., 2018; De Jonge et al., 2012; Sipos et al., 2014; Violante, 2013). Reducing conditions change the oxidation state of Fe and Mn, increase their solubility and may have indirect effects (known as reductive dissolution) on the mobility of associated metal cations (e.g. As, Cd, Cu, Ni, Pb, and Zn), releasing them from the solid phase to pore waters, depending on flood duration (Abgottspon et al., 2015; Ciszewski and Grygar, 2016; Du Laing et al., 2009; Frohne et al., 2011; Karimian et al., 2017; Rinklebe and Du Laing, 2011; Schulz-Zunkel et al., 2015; Shaheen et al., 2016, 2014b; Vaughan et al., 2009). Redox processes are a key factor for the reductive dissolution of Mn and Fe (hydr)oxides, these processes are often catalysed by microorganisms and result in the release of PTEs from the sediment (Du Laing et al., 2009; Frohne et al., 2011; Stafford et al., 2018; Yang et al., 2015). Relatively insoluble Fe(III) and Mn(IV) prevail under aerobic soil conditions providing sorption

surfaces for many metals, whereas under anaerobic conditions Mn(IV) and Fe(III) are reduced to more soluble forms (Mn(II) and Fe(II)) with consequential dissolution of Mn and Fe hydrous oxides, co-sorbed PTEs ions (e.g. As, Cd, Cr, Ni and Pb), are released into soil solution (Simmler et al., 2017; Stafford et al., 2018; Yang et al., 2015). After inundation, Fe and Mn may re-precipitate as oxides and can bind (by desorption or co-precipitation) the trace metals back into the solid state (Ciszewski and Grygar, 2016; Davranche et al., 2011; Du Laing et al., 2009).

Decreasing of E_H can initiate microbial sulphate reduction and this can reduce the mobility of some PTEs (e.g. As, Cd, Cu, Cr, Ni and Pb) through coprecipitation of metal cations with sulphides (Abgottspon et al., 2015; Borch et al., 2010; Weber et al., 2009), although many of these minerals are metastable and so prone to change (Karimian et al., 2018). Yang et al., (2015) put mixed sediment samples into a laboratory culture tanks and found that microbially induced release of sulphur with subsequent As precipitation was more important for controlling As adsorption/desorption than reductive dissolution of Fe/Mn oxides. As the flood recedes, the floodplain soils undergo drying and aeration that change the conditions from anoxic back to oxic. The now oxic environment causes sulphides to be oxidised, which then releases PTEs back into the pore waters (Abgottspon et al., 2015; Du Laing et al., 2007; Frohne et al., 2011). In addition to this, when exposed to oxygen and water, sulphides are oxidised to sulphates which leads to the formation of sulphuric acid thereby causing a decrease in pH and release of the PTEs (Emerson et al., 2017; Forstner and Wattman, 1981). Frohne et al. (2011) suggested that the mobility of Cd, Cu, Mn, Ni and Zn under oxidising conditions could be attributed to dissolution of sulphides and the resulting release of those metals. The extent to which the mineralogy of a floodplain soil is dominated by Fe/Mn oxides or sulphates may dictate whether PTEs are mobilised or immobilised during inundation, and the extent to which this phenomenon is reversed after floodwater recedes.

2.3.5 Soil pH

pH is a measure of the hydrogen ion concentration and can also be referred to as the degree of acidity or alkalinity. The soil pH is affected by flooding because of a well-established correlation between soil pH and changing redox conditions; as a soil becomes flooded, this creates reducing conditions where (H^+ ions) are consumed (for example due to reduction of Fe and Mn oxides) and the pH increases (Rinklebe and Shaheen, 2017; Weber et al., 2009). When the flood recedes, oxidation processes produce protons and decrease the pH (Adewuyi and Osobamiro, 2016; Frohne et al., 2015, 2011; Rinklebe and Shaheen, 2017; Shaheen and Rinklebe, 2017). Furthermore, on exposure to the atmosphere, when flooding recedes, dissolved organic carbon (DOC) is converted to CO_2 , which dissolves into porewater as carbonic acid, subsequently further reducing the soil pH (Peacock et al., 2015). However, this negative correlation between E_H and pH hasn't always been observed (Du Laing et al., 2009; Frohne et al., 2015). This is because the degradation of POM such as plant residues, by soil microbes, may increase the soil pH due to ammonification of the residue N (Xu et al., 2006).

As the pH changes, processes such as precipitation, co-precipitation and sorption/desorption of PTEs from organic matter or clay minerals occur, altering the chemical composition as well as reaction rates (Frohne et al., 2011). The soil pH plays an important role in mediating the mobility of PTEs and their availability for plant uptake, as the protons compete with metal cations for exchange sites on the surface of soils. Some of these exchange sites, particularly those associated with soil organic matter, are pH-dependent and thus only become deprotonated at high pH. A decrease in pH is generally accompanied by an increase in the mobility of most PTEs that are metal cations (e.g. Cd^{2+} , Cu^{2+} , Co^{2+} , Ni^{2+} , Pb^{2+} and Zn^{2+}) (Grönröft et al., 2005; Sherene, 2010). Thus, as pH increases there is a subsequent decrease in the mobility of these PTEs (Giacalone et al., 2005). The extent to which PTEs mobility decreases in soils during flooding, due to a redox-induced increase in pH, is likely to depend on the proportion of PTEs in the soil that are associated with pH-dependent exchange sites) which are typically associated with soil organic matter) and the pH of the soil prior to the flooding event.

2.3.6 Dissolved organic matter (DOM)

The increase in pH of soil solutions with lower redox potential (reducing conditions) is often accompanied by a release of dissolved organic matter (DOM) and the subsequent formation of soluble organo-metal complexes (Abgottspon et al., 2015; Alvim Ferraz and Lourenço, 2000; Frohne et al., 2011). The presence of DOM in floodplain soils acts as a chelating agent which has a strong binding ability and increases the mobility of PTEs into pore waters and subsequently into river water or groundwater (Dawson et al., 2010; Du Laing et al., 2009; Shaheen et al., 2014b). The greater the concentration of DOM in porewater, the more PTEs that are held in solution, and (to maintain an equilibrium) the more PTEs that desorb from the surfaces of the soil to replenish the free ion concentrations in the porewater, thus increasing PTEs mobility. Greater concentrations of DOM have been observed with decreasing E_H , which may be due to suppressed microbial carbon consumption under anoxic conditions (Frohne et al., 2015). Shaheen et al. (2014a) highlighted that increases in DOM associated with lower E_H may help to catalyse changes in the valence state of PTEs; for example, of Chromium (III) to (VI).

2.3.7 Temperature

As temperatures are predicted to increase as a result of climate change, they may become a factor that contributes to greater release of PTEs from the soil during a flood (Visser et al., 2012). Soils are affected by variations in air temperature which, in turn, affects the rate of biogeochemical processes during a flooding event, including decreasing redox potential and, ultimately, influences the rate and extent to which PTEs are released/ desorbed from POM into surface water and groundwater (Arnell et al., 2015; González-Alcaraz and van Gestel, 2015; Sánchez-Rodríguez et al., 2019; Shaheen et al., 2016; Stahl et al., 2013). Increases in temperature raise the ion activity in soil solution, and also make plants more active, which may lead to greater plant root uptake of soil water and dissolved/labile PTEs within this water (Sherene, 2010). Arsenic release from flooded soils was found to have temperature dependence, with As solubilisation increasing as temperature increased (Simmler et al., 2017; Visser et al., 2012). Temperature increases are attributed to a decrease in the

water viscosity resulting in dissociation of molecules and a subsequent increase in the number of ions in the solution. For every degree Celsius increase in temperature there is an observed increase in electrical conductivity of 1.9% (Ander et al., 2016; Ma et al., 2011).

Many of the processes (e.g. redox reactions) described in the previous sections are microbially mediated and temperature dependent, and so the extent to which they affect the mobility of PTEs depends on their kinetics and the duration that floodplain soils are inundated. Changes in soil pH and DOM have been shown to exert a greater influence than E_H on the mobility of PTEs when considering shorter flood-dry cycles (Shaheen et al., 2014b, 2014a). However, Dang, Liu, and Haigh (2002) found that with increasing flood duration, more trace elements were transformed from inert phase to exchangeable fractions, increasing mobilisation. Soil redox processes are important for protecting environmental health; however, the kinetics and mechanisms remain poorly characterised and understood (Abgottspon et al., 2015; Borch et al., 2010; Pulchalski, 2003). Many of the studies reviewed in this manuscript undertook experiments in the laboratory at temperatures that are higher than the soil temperatures typically found *in-situ* at the location where the soil samples were collected, and so the rates at which reactions occur and the subsequent mobilisation of PTEs may be overestimated in these experiments. It is not possible to verify the extent of the overestimation, so more *in-situ* experiments are needed to investigate and provide quantification of the differences between laboratory and *in-situ* experiments.

Groundwater and river temperatures may have a strong effect on floodplain soil temperatures (Andersen, 2018). In warmer seasons they will generally be cooler than soil temperatures, but the opposite occurs in cooler seasons. Also, changes in soil moisture content, as a result of flood events, will affect the soil thermal properties such as thermal conductivity and heat capacity (Lu et al., 2007), thereby also affecting the spatial and temporal variation in the soils' temperature regime. A laboratory microcosm experiment with mining-contaminated topsoil and subsoil samples saturated for up to 41 days at temperatures ranging between 10-25°C, found that soil temperature increased the

solubilisation of As, particularly in the topsoil when saturated for 1-2 weeks (Simmler et al., 2017). This means, for some PTEs, flooding during warmer seasons may result in greater mobilisation than flooding during cooler seasons. More seasonal field observations are required to understand the subtle interactions and feedbacks between soil moisture, floodwater temperature, and mobility of PTEs.

2. 4. Soil biological processes that influence PTEs mobility

2.4.1 Soil organisms

Floodplain soils contain a great diversity of organisms that are known to contribute to the physical structure of the soil/sediment through bioturbation which influences the biogeochemical cycling of PTEs through oxygen diffusion, redox gradient and decomposition of dissolved organic matter (Classen et al., 2015; He et al., 2019; Hooda, 2010; Selinus et al., 2005). As the soil pore spaces are filled with water, oxygen diffusion is low so microbial respiration relies on alternative electron acceptors (e.g. NO_3^- , Mn, Fe and S), resulting in reducing conditions (decreasing E_H) that simultaneously increase pH (Matern and Mansfeldt, 2016), and the changes to PTEs mobility (Figure 3) that are described in previous sections. Changes in the chemical speciation of PTEs can also occur due to microbial processes in reducing conditions, for example, sulphate reducing bacteria can methylate Hg in anoxic conditions (Ma et al., 2019).




Figure 3: Soil microbial processes during inundation of floodplain soil influences mobility of PTEs; (left-hand side) generic metals with valence state (denoted by M^{n+}) are coprecipitated to Fe oxides and are released due to reductive dissolution and (right-hand side) sulphate reduction (sulphate to sulphide) results in some metal (denoted by MS) precipitation, some of the metal (M^{n+}) remains in the pore water. Created with BioRender.com.

PTEs that are present in floodplain soils are often protected within the soils' aggregates, which are stabilised by POM. However, inundation can stimulate the soil microbial community, which is sensitive to disturbance, accelerating the refractory organic matter mineralisation and destabilisation of aggregates, exposing and increasing the mobility of PTEs in the soil (Du Laing et al., 2009; Gall et al., 2015; González-Macé et al., 2016; He et al., 2019; Rawlins et al., 2013). Tack et al. (2006) found that the drying of sandy soils caused an increase in soil solution metal concentrations, compared with the same soils maintained at field capacity. This observation was attributed to microbial effects, increasing the solubility of dissolved organic matter.

Flooding has been found to shift the soil biological community structure and function. These changes include a reduction of Gram-positive bacteria, mycorrhizal fungi and earthworms found under flooded conditions (Gregory et al., 2015; Harvey et al., 2019; Unger et al., 2009). Harvey et al., (2019) found that flooding induced short-term alterations to soil microbial biomass but these changes did

not persist in the long term; they concluded that temperate systems may be resilient to winter flood stress. The seasonal timing of floods influences the effect that flooding has on the soil microbial community, and so may result in different effects on, and recovery of, the soil microbial community. Sánchez-Rodríguez et al., (2019) subjected a UK agricultural grassland soil in an intact laboratory microcosm to flooding and found that summertime flooding (25°C), resulted in a loss of actinomycetes and arbuscular mycorrhizal fungi, and that these changes persisted post-flood. They expected microbial biomass to increase with flooding at higher temperatures, due to degradation of vegetation releasing labile carbon. However, they found that maintaining live roots and an active rhizosphere were more important for preserving the microbial community in grassland soils. Earthworms also play a role in increasing the mobility and availability of PTEs in floodplain soil through their activity causing changes to the soil microbial populations, pH, DOC or metal speciation (Sizmur et al., 2011; Sizmur and Hodson, 2009) which in turn influences PTEs mobility as discussed in the above sections.

As the PTEs are released into the aqueous phase and mobilised in the environment, they present a potential risk to soil organisms (Ehlers and Loibner, 2006; González-Alcaraz and van Gestel, 2015). Soil organisms uptake PTEs via ingestion of polluted soil, food or pore water and/or via dermal uptake or absorption of soil water, with the soil water being the more important of the two pathways (Chrzan, 2016; Hobbelen et al., 2006; Sivakumar and Subbhuraam, 2005). Vijver et al. (2007) found that the frequency of flooding did not result in consistent changes in the internal PTEs concentrations of earthworms. Earthworms accumulate PTEs in their chloragogenous tissue and have a mechanism that allows them to regulate their internal PTEs concentrations, so when they are introduced to contaminated soils the earthworms reach an equilibrium and when they are returned to uncontaminated/"clean" soils they are able to detoxify and eliminate essential metals through excretion (e.g. Cu and Zn), but not non-essential metals (e.g. Cd and Pb) as detoxification processes involve sequestration within an inorganic matrix or organic ligand (Sizmur and Hodson, 2009; Spurgeon and Hopkin, 1999). While microbes can tolerate larger quantities of essential PTEs, in excess both essential and non-essential PTEs (e.g., Al, As, Cd, Hg, Pb, Zn) can adversely affect microbial

communities by altering community structure and taxonomic richness; reducing the microbial biomass and lowering their enzyme activity which results in a decrease of soil diversity (Gadd, 2010; Gall et al., 2015; Wuana et al., 2011).

2.4.2 Plants

In many cases, PTEs are concentrated in the upper part of the soil profile where roots reside, meaning that increased mobility is likely to affect plants growing in floodplain soils. Wetland plants growing on inundated floodplain soils can also affect the mobility of PTEs because they are specially adapted to have air-filled tissues, or aerenchyma, which create patches of oxygenated soil around their roots, resulting in an increase in the volume of the oxic/anoxic interface and remobilising PTEs thus increasing their availability (Du Laing et al., 2009; Wright et al., 2017). However, in arable and pasture fields that are generally drier, flooding can cause crops to become stressed, as they are not adapted to wet soils. As oxygen levels decrease there is a build-up of carbon dioxide, methane and nitrogen gases that leads to the roots suffocating and dying (Hippolyte et al., 2012).

It is well established that symbiotic fungi, associated with plant roots, regulate the supply of micronutrients and reduce the uptake of non-essential PTEs by plants (Classen et al., 2015; Gadd, 2010; Tack, 2010). Plants, such as *Artemisia* and *Phalaris* species, on the floodplain excrete exudates during inundation which stimulates the activity of microbial symbionts in the rhizosphere, allowing PTEs to be taken up into the vegetation (Gall et al., 2015; Sullivan and Gadd, 2019; Violante et al., 2010; Xu et al., 2020). PTEs are often accumulated in plant root tissues and can sometimes be translocated into the plant shoots. However this is regulated in plants by the Casparian strip and therefore limited (Hooda, 2010; Nouri et al., 2009; Shahid et al., 2017). The uptake and accumulation of PTEs is element and plant-specific (Niu et al., 2007; Rinklebe et al., 2016; Tack, 2010; Violante et al., 2010; Xu et al., 2020). The mobilisation and uptake of PTEs by plants may pose a potential environmental risk (Shaheen and Rinklebe, 2014). European floodplains are most commonly used as grassland for grazing cattle or hay production, whereas in other regions e.g. India, they are used for

crops like rice, which raises concerns for possible pollutant transfer from the floodplain soil into the surrounding water bodies, then uptake and potential biomagnification of PTEs into the food chain (Martin et al., 2014; Overesch et al., 2007; Tóth et al., 2016a). However, the hyperaccumulation of PTEs by some plants (e.g. sunflower, mustard (Brassicaceae), alfalfa and Ricinus) has resulted in them being considered for phytoremediation of contaminated floodplain soils (Gall et al., 2015; Niu et al., 2007; Nouri et al., 2009; Shaheen et al., 2016; Violante et al., 2010).

Factors influencing plant uptake of PTEs include soil pH, electrical conductivity and the total concentrations of PTEs in the soil (Nouri et al., 2009). PTEs uptake also depends on the concentrations in the soil solution, governed by plant exudates and root-induced changes to pH and DOM (Gall et al., 2015). Quantifying the total content of PTEs transferred into the food chain via plants growing on contaminated soil is difficult (Gröngöft et al., 2005). The concentrations of PTEs found in floodplain plants are not always directly reflected in the PTEs content found in the soil, due to both physiological and biochemical differences between different plant species; for example differences in the age of the plant biomass (seasonal trends in growth and therefore uptake of nutrients). Moreover, the rooting depth influences metal mobilisation/immobilisation and element specific uptake into the roots which also affects the transfer into the shoots (Chrzan, 2016; Overesch et al., 2007). Thapa et al. (2016) also demonstrated a change in semi-arid Australian floodplain vegetation productivity in response to flooding and drying cycles; flooding brings nutrients which increases net primary productivity. These changes in vegetation productivity could also initiate structural changes in floodplain vegetation communities in natural and semi-natural ecosystems (Overesch et al., 2007).

3. Summary and further research needs

3.1. Summary of current understanding

Floodplain soils downstream of urban catchments contain elevated concentrations of PTEs as a legacy of human activity and these PTEs could potentially be remobilised by future flooding events. A number of processes occur within the soil, ultimately determining PTEs fate. These processes

include: sorption, desorption, complexation, precipitation and dissolution, transport of water and heat, and biological activity. The processes are influenced by the changing conditions that flooding brings particularly with regards to soil moisture content, temperature and redox potential. The mobility of PTEs in flooded soils is closely related to changes in redox potential which, in turn, is altered by flooding. These changes can have direct impacts on the mobility of redox sensitive PTEs (e.g. As and Cr). Furthermore, the reduction of Mn and Fe can cause reductive dissolution of co-precipitated PTEs, but the reduction of sulphate can result in the precipitation of PTEs as insoluble metal sulphides. Which of these processes dominates will depend on the mineralogy of the soil. PTEs precipitated as metal sulphides may oxidise after floodwaters recede and mobilise, accelerated by the pH reduction caused by production of sulfuric acid. There are important interactions between redox potential and other soil properties, such as soil pH, moisture content, POM, DOM, temperature, and salinity which also have a strong impact on PTEs mobility (Vaughan et al., 2009). Many of these reactions are microbially mediated, temperature dependent and the kinetics in real-world scenarios are poorly understood. However, it seems that changes associated with alterations to pH and dissolved organic carbon are relatively fast, while changes to E_H are slower and only become apparent after extended periods of flooding. In many cases, PTEs deposited due to legacy pollution events are concentrated in the upper part of the soil profile, meaning that increased mobility is likely to affect plants growing in floodplain soils and potentially lead to contamination of the surrounding environment, including overlying surface waters.

3.2 Knowledge gaps and recommendations for future research

Floods are dynamic events that expose floodplain soils to water with rapidly changing flows, chemical composition, and sediment load. They can be difficult to predict, due to their different types (e.g. overbanking or groundwater flooding), and the high variation in their magnitude, duration, and frequency of recurrence. Therefore, chemical, physical and biological data from floodplain soils immediately before and immediately after a flooding event are often lacking. However these data

would provide the necessary insights into the factors and processes involved in altering the mobility of PTEs during and after a real flooding event (Barber et al., 2017). The effect of flooding on PTEs mobility can be difficult to predict due to there being several factors (e.g. speciation, release through biological degradation and competitive action of other ions) or interactions between factors (e.g. changes in E_H caused degradation of POM) influencing PTEs mobility (Tack and Verloo, 1995).

Contamination of soil with PTEs receives most attention in highly contaminated urban, industrial, mining and waste disposal sites (Adamo et al., 2014; Resongles et al., 2015; Simmler et al., 2017; Wuana et al., 2011) with relatively little attention given to more ‘typical’ floodplains downstream of catchments with a history of urban and industrial development. Much of the work conducted to date (see Table S1) has been undertaken in Europe, America, Canada, China, Indonesia, Australia and New Zealand. Just over half of the studies cited in Table S1 were undertaken in Germany and Belgium (52%), with a particular research effort around the River Elbe and Wupper River in Germany (Du Laing et al., 2009; Förstner, 2004; Frohne et al., 2011; Overesch et al., 2007; Rennert et al., 2017; Rinklebe et al., 2013; Shaheen et al., 2017). However, research examining the relationship between PTEs mobility and flooding in other parts of the world that are expected to see an increase in the frequency and magnitude of flooding events, for example in Asia, Africa and India, is limited.

A number of factors were identified that contribute to whether the mobility of PTEs will increase or decrease during inundation of a floodplain, which may be interconnected or work in combination to affect PTEs mobility. As a result, different soils with differing mineralogy and thus different biogeochemical and physical properties, will likely respond differently to flooding. Individual studies tend to focus on one floodplain site. However, knowledge based on one river catchment may not be particularly useful for predicting the impacts of flooding at another site with different mineralogy and physical and chemical characteristics. A more fundamental mechanistic understanding is required to inform the development of predictive models. Therefore, more coordinated work encompassing multiple contrasting sites is required to understand the relative importance of key soil

properties (e.g. mineralogy, POM, soil pH, texture; and how these affect derived soil properties such as hydraulic and thermal soil properties) on influencing the impact of flooding on the mobility of PTEs.

Many of the findings in the literature are based on research of soils or sediments in laboratory-based artificial flooding environments (Figure 4), which often involve; homogenisation of samples and removal of plant roots, short-exposure time for soil microorganisms and incubation under controlled conditions, such as temperature (often higher than *in-situ* temperatures) and soil water conditions (often wetting the samples with deionised water which is slightly acidic) (Frohne et al., 2011; Izquierdo et al., 2017; Rinklebe et al., 2010; Weber et al., 2009). This makes extrapolation of laboratory-based findings to field situations difficult (Hooda, 2010). Attempts to model the concentration of PTEs in floodplain pore waters have demonstrated the complexity of predicting how different variables such as soil moisture content and temperature interact and alter mobility (Rennert et al., 2017), with site or catchment-specific information being of great importance to establish and capture spatial differences sufficiently (Schulz-Zunkel et al., 2015). While much research undertaken in controlled conditions in laboratory microcosms is undoubtedly useful because independent replicates can be assigned to treatments without confounding variables (Figure 4), there is a clear research need for on-site experiments on the effect of flooding on PTEs mobility using real-time field-based observations that capture the kinetics of processes before, during, and after a flooding event under ambient temperatures and in geochemically contrasting soils.




Figure 4: Strengths (+) and weaknesses (-) of laboratory-based studies for researching the impact of flooding on mobility of PTEs. Created with BioRender.com.

Acknowledgements

Jessica Ponting is supported by a NERC SCENARIO PhD studentship, with CASE support from the British Geological Survey Universities Funding Initiative.

Supplementary Material

One supplementary table (Table S1) is provided

References

- Abgottspon, F., Bigalke, M., Wilcke, W., 2015. Fast colloidal and dissolved release of trace elements in a carbonatic soil after experimental flooding. *Geoderma* 259–260, 156–163.
<https://doi.org/10.1016/J.GEODERMA.2015.06.005>

848 Acreman, M.C., Riddington, R., Booker, D.J., 2003. Hydrological impacts of floodplain restoration: a
849 case study of the River Cherwell, UK, European Geosciences Union.

850 Adamo, P., Iavazzo, P., Albanese, S., Agrelli, D., De Vivo, B., Lima, A., 2014. Bioavailability and soil-to-
851 plant transfer factors as indicators of potentially toxic element contamination in agricultural
852 soils. *Sci. Total Environ.* 500–501, 11–22. <https://doi.org/10.1016/J.SCITOTENV.2014.08.085>

853 Adekanmbi, A.A., Shaw, L.J., Sizmur, T., 2020. Effect of Sieving on Ex Situ Soil Respiration of Soils
854 from Three Land Use Types. *J. Soil Sci. Plant Nutr.* 1–5. [https://doi.org/10.1007/s42729-020-](https://doi.org/10.1007/s42729-020-00177-2)
855 [00177-2](https://doi.org/10.1007/s42729-020-00177-2)

856 Adewuyi, G.O., Osobamiro, M.T., 2016. Chemical speciation and potential mobility of some toxic
857 metals in tropical agricultural soil. *Res. J. Environ. Toxicol.* 10, 159–165.
858 <https://doi.org/10.3923/rjet.2016.159.165>

859 Alfieri, L., Bisselink, B., Dottori, F., Naumann, G., de Roo, A., Salamon, P., Wyser, K., Feyen, L., 2017.
860 Global projections of river flood risk in a warmer world. *Earth's Futur.* 5, 171–182.
861 <https://doi.org/10.1002/2016EF000485>

862 Alloway, B.J., 2013. Heavy Metals in Soils, Heavy Metals in Soils. [https://doi.org/10.1007/978-94-](https://doi.org/10.1007/978-94-007-4470-7_10)
863 [007-4470-7_10](https://doi.org/10.1007/978-94-007-4470-7_10)

864 Alloway, B.J., 1995. Heavy Metals in Soils: Trace Metals and Metalloids in Soils and their
865 Bioavailability, 3rd ed. Springer, Heidelberg. <https://doi.org/10.1007/978-94-011-1344-1>

866 Álvarez-Ayuso, E., Otones, V., Murciego, A., García-Sánchez, A., Regina, I.S., 2012. Antimony, arsenic
867 and lead distribution in soils and plants of an agricultural area impacted by former mining
868 activities. *Sci. Total Environ.* 439, 35–43. <https://doi.org/10.1016/J.SCITOTENV.2012.09.023>

869 Alvim Ferraz, M.C.M., Lourenço, J.C.N., 2000. Influence of organic matter content of contaminated
870 soils on the leaching rate of heavy metals. *Environ. Prog.* 19, 53–58.
871 <https://doi.org/10.1002/ep.670190118>

872 Amacher, M.C., Kotuby-Amacher, J., Selim, H.M., Iskandar, I.K., 1986. Retention and release of
873 metals by soils — Evaluation of several models. *Geoderma* 38, 131–154.
874 [https://doi.org/10.1016/0016-7061\(86\)90011-X](https://doi.org/10.1016/0016-7061(86)90011-X)

875 Ander, E.L., Watts, M.J., Smedley, P.L., Hamilton, E.M., Close, R., Crabbe, H., Fletcher, T., Rimell, A.,
876 Studden, M., Leonardi, G., 2016. Variability in the chemistry of private drinking water supplies

877 and the impact of domestic treatment systems on water quality. *Environ. Geochem. Health* 38,
878 1313–1332. <https://doi.org/10.1007/s10653-016-9798-0>

879 Andersen, D.C., 2018. Flood effects on soil thermal regimes in contrasting cold-desert river
880 floodplains (Yampa and Green rivers, Colorado). *Ecohydrology* 11, e1939.
881 <https://doi.org/10.1002/eco.1939>

882 Antoniadis, V., Shaheen, S.M., Tsadilas, C.D., Selim, M.H., 2018. Zinc sorption by different soils as
883 affected by selective removal of carbonates and hydrous oxides. *Appl. Geochemistry* 88, 49–58.
884 <https://doi.org/10.1016/J.APGEOCHEM.2017.04.007>

885 Arnell, N.W., Halliday, S.J., Battarbee, R.W., Skeffington, R.A., Wade, A.J., 2015. The implications of
886 climate change for the water environment in England. *Prog. Phys. Geogr. Earth Environ.* 39, 93–
887 120. <https://doi.org/10.1177/0309133314560369>

888 Baran, A., Tarnawski, M., 2015. Assessment of heavy metals mobility and toxicity in contaminated
889 sediments by sequential extraction and a battery of bioassays. *Ecotoxicology* 24, 1279–1293.
890 <https://doi.org/10.1007/s10646-015-1499-4>

891 Barber, L.B., Paschke, S.S., Battaglin, W.A., Douville, C., Fitzgerald, K.C., Keefe, S.H., Roth, D.A., Vajda,
892 A.M., 2017. Effects of an Extreme Flood on Trace Elements in River Water - From Urban Stream
893 to Major River Basin. *Environ. Sci. Technol.* 51, 10344–10356.
894 <https://doi.org/10.1021/acs.est.7b01767>

895 Bednářová, Z., Komprdová, K., Kalábová, T., Sánka, M., 2015. Impact of floods and their frequency on
896 content and distribution of risk elements in alluvial soils. *Water. Air. Soil Pollut.* 226.
897 <https://doi.org/10.1007/s11270-014-2253-x>

898 Bell, V.A., Kay, A.L., Cole, S.J., Jones, R.G., Moore, R.J., Reynard, N.S., 2012. How might climate
899 change affect river flows across the Thames Basin? An area-wide analysis using the UKCP09
900 Regional Climate Model ensemble. *J. Hydrol.* 442–443, 89–104.
901 <https://doi.org/10.1016/J.JHYDROL.2012.04.001>

902 Bellanger, B., Huon, S., Steinmann, P., Velasquez, F., Vall Es E, V., Arn, K., Clauer, N., Mariotti, A.E.,
903 2004. Oxic-anoxic conditions in the water column of a tropical freshwater reservoir (Peña-
904 Larga dam, NW Venezuela), Published in *Applied Geochemistry*.
905 <https://doi.org/10.1016/j.apgeochem.2003.11.007>

906 Berz, G., Kron, W., Loster, T., Rauch, E., Schimetschek, J., Schmieder, J., Siebert, A., Smolka, A., Wirtz,
 907 A., 2001. World Map of Natural Hazards-A Global View of the Distribution and Intensity of
 908 Significant Exposures, Natural Hazards.

909 Blackwell, M.S.A., Pilgrim, E.S., 2011. Ecosystem services delivered by small-scale wetlands. *Hydrol.*
 910 *Sci. J.* 56, 1467–1484. <https://doi.org/10.1080/02626667.2011.630317>

911 Blöschl, G., Hall, J., Parajka, J., Perdigão, R.A.P., Merz, B., Arheimer, B., Aronica, G.T., Bilibashi, A.,
 912 Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G.B., Claps, P., Fiala, K., Frolova, N.,
 913 Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T.R.,
 914 Kohnová, S., Koskela, J.J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L.,
 915 Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Rogger,
 916 M., Salinas, J.L., Sauquet, E., Šraj, M., Szolgay, J., Viglione, A., Volpi, E., Wilson, D., Zaimi, K.,
 917 Živković, N., 2017. Changing climate shifts timing of European floods. *Science* (80-.). 357, 588–
 918 590. <https://doi.org/10.1126/science.aan2506>

919 Borch, T., Kretzschmar, R., Skappler, A., Van Cappellen, P., Ginder-Vogel, M., Voegelin, A., Campbell,
 920 K., 2010. Biogeochemical redox processes and their impact on contaminant dynamics. *Environ.*
 921 *Sci. Technol.* <https://doi.org/10.1021/es9026248>

922 Bradley, S.B., Cox, J.J., 1990. The significance of the floodplain to the cycling of metals in the river
 923 Derwent Catchment, U.K. *Sci. Total Environ.* 97–98, 441–454. [https://doi.org/10.1016/0048-](https://doi.org/10.1016/0048-9697(90)90255-S)
 924 [9697\(90\)90255-S](https://doi.org/10.1016/0048-9697(90)90255-S)

925 Brönnimann, S., 2007. Impact of El Niño-Southern Oscillation on European climate. *Rev. Geophys.*
 926 45, n/a-n/a. <https://doi.org/10.1029/2006RG000199>

927 Bronstert, A., 2003. *Floods and Climate Change: Interactions and Impacts, Risk Analysis.*

928 Bufflap, S.E., Allen, H.E., 1995. Sediment pore water collection methods for trace metal analysis: A
 929 review. *Water Res.* 29, 165–177. [https://doi.org/10.1016/0043-1354\(94\)E0105-F](https://doi.org/10.1016/0043-1354(94)E0105-F)

930 Chan, S.C., Kendon, E.J., Fowler, H.J., Blenkinsop, S., Roberts, N.M., 2014. Projected increase in
 931 summer and winter UK sub-daily precipitation extremes from high-resolution regional climate
 932 models [WWW Document]. *Environ. Res. Lett.* URL
 933 [https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_](https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models)
 934 [winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models](https://www.researchgate.net/publication/265090772_Projected_increase_in_summer_and_winter_UK_sub-daily_precipitation_extremes_from_high-resolution_regional_climate_models)
 935 (accessed 3.2.20).

936 Chang, H., Franczyk, J., 2008. Climate Change, Land-Use Change, and Floods: Toward an Integrated
937 Assessment. *Geogr. Compass* 2, 1549–1579. <https://doi.org/10.1111/j.1749-8198.2008.00136.x>

938 Chrzan, A., 2016. Monitoring bioconcentration of potentially toxic trace elements in soils trophic
939 chains. *Environ. Earth Sci.* 75, 1–8. <https://doi.org/10.1007/s12665-016-5595-4>

940 Cipullo, S., Prpich, G., Campo, P., Coulon, F., 2018. Assessing bioavailability of complex chemical
941 mixtures in contaminated soils: Progress made and research needs. *Sci. Total Environ.* 615,
942 708–723. <https://doi.org/10.1016/j.scitotenv.2017.09.321>

943 Ciszewski, D., Grygar, T.M., 2016. A Review of Flood-Related Storage and Remobilization of Heavy
944 Metal Pollutants in River Systems. *Water. Air. Soil Pollut.* 227, 1–19.
945 <https://doi.org/10.1007/s11270-016-2934-8>

946 Classen, A.T., Sundqvist, M.K., Henning, J.A., Newman, G.S., Moore, J.A.M., Cregger, M.A.,
947 Moorhead, L.C., Patterson, C.M., 2015. Direct and indirect effects of climate change on soil
948 microbial and soil microbial-plant interactions: What lies ahead? *Ecosphere* 6, art130.
949 <https://doi.org/10.1890/ES15-00217.1>

950 Clemente, R., Dickinson, N.M., Lepp, N.W., 2008. Mobility of metals and metalloids in a multi-
951 element contaminated soil 20 years after cessation of the pollution source activity. *Environ.*
952 *Pollut.* 155, 254–261. <https://doi.org/10.1016/j.envpol.2007.11.024>

953 Czech, T., Baran, A., Wieczorek, J., 2014. Zawartość Metali Ciężkich W Glebach I Roślinach Z Terenu
954 Gminy Borzęcin (Województwo Małopolskie) Content of Heavy Metals in Soil and Plants From
955 an Area Borzęcin Municipality (Malopolska Province) 89–98.
956 <https://doi.org/10.12912/2081139X.20>

957 Dadson, S.J., Hall, J.W., Murgatroyd, A., Acreman, M., Bates, P., Beven, K., Heathwaite, L., Holden, J.,
958 Holman, I.P., Lane, S.N., O'connell, E., Penning-Rowsell, E., Reynard, N., Sear, D., Thorne, C.,
959 Wilby, R., 2017. A restatement of the natural science evidence concerning catchment-based
960 “natural” flood management in the UK. *Proc. R. Soc. A* 473.
961 <https://doi.org/10.1098/rspa.2016.0706>

962 Dang, Z., Liu, C., Haigh, M.J., 2002. Mobility of heavy metals associated with the natural weathering
963 of coal mine spoils. *Environ. Pollut.* 118, 419–426. [https://doi.org/10.1016/S0269-](https://doi.org/10.1016/S0269-7491(01)00285-8)
964 [7491\(01\)00285-8](https://doi.org/10.1016/S0269-7491(01)00285-8)

965 Davranche, M., Grybos, M., Gruau, G., Pédrot, M., Dia, A., Marsac, R., 2011. Rare earth element
 966 patterns: A tool for identifying trace metal sources during wetland soil reduction. *Chem. Geol.*
 967 284, 127–137. <https://doi.org/10.1016/j.chemgeo.2011.02.014>

968 Dawson, J.J.C., Tetzlaff, D., Carey, A.M., Raab, A., Soulsby, C., Killham, K., Meharg, A.A., 2010.
 969 Characterizing Pb mobilization from upland soils to streams using $^{206}\text{Pb}/^{207}\text{Pb}$ isotopic ratios.
 970 *Environ. Sci. Technol.* 44, 243–249. <https://doi.org/10.1021/es902664d>

971 De Jonge, M., Teuchies, J., Meire, P., Blust, R., Bervoets, L., 2012. The impact of increased oxygen
 972 conditions on metal-contaminated sediments part I: Effects on redox status, sediment
 973 geochemistry and metal bioavailability. *Water Res.* 46, 2205–2214.
 974 <https://doi.org/10.1016/J.WATRES.2012.01.052>

975 de Souza Machado, A.A., Spencer, K., Kloas, W., Toffolon, M., Zarfl, C., 2016. Metal fate and effects
 976 in estuaries: A review and conceptual model for better understanding of toxicity. *Sci. Total*
 977 *Environ.* 541, 268–281. <https://doi.org/10.1016/J.SCITOTENV.2015.09.045>

978 de Souza Machado, A.A., Spencer, K.L., Zarfl, C., O’Shea, F.T., 2018. Unravelling metal mobility under
 979 complex contaminant signatures. *Sci. Total Environ.* 622–623, 373–384.
 980 <https://doi.org/10.1016/J.SCITOTENV.2017.11.239>

981 De Vivo, B., Belkin, H.E., Lima, A., 2008. *Environmental Geochemistry: Site Characterization, Data*
 982 *Analysis And Case Histories*, *Environmental Geochemistry: Site Characterization, Data Analysis*
 983 *and Case Histories*. Elsevier. <https://doi.org/10.1016/B978-0-444-53159-9.X0001-0>

984 Degryse, F., Smolders, E., Parker, D.R., 2009. Partitioning of metals (Cd, Co, Cu, Ni, Pb, Zn) in soils:
 985 concepts, methodologies, prediction and applications - a review. *Eur. J. Soil Sci.* 60, 590–612.
 986 <https://doi.org/10.1111/j.1365-2389.2009.01142.x>

987 Domergue, F.L., Vedy, J.C., 1992. Mobility of heavy metals in soil profiles. *Int. J. Environ. Anal. Chem.*
 988 46, 13–23. <https://doi.org/10.1080/03067319208026993>

989 Du Laing, G., Rinklebe, J., Vandecasteele, B., Meers, E., Tack, F.M.G., 2009. Trace metal behaviour in
 990 estuarine and riverine floodplain soils and sediments: A review. *Sci. Total Environ.* 407, 3972–
 991 3985. <https://doi.org/10.1016/J.SCITOTENV.2008.07.025>

992 Du Laing, G., Vanthuyne, D.R.J., Vandecasteele, B., Tack, F.M.G., Verloo, M.G., 2007. Influence of
 993 hydrological regime on pore water metal concentrations in a contaminated sediment-derived

994 soil. Environ. Pollut. 147, 615–625. <https://doi.org/10.1016/J.ENVPOL.2006.10.004>

995 Eggleton, J., Thomas, K. V, 2004. A review of factors affecting the release and bioavailability of
 996 contaminants during sediment disturbance events. Environ. Int. 30, 973–980.
 997 <https://doi.org/10.1016/J.ENVINT.2004.03.001>

998 Ehlers, G.A.C., Loibner, A.P., 2006. Linking organic pollutant (bio)availability with geosorbent
 999 properties and biomimetic methodology: A review of geosorbent characterisation and
 1000 (bio)availability prediction. Environ. Pollut. 141, 494–512.
 1001 <https://doi.org/10.1016/J.ENVPOL.2005.08.063>

1002 Ellis, J.B., Shutes, R.B.E., Revitt, M.D., 2003. Constructed Wetlands and Links with Sustainable
 1003 Drainage Systems.

1004 Emerson, Kodak U, Scholastica Bejor, E., Ekeng, E.E., Ogarekpe, Nkpa M, Emerson, K U, Bejor, E.S.,
 1005 Ogarekpe, N M, Onuruka, A.U., 2017. TRANSPORT AND FATE OF SELECTED HEAVY METALS IN
 1006 CIRCUMNEUTRAL RIVER ENVIRONMENT: A CASE STUDY OF THE RIVER NENT CUMBRIA,
 1007 ENGLAND. Int. J. Res. 5, 159–169. <https://doi.org/10.5281/zenodo.818628>

1008 Euripidou, E., Murray, V., 2004. Public health impacts of floods and chemical contamination. J. Public
 1009 Health (Bangkok). 26, 376–383. <https://doi.org/10.1093/pubmed/fdh163>

1010 Evans, L.J., 1989. Chemistry of Metal Retention by Soils Chemistry of metal retention by soils Several
 1011 processes are explained. Artic. Environ. Sci. Technol. <https://doi.org/10.1021/es00067a001>

1012 Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coe, M.T.,
 1013 Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C.,
 1014 Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use.
 1015 Science (80-.). <https://doi.org/10.1126/science.1111772>

1016 Förstner, U., 2004. Traceability of sediment analysis. TrAC Trends Anal. Chem. 23, 217–236.
 1017 [https://doi.org/10.1016/S0165-9936\(04\)00312-7](https://doi.org/10.1016/S0165-9936(04)00312-7)

1018 Forstner, U., Wattman, G., 1981. Metal Pollution in the Aquatic Environment., 2nd ed. Springer-
 1019 Verlag, Berlin.

1020 Foulds, S.A., Griffiths, H.M., Macklin, M.G., Brewer, P.A., 2014. Geomorphological records of extreme
 1021 floods and their relationship to decadal-scale climate change. Geomorphology 216, 193–207.
 1022 <https://doi.org/10.1016/J.GEOMORPH.2014.04.003>

- 1023 Frohne, T., Diaz-Bone, R.A., Du Laing, G., Rinklebe, J., 2015. Impact of systematic change of redox
1024 potential on the leaching of Ba, Cr, Sr, and V from a riverine soil into water. *J. Soils Sediments*
1025 15, 623–633. <https://doi.org/10.1007/s11368-014-1036-8>
- 1026 Frohne, T., Rinklebe, J., Diaz-Bone, R.A., 2014. Contamination of Floodplain Soils along the Wupper
1027 River, Germany, with As, Co, Cu, Ni, Sb, and Zn and the Impact of Pre-definite Redox Variations
1028 on the Mobility of These Elements. *Soil Sediment Contam.* 23, 779–799.
1029 <https://doi.org/10.1080/15320383.2014.872597>
- 1030 Frohne, T., Rinklebe, J., Diaz-Bone, R.A., Du Laing, G., 2011. Controlled variation of redox conditions
1031 in a floodplain soil: Impact on metal mobilization and biomethylation of arsenic and antimony.
1032 *Geoderma* 160, 414–424. <https://doi.org/10.1016/J.GEODERMA.2010.10.012>
- 1033 Gadd, G.M., 2010. Metals, minerals and microbes: Geomicrobiology and bioremediation.
1034 *Microbiology.* <https://doi.org/10.1099/mic.0.037143-0>
- 1035 Gall, Jillian E, Boyd, Robert S, Rajakaruna, Nishanta, Gall, J E, Rajakaruna, N, Boyd, R S, 2015. Transfer
1036 of heavy metals through terrestrial food webs: a review, *Monitoring and Assessment.*
- 1037 Gambrell, R.P., 1994. Trace and Toxic Metals in Wetlands-A Review. *J. Environ. Qual.* 23, 883–891.
- 1038 Gao, H., Shao, M., 2015. Effects of temperature changes on soil hydraulic properties. *Soil Tillage Res.*
1039 153, 145–154. <https://doi.org/10.1016/j.still.2015.05.003>
- 1040 Giacalone, A., Gianguzza, A., Orecchio, S., Piazzese, D., Dongarrà, G., Sciarrino, S., Varrica, D., 2005.
1041 Metals distribution in the organic and inorganic fractions of soil: A case study on soils from
1042 Sicily. *Chem. Speciat. Bioavailab.* 17, 83–93. <https://doi.org/10.3184/095422905782774892>
- 1043 González-Alcaraz, M.N., van Gestel, C.A.M., 2015. Climate change effects on enchytraeid
1044 performance in metal-polluted soils explained from changes in metal bioavailability and
1045 bioaccumulation. *Environ. Res.* 142, 177–184. <https://doi.org/10.1016/j.envres.2015.06.027>
- 1046 González-Macé, O., Steinauer, K., Jousset, A., Eisenhauer, N., Scheu, S., 2016. Flood-induced changes
1047 in soil microbial functions as modified by plant diversity. *PLoS One* 11.
1048 <https://doi.org/10.1371/journal.pone.0166349>
- 1049 Gregory, A.S., Ritz, K., McGrath, S.P., Quinton, J.N., Goulding, K.W.T., Jones, R.J.A., Harris, J.A., Bol,
1050 R., Wallace, P., Pilgrim, E.S., Whitmore, A.P., 2015. A review of the impacts of degradation
1051 threats on soil properties in the UK. *Soil Use Manag.* 31, 1–15.

1052 <https://doi.org/10.1111/sum.12212>

1053 Grimm, A.M., Tedeschi, R.G., 2009. ENSO and extreme rainfall events in South America. *J. Clim.* 22,
1054 1589–1609. <https://doi.org/10.1175/2008JCLI2429.1>

1055 Gröngröft, A., Krüger, F., Grunewald, K., Meißner, R., Miehl, G., 2005. Plant and soil
1056 contamination with trace metals in the Elbe floodplains: A case study after the flood in August
1057 2002. *Acta Hydrochim. Hydrobiol.* 33, 466–474. <https://doi.org/10.1002/aheh.200400596>

1058 Hahne, H.C.H., Kroontje, W., 1973. Significance of pH and Chloride Concentration on Behavior of
1059 Heavy Metal Pollutants: Mercury(II), Cadmium(II), Zinc(II), and Lead(II). *J. Environ. Qual.* 2, 444–
1060 450. <https://doi.org/10.2134/jeq1973.00472425000200040007x>

1061 Harvey, R.J., Chadwick, D.R., Sánchez-Rodríguez, A.R., Jones, D.L., 2019. Agroecosystem resilience in
1062 response to extreme winter flooding. *Agric. Ecosyst. Environ.* 279, 1–13.
1063 <https://doi.org/10.1016/J.AGEE.2019.04.001>

1064 He, Y., Men, B., Yang, X., Li, Y., Xu, H., Wang, D., 2019. Relationship between heavy metals and
1065 dissolved organic matter released from sediment by bioturbation/bioirrigation. *J. Environ. Sci.*
1066 (China) 75, 216–223. <https://doi.org/10.1016/j.jes.2018.03.031>

1067 Hillel, D., 1998. *Introduction to Environmental Soil Physics*, 1st ed. Academic Press.

1068 Hippolyte, I., Jenny, C., Gardes L., Bakry F., Rivallan R., Pomies V., Cubry P., Tomekpe K., Risterucci A.
1069 M., Roux N., Rouard M., Arnaud E., Kolesnikova-Allen M., Perrier X., 2012. Foundation
1070 characteristics of edible *Musa* triploids revealed from allelic distribution of SSR markers. *Ann.*
1071 *Bot.* 109, 937–951. <https://doi.org/10.1093/aob>

1072 Hirabayashi, Y., Kanae, S., 2009. First estimate of the future global population at risk of flooding.
1073 *Hydrol. Res. Lett.* 3, 6–9. <https://doi.org/10.3178/hrl.3.6>

1074 Hirabayashi, Y., Kanae, S., Emori, S., Oki, T., Kimoto, M., 2008. Global projections of changing risks of
1075 floods and droughts in a changing climate. *Hydrol. Sci. J.* 53, 754–772.
1076 <https://doi.org/10.1623/hysj.53.4.754>

1077 Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H.,
1078 Kanae, S., 2013. Global flood risk under climate change. *Nat. Clim. Chang.* 3, 816–821.
1079 <https://doi.org/10.1038/nclimate1911>

1080 Hobbelen, P.H.F., Koolhaas, J.E., van Gestel, C.A.M., 2006. Bioaccumulation of heavy metals in the
 1081 earthworms *Lumbricus rubellus* and *Aporrectodea caliginosa* in relation to total and available
 1082 metal concentrations in field soils. *Environ. Pollut.* 144, 639–646.
 1083 <https://doi.org/10.1016/J.ENVPOL.2006.01.019>

1084 Hooda, P.S., 2010. Trace Elements in Soils, Trace Elements in Soils. John Wiley and Sons.
 1085 <https://doi.org/10.1002/9781444319477>

1086 Huddart, D., Stott, T., Huddart, D., Stott, T., 2020. Climate Change and Adventure Tourism, in:
 1087 Adventure Tourism. Springer International Publishing, pp. 437–469.
 1088 https://doi.org/10.1007/978-3-030-18623-4_13

1089 Hurley, R.R., Rothwell, J.J., Woodward, J.C., 2017. Metal contamination of bed sediments in the
 1090 Irwell and Upper Mersey catchments, northwest England: exploring the legacy of industry and
 1091 urban growth. *J. Soils Sediments* 17, 2648–2665. <https://doi.org/10.1007/s11368-017-1668-6>

1092 Ibragimow, A., Walna, B., Siepak, M., 2013. Effects of Flooding on the Contamination of Floodplain
 1093 Sediments with Available Fractions of Trace Metals (Western Poland), *J. Environ. Stud.*

1094 IPCC, 2012. Managing the Risks of Extreme Events and Disasters to Advance Climate Change
 1095 Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on
 1096 Climate Change. [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.

1097 Izquierdo, M., Tye, A.M., Chenery, S.R., 2017. Using isotope dilution assays to understand speciation
 1098 changes in Cd, Zn, Pb and Fe in a soil model system under simulated flooding conditions.
 1099 *Geoderma* 295, 41–52. <https://doi.org/10.1016/J.GEODERMA.2017.02.006>

1100 Izquierdo, M., Tye, A.M., Chenery, S.R., 2013. Lability, solubility and speciation of Cd, Pb and Zn in
 1101 alluvial soils of the River Trent catchment UK. *Environ. Sci. Process. Impacts* 15, 1844–1858.
 1102 <https://doi.org/10.1039/c3em00370a>

1103 Jenkins, G.J., Murphy, J.M., Sexton, D.M.H., Lowe, J.A., Jones, P., Kilsby, C.G., 2009. UK Climate
 1104 Projection Briefing Report [WWW Document]. Met Off. Hadley Cent. URL
 1105 https://www.researchgate.net/publication/257343385_UK_Climate_Projection_Briefing_Report
 1106 rt (accessed 3.2.20).

1107 Jiao, W., Ouyang, W., Hao, F., Huang, H., Shan, Y., Geng, X., 2014. Combine the soil water assessment
 1108 tool (SWAT) with sediment geochemistry to evaluate diffuse heavy metal loadings at watershed

1109 scale. *J. Hazard. Mater.* 280, 252–259. <https://doi.org/10.1016/J.JHAZMAT.2014.07.081>

1110 Johnson, Z.F., Chikamoto, Y., Wang, S.Y.S., McPhaden, M.J., Mochizuki, T., 2020. Pacific decadal
1111 oscillation remotely forced by the equatorial Pacific and the Atlantic Oceans. *Clim. Dyn.* 55,
1112 789–811. <https://doi.org/10.1007/s00382-020-05295-2>

1113 Johnston, S.G., Burton, E.D., Aaso, T., Tuckerman, G., 2014. Sulfur, iron and carbon cycling following
1114 hydrological restoration of acidic freshwater wetlands. *Chem. Geol.* 371, 9–26.
1115 <https://doi.org/10.1016/j.chemgeo.2014.02.001>

1116 Junk, W.J., Bayley, P.B., Sparks, R.E., 1989. The Flood Pulse Concept in River-Floodplain Systems. *Can.*
1117 *Spec. Publ. Fish. Aquat. Sci.* 106, 110–127. https://doi.org/10.1007/978-3-662-03416-3_4

1118 Kalbitz, K., Wennrich, R., 1998. Mobilization of heavy metals and arsenic in polluted wetland soils
1119 and its dependence on dissolved organic matter. *Sci. Total Environ.* 209, 27–39.
1120 [https://doi.org/10.1016/S0048-9697\(97\)00302-1](https://doi.org/10.1016/S0048-9697(97)00302-1)

1121 Karimian, N., Johnston, S.G., Burton, E.D., 2018. Iron and sulfur cycling in acid sulfate soil wetlands
1122 under dynamic redox conditions: A review. *Chemosphere* 197, 803–816.
1123 <https://doi.org/10.1016/j.chemosphere.2018.01.096>

1124 Karimian, N., Johnston, S.G., Burton, E.D., 2017. Effect of cyclic redox oscillations on water quality in
1125 freshwater acid sulfate soil wetlands. *Sci. Total Environ.*
1126 <https://doi.org/10.1016/j.scitotenv.2016.12.131>

1127 Karl, T.R., Trenberth, K.E., 2003. Modern Global Climate Change. *Science* (80-.). 302, 1719–1723.
1128 <https://doi.org/10.1126/science.1090228>

1129 Kelly, T.J., Hamilton, E., Watts, M.J., Ponting, J., Sizmur, T., 2020. The effect of flooding and drainage
1130 duration on the release of trace elements from floodplain soils. *Environ. Toxicol. Chem.*
1131 etc.4830. <https://doi.org/10.1002/etc.4830>

1132 Kharin, V. V., Zwiers, F.W., Zhang, X., Hegerl, G.C., 2007. Changes in temperature and precipitation
1133 extremes in the IPCC ensemble of global coupled model simulations. *J. Clim.* 20, 1419–1444.
1134 <https://doi.org/10.1175/JCLI4066.1>

1135 Kirk, G., 2004. *The Biogeochemistry of Submerged Soils, The Biogeochemistry of Submerged Soils.*
1136 Wiley & Sons, Ltd: Chichester. <https://doi.org/10.1002/047086303x>

1137 Kiss, T.B.W., Chen, X., Ponting, J., Sizmur, T., Hodson, M.E., (In Review) Dual stresses of flooding and
 1138 agricultural land use reduce earthworm populations more than the individual stressors. *Sci.*
 1139 *Total Environ.* STOTEN-D-20-17138

1140 Koretsky, C.M., Haveman, M., Beuving, L., Cuellar, A., Shattuck, T., Wagner, M., 2007. Spatial
 1141 variation of redox and trace metal geochemistry in a minerotrophic fen. *Biogeochemistry* 86,
 1142 33–62. <https://doi.org/10.1007/s10533-007-9143-x>

1143 Kowalik, C., Kraft, J., Einax, J.W., 2004. The situation of the German Elbe tributaries - Development of
 1144 the loads in the last 10 years. *Acta Hydrochim. Hydrobiol.* 31, 334–345.
 1145 <https://doi.org/10.1002/aheh.200300507>

1146 Kulp, S.A., Strauss, B.H., 2019. New elevation data triple estimates of global vulnerability to sea-level
 1147 rise and coastal flooding. *Nat. Commun.* 10, 1–12. [https://doi.org/10.1038/s41467-019-12808-](https://doi.org/10.1038/s41467-019-12808-z)
 1148 [z](https://doi.org/10.1038/s41467-019-12808-z)

1149 Kundzewicz, Z.W., Kanae, S., Seneviratne, S.I., Handmer, J., Nicholls, N., Peduzzi, P., Mechler, R.,
 1150 Bouwer, L.M., Arnell, N., Mach, K., Muir-Wood, R., Brakenridge, G.R., Kron, W., Benito, G.,
 1151 Honda, Y., Takahashi, K., Sherstyukov, B., 2014. Flood risk and climate change: global and
 1152 regional perspectives. *Hydrol. Sci. J.* 59, 1–28. <https://doi.org/10.1080/02626667.2013.857411>

1153 Lane, S.N., 2017. Natural flood management. *Wiley Interdiscip. Rev. Water* 4, e1211.
 1154 <https://doi.org/10.1002/wat2.1211>

1155 Le Gall, M., Ayrault, S., Evrard, O., Laceby, J.P., Gateuille, D., Lefèvre, I., Mouchel, J.-M., Meybeck, M.,
 1156 2018. Investigating the metal contamination of sediment transported by the 2016 Seine River
 1157 flood (Paris, France). *Environ. Pollut.* 240, 125–139.
 1158 <https://doi.org/10.1016/J.ENVPOL.2018.04.082>

1159 Lee, D.Y., Huang, J.C., Juang, K.W., Tsui, L., 2005. Assessment of phytotoxicity of chromium in
 1160 flooded soils using embedded selective ion exchange resin method. *Plant Soil* 277, 97–105.
 1161 <https://doi.org/10.1007/s11104-005-5997-7>

1162 Lewin, J., Macklin, M.G., 2010. Floodplain catastrophes in the UK Holocene: Messages for managing
 1163 climate change. *Hydrol. Process.* 24, 2900–2911. <https://doi.org/10.1002/hyp.7704>

1164 Lloyd, J.R., 2003. Microbial reduction of metals and radionuclides. *FEMS Microbiol. Rev.* 27, 411–
 1165 425. [https://doi.org/10.1016/S0168-6445\(03\)00044-5](https://doi.org/10.1016/S0168-6445(03)00044-5)

- 1166 Lu, S., Ren, T., Gong, Y., Horton, R., 2007. An Improved Model for Predicting Soil Thermal
1167 Conductivity from Water Content at Room Temperature. *Soil Sci. Soc. Am. J.* 71, 8–14.
1168 <https://doi.org/10.2136/sssaj2006.0041>
- 1169 Lučić, M., Jurina, I., Ščančar, J., Mikac, N., Vdović, N., 2019. Sedimentological and geochemical
1170 characterization of river suspended particulate matter (SPM) sampled by time-integrated mass
1171 flux sampler (TIMS) in the Sava River (Croatia). *J. Soils Sediments* 19, 989–1004.
1172 <https://doi.org/10.1007/s11368-018-2104-2>
- 1173 Ma, M., Du, H., Wang, D., 2019. Mercury methylation by anaerobic microorganisms: A review. *Crit.*
1174 *Rev. Environ. Sci. Technol.* 49, 1893–1936. <https://doi.org/10.1080/10643389.2019.1594517>
- 1175 Ma, R., McBratney, A., Whelan, B., Minasny, B., Short, M., 2011. Comparing temperature correction
1176 models for soil electrical conductivity measurement. *Precis. Agric.*
1177 <https://doi.org/10.1007/s11119-009-9156-7>
- 1178 MacDonald, D., Dixon, A., Newell, A., Hallaways, A., 2012. Groundwater flooding within an urbanised
1179 flood plain. *J. Flood Risk Manag.* 5, 68–80. <https://doi.org/10.1111/j.1753-318X.2011.01127.x>
- 1180 Macklin, M.G., Rumsby, B.T., 2007. Changing Climate and Extreme Floods in the British Uplands, New
1181 Series.
- 1182 Madsen, H., Lawrence, D., Lang, M., Martinkova, M., Kjeldsen, T.R., 2014. Review of trend analysis
1183 and climate change projections of extreme precipitation and floods in Europe. *J. Hydrol.* 519,
1184 3634–3650. <https://doi.org/10.1016/J.JHYDROL.2014.11.003>
- 1185 Maggioni, V., Massari, C., 2018. On the performance of satellite precipitation products in riverine
1186 flood modeling: A review. *J. Hydrol.* 558, 214–224.
1187 <https://doi.org/10.1016/J.JHYDROL.2018.01.039>
- 1188 Malmon, D. V., Dunne, T., Reneau, S.L., 2002. Predicting the Fate of Sediment and Pollutants in River
1189 Floodplains. *Environ. Sci. Technol* 36, 2026–2032. <https://doi.org/10.1021/ES010509+>
- 1190 Malmon, D. V., Reneau, S.L., Dunne, T., 2004. Sediment sorting and transport by flash floods. *J.*
1191 *Geophys. Res. Earth Surf.* 109, n/a–n/a. <https://doi.org/10.1029/2003jf000067>
- 1192 Maluckov, B.S., 2017. The Catalytic Role of Acidithiobacillus ferrooxidans for Metals Extraction from
1193 Mining - Metallurgical Resource. *Biodivers. Int. J.* 1.
1194 <https://doi.org/10.15406/bij.2017.01.00017>

1195 Martin, M., Bonifacio, E., Hossain, K.M.J., Huq, S.M.I., Barberis, E., 2014. Arsenic fixation and
 1196 mobilization in the soils of the Ganges and Meghna floodplains. Impact of pedoenvironmental
 1197 properties. *Geoderma* 228–229, 132–141. <https://doi.org/10.1016/j.geoderma.2013.09.020>

1198 Meegoda, J.N., Martin, L., 2019. In-Situ Determination of Specific Surface Area of Clays. *Geotech*
 1199 *Geol Eng* 37, 465–474. <https://doi.org/https://doi.org/10.1007/s10706-018-0623-7>

1200 Mullan, D., Matthews, T., Vandaele, K., Barr, I.D., Swindles, G.T., Meneely, J., Boardman, J., Murphy,
 1201 C., 2019. Climate impacts on soil erosion and muddy flooding at 1.5 versus 2°C warming. *L.*
 1202 *Degrad. Dev.* 30, 94–108. <https://doi.org/10.1002/ldr.3214>

1203 Naidu, R., Channey, R., McConnell, S., Johnston, N., Semple, K.T., McGrath, S., Dries, V., Nathanail, P.,
 1204 Harmsen, J., Pruszinski, A., MacMillan, J., Palanisami, T., 2015. Towards bioavailability-based
 1205 soil criteria: past, present and future perspectives. *Environ. Sci. Pollut. Res.* 22, 8779–8785.
 1206 <https://doi.org/10.1007/s11356-013-1617-x>

1207 Naidu, R., Semple, K.T., Megharaj, M., Juhasz, A.L., Bolan, N.S., Gupta, S.K., Clothier, B.E., Schulin, R.,
 1208 2008. Bioavailability: Definition, assessment and implications for risk assessment. *Dev. Soil Sci.*
 1209 32, 39–51. [https://doi.org/10.1016/S0166-2481\(07\)32003-5](https://doi.org/10.1016/S0166-2481(07)32003-5)

1210 Neal, C., Smith, C.J., Jeffery, H.A., Jarvie, H.P., Robson, A.J., 1996. Trace element concentrations in
 1211 the major rivers entering the Humber estuary, NE England. *J. Hydrol.* 182, 37–64.
 1212 [https://doi.org/10.1016/0022-1694\(95\)02940-0](https://doi.org/10.1016/0022-1694(95)02940-0)

1213 Ng, J.C., Juhasz, A., Smith, E., Naidu, R., 2015. Assessing the bioavailability and bioaccessibility of
 1214 metals and metalloids. *Environ. Sci. Pollut. Res.* 22, 8802–8825.
 1215 <https://doi.org/10.1007/s11356-013-1820-9>

1216 Niu, Z., Sun, L., Sun, T., Li, Y., Wang, H., 2007. Evaluation of phytoextracting cadmium and lead by
 1217 sunflower, ricinus, alfalfa and mustard in hydroponic culture. *J. Environ. Sci.* 19, 961–967.
 1218 [https://doi.org/10.1016/S1001-0742\(07\)60158-2](https://doi.org/10.1016/S1001-0742(07)60158-2)

1219 Nouri, J., Khorasani, N., Lorestani, B., Karami, M., Hassani, A.H., Yousefi, N., 2009. Accumulation of
 1220 heavy metals in soil and uptake by plant species with phytoremediation potential. *Environ.*
 1221 *Earth Sci.* 59, 315–323. <https://doi.org/10.1007/s12665-009-0028-2>

1222 Nriagu, J.O., Bhattacharya, P., Mukherjee, A.B., Bundschuh, J., Zevenhoven, R., Loeppert, R.H., 2007.
 1223 Arsenic in soil and groundwater: an overview. *Trace Met. other Contam. Environ.* 9, 3–60.

- 1224 [https://doi.org/10.1016/S1875-1121\(06\)09001-8](https://doi.org/10.1016/S1875-1121(06)09001-8)
- 1225 Nshimiyimana, F.X., Faciu, M.-E., Abidi, A. El, Blidi, S. El, Fekhaoui, M., Ifrim, I.L., Soulaymani, A.,
 1226 Lazar, G., 2014. ANALYSIS OF SEASONAL VARIATION ON DEGREE OF CONTAMINATION WITH
 1227 HEAVY METALS IN AARJATE VILLAGE, MOROCCO. AN INDEX APPROACH. Sci. Study Res. Chem.
 1228 Chem. Eng. Biotechnol. Food Ind. 15, 337–344.
- 1229 Ostergren, J.D., Brown, G.E., Parks, G.A., Persson, P., 2000. Inorganic ligand effects on Pb(II) sorption
 1230 to goethite (α -FeOOH). II. Sulfate. J. Colloid Interface Sci. 225, 483–493.
 1231 <https://doi.org/10.1006/jcis.1999.6702>
- 1232 Overesch, M., Rinklebe, J., Broll, G., Neue, H.U., 2007. Metals and arsenic in soils and corresponding
 1233 vegetation at Central Elbe river floodplains (Germany). Environ. Pollut. 145, 800–812.
 1234 <https://doi.org/10.1016/j.envpol.2006.05.016>
- 1235 Pan, L., Fang, G., Wang, Y., Wang, L., Su, B., Li, D., Xiang, B., 2018. Potentially toxic element pollution
 1236 levels and risk assessment of soils and sediments in the upstream river, miyun reservoir, china.
 1237 Int. J. Environ. Res. Public Health 15. <https://doi.org/10.3390/ijerph15112364>
- 1238 Pappenberger, F., Dutra, E., Wetterhall, F., Cloke, H.L., 2012. Deriving global flood hazard maps of
 1239 fluvial floods through a physical model cascade. Hydrol. Earth Syst. Sci. 16, 4143–4156.
 1240 <https://doi.org/10.5194/hess-16-4143-2012>
- 1241 Pathirana, A., Deneke, H.B., Veerbeek, W., Zevenbergen, C., Banda, A.T., 2014. Impact of urban
 1242 growth-driven landuse change on microclimate and extreme precipitation - A sensitivity study.
 1243 Atmos. Res. 138, 59–72. <https://doi.org/10.1016/j.atmosres.2013.10.005>
- 1244 Peacock, M., Freeman, C., Gauci, V., Lebron, I., Evans, C.D., 2015. Investigations of freezing and cold
 1245 storage for the analysis of peatland dissolved organic carbon (DOC) and absorbance properties.
 1246 Environ. Sci. Process. Impacts 17, 1290–1301. <https://doi.org/10.1039/c5em00126a>
- 1247 Peijnenburg, W.J.G.M., Zablotzka, M., Vijver, M.G., 2007. Monitoring metals in terrestrial
 1248 environments within a bioavailability framework and a focus on soil extraction. Ecotoxicol.
 1249 Environ. Saf. 67, 163–179. <https://doi.org/10.1016/J.ECOENV.2007.02.008>
- 1250 Pendergrass, A.G., 2018. What precipitation is extreme? Science (80-.). 360, 1072–1073.
 1251 <https://doi.org/10.1126/science.aat1871>
- 1252 Plum, N.M., Filser, J., 2008. Worms and wetland water: the role of lumbricids and enchytraeids in

- 1253 nutrient mobilization from flooded soils, undefined.
- 1254 Poot, A., Gillissen, F., Koelmans, A.A., 2007. Effects of flow regime and flooding on heavy metal
1255 availability in sediment and soil of a dynamic river system. *Environ. Pollut.* 148, 779–787.
1256 <https://doi.org/10.1016/j.envpol.2007.01.045>
- 1257 Prudhomme, C., Davies, H., 2009. Assessing uncertainties in climate change impact analyses on the
1258 river flow regimes in the UK. Part 2: Future climate. *Clim. Change* 93, 197–222.
1259 <https://doi.org/10.1007/s10584-008-9461-6>
- 1260 Prunty, L., Bell, J., 2005. Soil Temperature Change over Time during Infiltration. *Soil Sci. Soc. Am. J.*
1261 69, 766–775. <https://doi.org/10.2136/sssaj2004.0219>
- 1262 Puchalski, W., 2003. THE IMPORTANCE OF ACCUMULATED ORGANIC MATTER IN NUTRIENT
1263 TRANSPORT AND STORAGE IN FLOODPLAINS, in: International Conference “EcoFlood: Towards
1264 Natural Flood Reduction Strategies”, Warsaw, EU Framework 5.
- 1265 Pulchalski, W., 2003. The Importance of Accumulated Organic Matter in Nutrient Transport and
1266 Storage in Floodplains [WWW Document]. URL
1267 https://www.researchgate.net/publication/237333548_THE_IMPORTANCE_OF_ACCUMULATE
1268 D_ORGANIC_MATTER_IN_NUTRIENT_TRANSPORT_AND_STORAGE_IN_FLOODPLAINS (accessed
1269 2.24.20).
- 1270 Pulley, S., Collins, A.L., 2019. Field-based determination of controls on runoff and fine sediment
1271 generation from lowland grazing livestock fields. *J. Environ. Manage.* 249, 109365.
1272 <https://doi.org/10.1016/j.jenvman.2019.109365>
- 1273 Qiao, P., Lei, M., Yang, S., Yang, J., Zhou, X., Dong, N., Guo, G., 2019. Development of a model to
1274 simulate soil heavy metals lateral migration quantity based on SWAT in Huanjiang watershed,
1275 China. *J. Environ. Sci.* 77, 115–129. <https://doi.org/10.1016/J.JES.2018.06.020>
- 1276 Rawlins, B.G., Wragg, J., Lark, R.M., 2013. Application of a novel method for soil aggregate stability
1277 measurement by laser granulometry with sonication. *Eur. J. Soil Sci.* 64, 92–103.
1278 <https://doi.org/10.1111/ejss.12017>
- 1279 Rennert, T., Rabus, W., Rinklebe, J., 2017. Modelling the concentrations of dissolved contaminants
1280 (Cd, Cu, Ni, Pb, Zn) in floodplain soils. *Environ. Geochem. Health* 39, 331–344.
1281 <https://doi.org/10.1007/s10653-016-9859-4>

- 1282 Resongles, E., Casiot, C., Freydier, R., Le Gall, M., Elbaz-Poulichet, F., 2015. Variation of dissolved and
1283 particulate metal(loid) (As, Cd, Pb, Sb, Tl, Zn) concentrations under varying discharge during a
1284 Mediterranean flood in a former mining watershed, the Gardon River (France). *J. Geochemical*
1285 *Explor.* 158, 132–142. <https://doi.org/10.1016/J.GEXPLO.2015.07.010>
- 1286 Rinklebe, J., Du Laing, G., 2011. Factors Controlling the Dynamics of Trace Metals in Frequently
1287 Flooded Soils, in: *Dynamics and Bioavailability of Heavy Metals in the Rootzone*. CRC Press, pp.
1288 245–270. <https://doi.org/10.1201/b10796-10>
- 1289 Rinklebe, J., During, A., Overesch, M., Du Laing, G., Wennrich, R., Stärk, H.J.H.J., Mothes, S., 2010.
1290 Dynamics of mercury fluxes and their controlling factors in large Hg-polluted floodplain areas.
1291 *Environ. Pollut.* 158, 308–318. <https://doi.org/10.1016/j.envpol.2009.07.001>
- 1292 Rinklebe, J., Franke, C., Neue, H.-U., 2007. Aggregation of floodplain soils based on classification
1293 principles to predict concentrations of nutrients and pollutants. *Geoderma* 141, 210–223.
1294 <https://doi.org/10.1016/J.GEODERMA.2007.06.001>
- 1295 Rinklebe, J., Knox, A.S., Paller, M., Knox, A.S., Paller, M., 2016. Potential Mobility, Bioavailability, and
1296 Plant Uptake of Toxic Elements in Temporary Flooded Soils In book: *Trace elements in*
1297 *waterlogged soils and sediments*, 1st ed. CRC Press, Taylor & Francis Group, New York.
1298 <https://doi.org/10.1201/9781315372952-23>
- 1299 Rinklebe, J., Shaheen, S.M., 2017. Redox chemistry of nickel in soils and sediments: A review.
1300 *Chemosphere* 179, 265–278. <https://doi.org/10.1016/J.CHEMOSPHERE.2017.02.153>
- 1301 Rinklebe, J., Wennrich, R., Du Laing, G., Stark, H.J., Mothes, S., 2013. Mercury emissions from
1302 flooded soils and sediments in Germany are an underestimated problem: challenges for
1303 reliable risk assessments and management strategies. *E3S Web Conf.* 1.
- 1304 Rizzo, A., Bresciani, R., Masi, F., Boano, F., Revelli, R., Ridolfi, L., 2018. Flood reduction as an
1305 ecosystem service of constructed wetlands for combined sewer overflow. *J. Hydrol.* 560, 150–
1306 159. <https://doi.org/10.1016/j.jhydrol.2018.03.020>
- 1307 Rodgers, K., Hursthouse, A., Cuthbert, S., 2015. The Potential of Sequential Extraction in the
1308 Characterisation and Management of Wastes from Steel Processing: A Prospective Review. *Int.*
1309 *J. Environ. Res. Public Health* 12, 11724–11755. <https://doi.org/10.3390/ijerph120911724>
- 1310 Rudiš, M., Valenta, P., Valentová, J., Nöl, O., 2009. Assessment of the deposition of polluted

1311 sediments transferred by a catastrophic flood and related changes in groundwater quality. J.
 1312 Hydrol. 369, 326–335. <https://doi.org/10.1016/J.JHYDROL.2009.02.023>

1313 Sánchez-Rodríguez, A.R., Nie, C., Hill, P.W., Chadwick, D.R., Jones, D.L., 2019. Extreme flood events at
 1314 higher temperatures exacerbate the loss of soil functionality and trace gas emissions in
 1315 grassland. *Soil Biol. Biochem.* 130, 227–236. <https://doi.org/10.1016/J.SOILBIO.2018.12.021>

1316 Schaller, N., Kay, A.L., Lamb, R., Massey, N.R., Van Oldenborgh, G.J., Otto, F.E.L., Sparrow, S.N.,
 1317 Vautard, R., Yiou, P., Ashpole, I., Bowery, A., Crooks, S.M., Haustein, K., Huntingford, C., Ingram,
 1318 W.J., Jones, R.G., Legg, T., Miller, J., Skeggs, J., Wallom, D., Weisheimer, A., Wilson, S., Stott,
 1319 P.A., Allen, M.R., 2016. Human influence on climate in the 2014 southern England winter floods
 1320 and their impacts. *Nat. Clim. Chang.* 6, 627–634. <https://doi.org/10.1038/nclimate2927>

1321 Schulz-Zunkel, C., Krueger, F., 2009. Trace Metal Dynamics in Floodplain Soils of the River Elbe: A
 1322 Review. *J. Environ. Qual.* 38, 1349–1362. <https://doi.org/10.2134/jeq2008.0299>

1323 Schulz-Zunkel, C., Rinklebe, J., Bork, H.-R., 2015. Trace element release patterns from three
 1324 floodplain soils under simulated oxidized–reduced cycles. *Ecol. Eng.* 83, 485–495.
 1325 <https://doi.org/10.1016/J.ECOLENG.2015.05.028>

1326 Selinus, O., Alloway, B., Centeno, J.A., Robert B. Finkelman, Fuge, R., Lindh, U., Smedley, P., 2005.
 1327 Essentials of Medical Geology: Impacts of the Natural Environment on Public Health. Elsevier
 1328 Academic Press.

1329 Shaheen, S.M., Frohne, T., White, J.R., DeLaune, R.D., 2017. Redox-induced mobilization of copper,
 1330 selenium, and zinc in deltaic soils originating from Mississippi (U.S.A.) and Nile (Egypt) River
 1331 Deltas: A better understanding of biogeochemical processes for safe environmental
 1332 management. *J. Environ. Manage.* 186, 131–140.
 1333 <https://doi.org/10.1016/J.JENVMAN.2016.05.032>

1334 Shaheen, S.M., Rinklebe, J., 2017. Sugar beet factory lime affects the mobilization of Cd, Co, Cr, Cu,
 1335 Mo, Ni, Pb, and Zn under dynamic redox conditions in a contaminated floodplain soil. *J.*
 1336 *Environ. Manage.* 186, 253–260. <https://doi.org/10.1016/j.jenvman.2016.07.060>

1337 Shaheen, S.M., Rinklebe, J., 2014. Geochemical fractions of chromium, copper, and zinc and their
 1338 vertical distribution in floodplain soil profiles along the Central Elbe River, Germany. *Geoderma*
 1339 228–229, 142–159. <https://doi.org/10.1016/J.GEODERMA.2013.10.012>

1340 Shaheen, S.M., Rinklebe, J., Frohne, T., White, J.R., DeLaune, R.D., 2016. Redox effects on release
 1341 kinetics of arsenic, cadmium, cobalt, and vanadium in Wax Lake Deltaic freshwater marsh soils.
 1342 Chemosphere 150, 740–748. <https://doi.org/10.1016/J.CHEMOSPHERE.2015.12.043>

1343 Shaheen, S.M., Rinklebe, J., Rupp, H., Meissner, R., 2014a. Temporal dynamics of pore water
 1344 concentrations of Cd, Co, Cu, Ni, and Zn and their controlling factors in a contaminated
 1345 floodplain soil assessed by undisturbed groundwater lysimeters. Environ. Pollut. 191, 223–231.
 1346 <https://doi.org/10.1016/J.ENVPOL.2014.04.035>

1347 Shaheen, S.M., Rinklebe, J., Rupp, H., Meissner, R., 2014b. Lysimeter trials to assess the impact of
 1348 different flood–dry-cycles on the dynamics of pore water concentrations of As, Cr, Mo and V in
 1349 a contaminated floodplain soil. Geoderma 228–229, 5–13.
 1350 <https://doi.org/10.1016/J.GEODERMA.2013.12.030>

1351 Shahid, M., Shamshad, S., Rafiq, M., Khalid, S., Bibi, I., Niazi, N.K., Dumat, C., Rashid, M.I., 2017.
 1352 Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: A
 1353 review. Chemosphere 178, 513–533. <https://doi.org/10.1016/J.CHEMOSPHERE.2017.03.074>

1354 Sherene, T., 2010. Mobility and Transport of Heavy Metals in Polluted Soil Environment. Biol.
 1355 Forum—An Int. J. 2, 112–121.

1356 Simmler, M., Bommer, J., Frischknecht, S., Christl, I., Kotsev, T., Kretzschmar, R., 2017. Reductive
 1357 solubilization of arsenic in a mining-impacted river floodplain: Influence of soil properties and
 1358 temperature. Environ. Pollut. 231, 722–731. <https://doi.org/10.1016/J.ENVPOL.2017.08.054>

1359 Sipos, P., Choi, C., Németh, T., Szalai, Z., Póka, T., 2014. Relationship between iron and trace metal
 1360 fractionation in soils. Chem. Speciat. Bioavailab. 26, 21–30.
 1361 <https://doi.org/10.3184/095422914X13887685052506>

1362 Sivakumar, S., Subbhuraam, C.V., 2005. Toxicity of chromium(III) and chromium(VI) to the
 1363 earthworm *Eisenia fetida*. Ecotoxicol. Environ. Saf. 62, 93–98.
 1364 <https://doi.org/10.1016/J.ECOENV.2004.08.006>

1365 Sizmur, T., Hodson, M.E., 2009. Do earthworms impact metal mobility and availability in soil? – A
 1366 review. Environ. Pollut. 157, 1981–1989. <https://doi.org/10.1016/J.ENVPOL.2009.02.029>

1367 Sizmur, T., Palumbo-Roe, B., Watts, M.J., Hodson, M.E., 2011. Impact of the earthworm *Lumbricus*
 1368 *terrestris* (L.) on As, Cu, Pb and Zn mobility and speciation in contaminated soils. Environ.

- 1369 Pollut. 159, 742–748. <https://doi.org/10.1016/J.ENVPOL.2010.11.033>
- 1370 Sizmur, T., Richardson, J., 2020. Earthworms accelerate the biogeochemical cycling of potentially
1371 toxic elements: Results of a meta-analysis. *Soil Biol. Biochem.* 148, 107865.
1372 <https://doi.org/10.1016/j.soilbio.2020.107865>
- 1373 Spurgeon, D.J., Hopkin, S.P., 1999. Comparisons of metal accumulation and excretion kinetics in
1374 earthworms (*Eisenia fetida*) exposed to contaminated ®eld and laboratory soils. *Appl. Soil Ecol.*
1375 11, 227–243.
- 1376 Srivastava, V., Sarkar, A., Singh, S.K., Singh, P.K., De Araujo, A.S.F., Singh, R.P., 2017. Agroecological
1377 Responses of Heavy Metal Pollution with Special Emphasis on Soil Health and Plant
1378 Performances. *Front. Environ. Sci* 5, 64. <https://doi.org/10.3389/fenvs.2017.00064>
- 1379 Stafford, A., Jeyakumar, P., Hedley, M., Anderson, C., 2018. Influence of Soil Moisture Status on Soil
1380 Cadmium Phytoavailability and Accumulation in Plantain (*Plantago lanceolata*). *Soil Syst.* 2, 9.
1381 <https://doi.org/10.3390/soils2010009>
- 1382 Stagl, J., Mayr, E., Koch, H., Hattermann, F.F., Huang, S., 2014. Effects of Climate Change on the
1383 Hydrological Cycle in Central and Eastern Europe, in: *Advances in Global Change Research*.
1384 Springer International Publishing, pp. 31–43. https://doi.org/10.1007/978-94-007-7960-0_3
- 1385 Stahl, R.G., Hooper, M.J., Balbus, J.M., Clements, W., Fritz, A., Gouin, T., Helm, R., Hickey, C., Landis,
1386 W., Moe, S.J., 2013. The influence of global climate change on the scientific foundations and
1387 applications of Environmental Toxicology and Chemistry: Introduction to a SETAC international
1388 workshop. *Environ. Toxicol. Chem.* 32, 13–19. <https://doi.org/10.1002/etc.2037>
- 1389 Steinnes, E., 2013. Heavy metal contamination of the terrestrial environment from long-range
1390 atmospheric transport: Evidence from 35 years of research in Norway. *E3S Web Conf.* 1, 35001.
1391 <https://doi.org/10.1051/E3SCONF/20130135001>
- 1392 Stuart, M., Lapworth, D., 2011. A review of processes important in the floodplain setting. *Br. Geol.*
1393 *Surv.* 32 (OR/11/030).
- 1394 Stürck, J., Poortinga, A., Verburg, P.H., 2014. Mapping ecosystem services: The supply and demand
1395 of flood regulation services in Europe. *Ecol. Indic.* 38, 198–211.
1396 <https://doi.org/10.1016/j.ecolind.2013.11.010>
- 1397 Sullivan, T.S., Gadd, G.M., 2019. Metal bioavailability and the soil microbiome, in: *Advances in*

- 1398 Agronomy. Academic Press Inc., pp. 79–120. <https://doi.org/10.1016/bs.agron.2019.01.004>
- 1399 Tack, F.M., Verloo, M.G., 1995. Chemical Speciation and Fractionation in Soil and Sediment Heavy
1400 Metal Analysis: A Review, Chemical. International Journal of Environmental Analytical
1401 Chemistry.
- 1402 Tack, F.M.G., 2010. Trace elements: General soil chemistry, principles and processes, in: Trace
1403 Elements in Soils. John Wiley and Sons, pp. 9–37. <https://doi.org/10.1002/9781444319477.ch2>
- 1404 Tack, F.M.G., Van Ranst, E., Lievens, C., Vandenberghe, R.E., 2006. Soil solution Cd, Cu and Zn
1405 concentrations as affected by short-time drying or wetting: The role of hydrous oxides of Fe
1406 and Mn. *Geoderma* 137, 83–89. <https://doi.org/10.1016/j.geoderma.2006.07.003>
- 1407 Tedeschi, R.G., Collins, M., 2016. The influence of ENSO on South American precipitation during
1408 austral summer and autumn in observations and models. *Int. J. Climatol.* 36, 618–635.
1409 <https://doi.org/10.1002/joc.4371>
- 1410 Thapa, R., Thoms, M.C., Parsons, M., Reid, M., 2016. Adaptive cycles of floodplain vegetation
1411 response to flooding and drying. *Earth Surf. Dyn.* 4, 175–191. [https://doi.org/10.5194/esurf-4-](https://doi.org/10.5194/esurf-4-175-2016)
1412 [175-2016](https://doi.org/10.5194/esurf-4-175-2016)
- 1413 Thomas, S.K., Severson, E.D., Galbraith, J.M., 2016. Measuring Saturated Hydraulic Conductivity in
1414 Soil. Virginia Coop. Extension, Blacksburg, VA. 1–13.
- 1415 Tockner, K., Lorang, M.S., Stanford, J.A., 2010. River flood plains are model ecosystems to test
1416 general hydrogeomorphic and ecological concepts. *River Res. Appl.* 26, 76–86.
1417 <https://doi.org/10.1002/rra.1328>
- 1418 Tockner, K., Stanford, J.A., 2002. Riverine flood plains: Present state and future trends. *Environ.*
1419 *Conserv.* 29, 308–330. <https://doi.org/10.1017/S037689290200022X>
- 1420 Tóth, G., Hermann, T., Da Silva, M.R., Montanarella, L., 2016a. Heavy metals in agricultural soils of
1421 the European Union with implications for food safety. *Environ. Int.* 88, 299–309.
1422 <https://doi.org/10.1016/J.ENVINT.2015.12.017>
- 1423 Tóth, G., Hermann, T., Szatmári, G., Pásztor, L., 2016b. Maps of heavy metals in the soils of the
1424 European Union and proposed priority areas for detailed assessment. *Sci. Total Environ.* 565,
1425 1054–1062. <https://doi.org/10.1016/J.SCITOTENV.2016.05.115>

- 1426 Unger, I.M., Kennedy, A.C., Muzika, R.-M., 2009. Flooding effects on soil microbial communities.
1427 Appl. Soil Ecol. 42, 1–8. <https://doi.org/10.1016/J.APSOIL.2009.01.007>
- 1428 Vaughan, K.L., Rabenhorst, M.C., Needelman, B.A., 2009. Saturation and Temperature Effects on the
1429 Development of Reducing Conditions in Soils. Soil Sci. Soc. Am. J. 73, 663–667.
1430 <https://doi.org/10.2136/sssaj2007.0346>
- 1431 Vijver, M.G., Vink, J.P.M., Miermans, C.J.H., van Gestel, C.A.M., 2007. Metal accumulation in
1432 earthworms inhabiting floodplain soils. Environ. Pollut. 148, 132–140.
1433 <https://doi.org/10.1016/j.envpol.2006.10.046>
- 1434 Vink, J.P.M., Meeussen, J.C.L., 2007. BIOCHEM–ORCHESTRA: A tool for evaluating chemical
1435 speciation and ecotoxicological impacts of heavy metals on river flood plain systems. Environ.
1436 Pollut. 148, 833–841. <https://doi.org/10.1016/J.ENVPOL.2007.01.041>
- 1437 Violante, A., 2013. Elucidating Mechanisms of Competitive Sorption at the Mineral/Water Interface.
1438 Adv. Agron. 118, 111–176. <https://doi.org/10.1016/B978-0-12-405942-9.00003-7>
- 1439 Violante, A., Cozzolino, V., Perelomov, L., Caporale, A.G., Pigna, M., 2010. Mobility and bioavailability
1440 of heavy metals and metalloids in soil environments. J. Soil Sci. Plant Nutr. 10, 268–292.
1441 <https://doi.org/10.4067/S0718-95162010000100005>
- 1442 Visser, A., Kroes, J., Van Vliet, M.T.H., Blenkinsop, S., Fowler, H.J., Broers, H.P., 2012. Climate change
1443 impacts on the leaching of a heavy metal contamination in a small lowland catchment, in:
1444 Journal of Contaminant Hydrology. Elsevier, pp. 47–64.
1445 <https://doi.org/10.1016/j.jconhyd.2011.04.007>
- 1446 Weber, F.-A., Voegelin, A., Kretzschmar, R., 2009. Multi-metal contaminant dynamics in temporarily
1447 flooded soil under sulfate limitation. Geochim. Cosmochim. Acta 73, 5513–5527.
1448 <https://doi.org/10.1016/J.GCA.2009.06.011>
- 1449 Wheeler, H., Evans, E., 2009. Land use, water management and future flood risk. Land use policy 26,
1450 S251–S264. <https://doi.org/10.1016/J.LANDUSEPOL.2009.08.019>
- 1451 Wijngaard, R.R., van der Perk, M., van der Grift, B., de Nijs, T.C.M., Bierkens, M.F.P., 2017. The
1452 Impact of Climate Change on Metal Transport in a Lowland Catchment. Water. Air. Soil Pollut.
1453 228, 107. <https://doi.org/10.1007/s11270-017-3261-4>
- 1454 Wilby, R.L., Beven, K.J., Reynard, N.S., 2008. Climate change and fluvial flood risk in the UK: more of

1455 the same? *Hydrol. Process.* 22, 2511–2523. <https://doi.org/10.1002/hyp.6847>

1456 Winger, P.V. V, Lasier, P.J.J., Jackson, B.P.P., 1998. The Influence of Extraction Procedure on Ion
 1457 Concentrations in Sediment Pore Water. *Arch. Environ. Contam. Toxicol.* 35, 8–13.
 1458 <https://doi.org/10.1007/s002449900341>

1459 Wright, A.J., de Kroon, H., Visser, E.J.W., Buchmann, T., Ebeling, A., Eisenhauer, N., Fischer, C.,
 1460 Hildebrandt, A., Ravenek, J., Roscher, C., Weigelt, A., Weisser, W., Voeselek, L.A.C.J., Mommer,
 1461 L., 2017. Plants are less negatively affected by flooding when growing in species-rich plant
 1462 communities. *New Phytol.* 213, 645–656. <https://doi.org/10.1111/nph.14185>

1463 Wuana, R.A., Okieimen, F.E., Montuelle, B., Steinman, A.D., 2011. Heavy Metals in Contaminated
 1464 Soils: A Review of Sources, Chemistry, Risks and Best Available Strategies for Remediation. *Int.*
 1465 *Sch. Res. Netw. ISRN Ecol.* 2011, 20. <https://doi.org/10.5402/2011/402647>

1466 Wyszowska, J., Borowik, A., Kucharski, M., Kucharski, J., 2013. Effect of cadmium, copper and zinc
 1467 on plants, soil microorganisms and soil enzymes. *J. Elem.*
 1468 <https://doi.org/10.5601/jelem.2013.18.4.455>

1469 Xu, J., Zheng, L., Xu, L., Wang, X., 2020. Uptake and allocation of selected metals by dominant
 1470 vegetation in Poyang Lake wetland: From rhizosphere to plant tissues. *Catena* 189, 104477.
 1471 <https://doi.org/10.1016/j.catena.2020.104477>

1472 Xu, J.M., Tang, C., Chen, Z.L., 2006. The role of plant residues in pH change of acid soils differing in
 1473 initial pH. *Soil Biol. Biochem.* 38, 709–719. <https://doi.org/10.1016/j.soilbio.2005.06.022>

1474 Xu, Y., Feng, L., Jeffrey, P.D., Shi, Y., Morel, F.M.M., 2008. Structure and metal exchange in the
 1475 cadmium carbonic anhydrase of marine diatoms. *Nature* 452, 56–61.
 1476 <https://doi.org/10.1038/nature06636>

1477 Yang, C., Li, S., Liu, R., Sun, P., Liu, K., 2015. Effect of reductive dissolution of iron (hydr)oxides on
 1478 arsenic behavior in a water-sediment system: First release, then adsorption. *Ecol. Eng.* 83, 176–
 1479 183. <https://doi.org/10.1016/j.ecoleng.2015.06.018>

1480 Zhao, Y., Marriott, S., Rogers, J., Iwugo, K., 1999. A preliminary study of heavy metal distribution on
 1481 the floodplain of the River Severn, UK by a single flood event. *Sci. Total Environ.* 243–244, 219–
 1482 231. [https://doi.org/10.1016/S0048-9697\(99\)00386-1](https://doi.org/10.1016/S0048-9697(99)00386-1)

1483 Zhao, Y., Marriott, S.B., 2013. Dispersion and Remobilisation of Heavy Metals in the River Severn

1484 System, UK. Procedia Environ. Sci. 18, 167–173.
1485 <https://doi.org/10.1016/J.PROENV.2013.04.022>

1486 Zia, A., van den Berg, L., Ahmad, M.N., Riaz, M., Zia, D., Ashmore, M., 2018. Controls on
1487 accumulation and soil solution partitioning of heavy metals across upland sites in United
1488 Kingdom (UK). J. Environ. Manage. 222, 260–267.
1489 <https://doi.org/10.1016/J.JENVMAN.2018.05.076>

1490

1491

1492

1493